

RECORDS
of the
INDIAN MUSEUM

(A JOURNAL OF INDIAN ZOOLOGY)

Vol. VIII, 1912-1922.

EDITED BY
THE DIRECTOR,
ZOOLOGICAL SURVEY OF INDIA.

Calcutta :

•PUBLISHED BY THE DIRECTOR, ZOOLOGICAL SURVEY OF INDIA.
PRINTED AT THE BAPTIST MISSION PRESS.

1923.

ZOOLOGICAL RESULTS
OF THE
ABOR EXPEDITION.

EDITORIAL NOTE.

We had hoped to have the collections made on the Abor Expedition worked out completely, and to have published a final summary with figures of characteristic animals. The war, however, interrupted the progress of the work, which has continued for 10 years. As there seems little prospect of completing it now and as the delay is causing serious bibliographical inconvenience owing to the lack of an index to the volume, we have decided to bring it to a close. At some future date it may be utilised in a general study of the Himalayan fauna.

N. ANNANDALE,
Director,
Zoological Survey of India.

CALCUTTA :
February 22nd, 1922.

CONTENTS



PART I. *Published October, 1912.*

	<i>Page</i>
Introduction	.. I
I. Batrachia	. 7
II. Reptilia	.. 37
III. Lepidoptera	61
IV. Porifera	67
V. Scolopendridae	69
VI. Hymenoptera Anthophila	.. 75

PART II. *Published March, 1913.*

VII. Mammals	85
VIII. Coleoptera, I-III : Rhysodidae, Nitidulidae, Colydiidae, Cucujidae, Passandridae, Discolomidae, Cryptophagidae, Mycetophagidae and Dryopidae.	99
IX. Arachnida, I : Chelifera, Pedipalpi and Scorpiones	127
X. Dermaptera	135
XI. Diptera	149

PART III. *Published September, 1913.*

XII. Coleoptera, IV-VI : Lamellicornia, Anthribidae and Une nouvelle espèce de Malacodermes des Indes.	191
XIII. Thysanoptera	201
XIV. Arachnida, II : Opiliones	203
XV. Mollusca, I : Rathouisiidae	209
XVI. Temnocephalidae	229
XVII. Hymenoptera, II, III : Ants (Formicidae) and Tenthredinidae	233
XVIII. Fish	243
XIX. Birds	259
XX. Crustacea Decapoda	289
XXI. Orthoptera, I : Tetriginæ (Acridiinae).	311
XXII. Freshwater Planaria	317

PART IV. *Published March, 1914*

XXIII. Hymenoptera, IV : Ichneumonidae	323
XXIV. Orthoptera, II : Gryllacridae and Stenopelmatidae	329
XXV. Odonata	335
XXVI. Neuropteroid Insects	351
XXVII. Reptilia (Supplement)	357
XXVIII. Mollusca, II : Zonitidae and Helicidae (in part)	359
XXIX. Oligochaeta	365

PART V *Published July, 1914.*

XXX. Coleoptera, VII : Ipidæ and Platypodidae	411
XXXI. Orthoptera, III : Mantidae and Phasmidae	415

	<i>Page</i>
XXXII. Termitidae	425
XXXIII. Lepidoptera, II: Moths	437
XXXIV. Hymenoptera, V: Fossores, Diploptera and Chrysi- didae	443
XXXV. Crustacea Amphipoda	449
PART VI. <i>Published December, 1914.</i>	
XXXVI. Land Planarians	455
XXXVII. Terrestrial Isopoda	465
XXXVIII. Onychophora	471
PART VII. <i>Published August, 1915.</i>	
XXXIX. Mollusca, III: Cyclophoridae (in part)	493
XL. Mollusca, IV: Helicidae	505
XLI. Coleoptera, VIII: Passalidae and Lucanidae	515
PART VIII. <i>Published November, 1915.</i>	
XLII. Coleoptera, IX: Tenebrionidae	519
XLIII. Mollusca, V	537
PART IX. <i>Published August, 1916.</i>	
XLIV. Terrestrial Isopoda, II	543
XLV. Mollusca, VI	547
PART X. <i>Published April, 1917.</i>	
XLVI. Collembola	561
XLVII. Mollusca, VII: Cyclophoridae (in part)	569
PART XI. <i>Published February, 1918.</i>	
XLVIII. Mollusca, VIII: Macrochlamyinae (in part)	581
PART XII. <i>Published November, 1918.</i>	
XLIX. Mollusca, IX	601
PART XIII. <i>Published May, 1922.</i>	
L. Coleoptera, X: Hydrophilidae	623

LIST OF PLATES.



	<i>Follow Page</i>
Plate I (Map)	6
Plates II-IV (Batrachia)	36
Plate V (Reptilia)	60
Plate VI (Diptera)	100
Plates VII-IX (Fish)	258
Plates X-XIII (Mollusca)	228
Plate XIV (Temnocephalidae)	232
Plate XV (Orthoptera)	316
Plate XVI (Odonata)	350
Plates XVII-XXI (Crustacea Decapoda)	310
Plate XXII (Freshwater Planaria)	322
Plates XXIII-XXIV (Mollusca)	364
Plate XXV (Neuropteroid insects)	356
Plates XXVI-XXVII (Oligochaeta)	410
Plate XXVIII (Crustacea Amphipoda)	454
Plates XXIX-XXX (Land Planarians)	464
Plates XXXI-XXXIII (Terrestrial Isopoda)	470
Plates XXXIV-XXXVII (Onychophora)	492
Plates XXXVIII-XL (Mollusca)	504
Plates XLI-XLII (Mollusca)	514
Plates XLIII-XLIV (Tenebrionidae)	536
Plates XLV-XLVIII (Mollusca)	560
Plates XLIX-L (Terrestrial Isopoda)	546
Plates LI-LIV (Mollusca)	600
Plates LV-LVII (Collembola)	568

LIST OF AUTHORS.

	<i>Page</i>		<i>Page</i>
Annandale, N., <i>D.Sc., F.A.S.B.</i>		Mollusca, VI: Zonitidae	547
Batrachia	7	Mollusca, VII: Cyclophoridae (in part)	569
Reptilia	37	Mollusca, VIII: Macrochlamyinae (in part)	581
Porifera	67	Mollusca, IX: Zonitidae and Helicidae	601
Reptilia (Supplement)	357	Gravely, F. H., <i>M.Sc.</i>	
Arrow, G. J., <i>F.E.S.</i>		Scolopendridae	69
Coleoptera, IV: Lamellicornia	191	Arachnida, I: Pedipalpi	127
Bagnall, Richard S., <i>F.L.S.</i>		Temnocephalidae	229
Thysanoptera	201	Coleoptera, VIII: Passalidae and Lucanidae	515
Baker, E. C. Stuart.		Coleoptera, IX: Terrebriionidae	519
Birds	259	Griffini, A.	
Banks, Nathan.		Orthoptera, II: Gryllacridae and Stenopelmattidae	329
Neuropteroid Insects	351	Grouvelle, A.	
Bickhardt, H.		Coleoptera, I: Rhyssodidae, Nitidulidae, Colydiidae, Cucujidae, Passandridae, Discolomidae, Cryptophagidae, Mycetophagidae, Dryopidae	99
Coleoptera, III: Histeridae	121	Gude, G. K.	
Brunetti, E.		Mollusca, IV: Helicidae.	505
Diptera	149	Hancock, J. L., <i>F.E.S.</i>	
Burr, Malcolm, <i>D.Sc., F.L.S., etc.</i>		Orthoptera, I: Tetriginae (Acridiinae)	311
Dermaptera	135	Henderson, J. R., <i>M.B., C.M., F.L.S.</i>	
Carpenter, G. H., <i>M.Sc., M.R.I.A.</i>		Arachnida, I: Scorpiones	128
Collembola	561	Jordan, K.	
Chaudhuri, B. L., <i>B.A., B.Sc.</i>		Coleoptera, V: Anthribidae	197
Fish	243	Kemp, S., <i>B.A.</i>	
Collinge, Walter E., <i>M.Sc., F.L.S.</i>		Crustacea Decapoda	289
Terrestrial Isopoda	465	Onychophora	471
" " II	543	Laidlaw, F. F.	
d'Orchymont, A.		Odonata	335
Coleoptera, X: Hydrophilidae	623	Morley, Claude, <i>F.Z.S., E.E.S.</i>	
Elligsen, E.		Hymenoptera, IV: Ichneumonidae	32
Arachnida, I: Chelifera.	127		
Evans, W. H., Capt., <i>R.E.</i>			
Lepidoptera	61		
Ghosh, Ekendranath, <i>B.Sc., L.M.S.</i>			
Mollusca, I: Rathouisidae	209		
Godwin-Austen, H. H., <i>F.R.S.</i>			
Mollusca, II: Zonitidae and Helicidae (in part)	359		
Mollusca, III: Cyclophoridae (in part)	493		

List of Authors.

	<i>Page</i>		<i>Page</i>
Olivier, Ernest.		Roewer, C. Fr.	
Coleoptera, II: Malaco-		Arachnida, II: Opiliones	203
dermidae	119	Rohwer, S. A.	
Paiva, C. A.		Hymenoptera, III: Ten-	
Hymenoptera	Antho-	thredinidae	239
phila	75	Stephenson, J, <i>D.Sc.</i> , <i>Major</i> ,	
Pic, Maurice.		<i>I.M.S.</i>	
Coleoptera, VI: Une nou-		Oligochaeta	365
velle espèce de Malaco-		Wheeler, William Morton.	
dermes des Indes	199	Hymenoptera, II: Ants	
Preston, H. B., <i>F.Z.S.</i>		(Formicidae)	233
Mollusca, V	537	Whitehouse, Richard H.,	
Robinson, Herbert C.,		<i>M.Sc.</i>	
<i>C.M.Z.S.</i>		Freshwater Planaria	317
Mammals	85	Land Planarians	455

INDEX.



[*N.B.*—An asterisk (*) preceding a line denotes a new subspecies or variety; a dagger (†), a new species; a double (‡), a new genus or subgenus; synonyms are printed in italics.]

	<i>Page</i>		<i>Page</i>
A			
Ablabes frenatus ..	37, 47, 53	Agrioninae ..	339, 341
†pavo	37, 47, 53	Agrotis flavirena ..	438
porphyraceus ..	37, 47, 53	Alaopone	234
Abyomma testudinarium	98	conradi ..	234
‡Aborichthys 244	curtisi ..	234
†kempi ..	243, 245	oberthurii ..	234
Abrornis albigularis	271	orientalis fusca	234
superciliaris .	271	orientalis longicornis	234
Acalyptrata	173, 191	vishnui	234
Acanthocephala ..	98	Alcippe nepalensis ..	263
Acantholipes pansalis	438	Alectoropodes ..	287
Acanthopterygii	256	Allactoneura cincta	149
Acanthosaura minor	37, 41, 52	Allocamptus sinuatus	324
Acantholephusa 301	Allodablia ..	145
Acanthothrips sanguineus	.. 201	coriacea .	146
Accipitres 286	scabriuscula	145
Aceraius 516	Allodape ..	83
grandis ..	515, 516	Allograpta ..	163
grandis hirsutus	.. 516	Alphitobius (Diachina) quadri-	
himalayensis .	515, 516	maculatus ..	526
Achorutes 561	Alucurus striatus ..	268
Aciagrion borneensis	344	Alycaei	502, 503
pallidum 344	Alycaeus	494, 502, 569, 570
Acidia fossata 185	aborensis	503
Acisoma panorpoides	337	brahma ..	503
Acorynus carinifrons	.. 197	chanjukensis	503
Actinodura egertoni egertoni	.. 265	distinctus ..	503
Acridiinae ..	311	duoculmen	503
Acromantis 417	graphicus dihiugensis	503
japonica	415, 417	lohitisensis	503
Adiathetus	144	luyorensis	503
glaucopterus	.. 144	oakesi ..	503
Adoretus	191	oglei ..	503
Aegus labilis ..	518	panggiana ..	503
Aetheriidae ..	491	sibbumensis	503
Aethopyga andersoni	259, 281	vesica ..	503
saturata	281	yamneyensis ..	503
seheriae seheriae	259, 281	Alycaeus (Cyclorix) costatus	503
Aeschnidae 339	granum ..	503
Aeschninae ..	339	Alycaeus (Dioryx) globulosus	503
Agamidae 40	globulosus ..	503
Agonosoma	157	urceolus ..	503
Agriocnemis	.. 347	Alycaeus (Raptomphalus) magni-	
†aborense	.. 347	ficus	503
incisa	349	Alyceinae ..	494, 569
lacteola ..	347	Amalopsis glabripennis	.. 153
lunulata ..	347	Amarygmus pilipes ..	532
minima	348	Ambassis rauga	244, 256
†nana	348	Amblycephalidae .	.. 50
†obscura	346	Amblycephalus monticola	37, 50, 54
pygmaea ..	348	mangois ..	243, 244, 252
rubeola ..	347	Ammophila atripes	.. 445
Agrion 343	Amorphoscelis	.. 415
		annulicornis	415

	Page		Page
Amphidromus ..	360	†Aproaspidops ..	45, 54
Amphiops gibbus ..	628, 629	†antecursorum	37, 46, 53
mirabilis ..	623, 628, 629	Apterygota	567
mirabilis variolosus ..	628	Arachnida	97, 98, 127, 203, 466
pedestris ..	628, 629	Arachnothera magna	281
*pedestris varians ..	629	Arachnotherinae ..	281
Amphipoda	449, 453	Arboricola rufigularis intermedia	287
Amphithemis vacillans	336	rufigularis rufigularis	287
Ampittia maro	65	Archibasis oscillans	343
Ampulex	443	<i>Archisometrus mucronatus</i>	128
†aborensis ..	445	<i>scutillus</i> ..	129
Amyna selenampha ..	438	Argynnis hyperbius	63
†Amystrops montana	103	Arhopala silhetensis	65
Anatidae ..	288	Artaminae ..	273
Anax guttatus	339	Artamus fuscus ..	273
Ancaria fuscinervis	327	Artemisia ..	171
Andrena mollis ..	75	Asarkina aegrotus ..	164
Anchurinae ..	145	salviae ..	164
Aneleus minimus	236	Asilidae	157
pygmaeus	236	Asilinae ..	157
sarasinorum	236	Asilus ..	157
similis ..	236	†Assamia aborensis	204
Angitia	325	sexdentata	204
Anguidae ..	42	Assamiidae	204
Anisodactyli	284	Atopos ..	227
Anisolabis	137	†(Podangia) kempii	209
annulipes	138	sanguiolenta	216, 226
gaudens	138	Atteva niveigutta ..	441
pervicina ..	137	subaurata	441
Anisoptera ..	354	Atyidae ..	305, 491
Ankylopteryx octopunctata ..	354	Austenia 360, 548, 549, 550, 554, 556	
Anomala ..	191	†aborensis ..	548
Anomalon ? binghami	325	†alba	549
decorum ..	325	gigas ..	361
Anomoea ..	185	resplendens	360, 361
Anseres	288	rotunda ..	551
Anthomyidae ..	173	†rotungensis	360
Anthomyiinae ..	171	†siyomensis ..	550
<i>Anthophora subcaerulea</i>	78	tigris	548
zonata ..	78		
Anthracias fairmairei	529, 530	B	
†punctipennis	529	Baccha flavopunctata ..	165
tenuis ..	530	gratiosa ..	165
Anthribidae ..	197	<i>Bacillus artemis</i>	419, 420
Anthribus macrocerus ..	198	<i>Bacillus (Baculum) artemis</i>	419, 420
Anthus trivialis ..	281	<i>Bacteria baucis</i> ..	422
trivialis maculatus	280, 281	Badis badis	244, 256
richardi richardi	281	Bactis ..	354
Anurida ..	562	†Bapuia ..	589
Apachyidae	140	†rengingensis ..	589
Apachyus ..	140	Barbus chagunio ..	250, 251
feae	140	chola	243, 244, 249
Aphodiinae ..	195	hexagonolepis	243, 244, 249
Aphodius chinensis	195	hexastichus	243, 244, 249
elegans	195	kunta ..	250
urostigma ..	193	savana ..	244, 250
Apis dorsata	80	sophore	243, 244, 250
latipes	79	spilopholus	244, 250
testacea	80	ticto	244, 251
zonata	78, 80	tor	244, 251
Apobletes macilentus	122	Barytelphusa	302
planidorsum	122	Basitropis hamata	198
tener ..	122	Bathia ..	593
Apodes ..	255	madagascariensis ..	593
Apogonia	192	praetumida	593
Apophua carinata	323	Batrachia	7, 29, 35, 36, 37, 54, 55
Appias lalage ..	63		

	<i>Page</i>		<i>Page</i>
Belone cancila ..	244, 256	Camponotus nicobarensis ..	237
Bensonia 596	taylori albosparsus ..	237
†aborensis ..	596, 600, 621	Campoplex tyrannus ..	325
Berotha insolita 354	Camptolynx fuscipennis ..	327
Bibio obscuripennis 150	striatus ..	327
Bibionidae 150	†Camptoprosopella notatifrons ..	181
Bipalium 455	Capitonidae ..	283
ceres 462	Capo-peripatus ..	483
delicatum ..	458, 459	Capricornis sumatrensis jamrachi ..	91
diana 462	sumatraensis milne-edwardsi ..	92
†dihangense ..	455, 457	sumatraensis thar ..	92
†giganteum ..	456, 457	Capritermes incola ..	435
proserpina 462	†laetus ..	434, 435
robiginosum ..	459, 460	nemorosi ..	435
†rotungense 459	Caracolla ..	537
smithi 457	Cardanus ..	518
†sordidum ..	459, 460	Caridina 289, 290, 305, 306, 308, 310 ..	310
†Bithynia dibrugarhensis ..	540, 541	acutirostris ..	308
pulchella 540	†excavata 289, 290, 306, 308, 310 ..	310
Bithynis (Parapalaemon) hendersoni ..	303	grandirostris ..	310
Blattoidea 3	†hodgarti 289, 290, 308, 309, 310 ..	310
Blennocampa 239	nilotica ..	307
†gracilicornis 239	weberi 231, 289, 290, 305, 306, 308 ..	308
tenuicornis 239	weberi sumatrensis ..	305, 306
Blythia 45, 54	†Caridinicola ..	231, 232, 290
reticulata ..	37, 45, 46, 53	indica ..	231
Boarmia ochrifasciata 440	Carpophaginae ..	286
Bombus buccinatoris 80	Carpophilus dimidiatus ..	100
haemorrhoidalis 80	dimidiatus contingens ..	100
orientalis 80	obsoletus ..	100
simulus 80	Castalius elna ..	64
Barborinae 173	rosimon ..	64
Brachydiplax sobrina 337	Catapiestus indicus ..	528
Brachylabinae 139	Catharsius molossus ..	194
Brachypodinae 267	Catochrysops strabo ..	64
Brachypteryginae 264	Celyphinae ..	184
Brachypteryx 278	Celyphus ? lucidus ..	184
Brachythemis contaminata 338	scutatus ..	184
Brackenridgia cavernarum 467	Centropus bengalensis ..	286
Bringa remifer 270	sinensis ..	285
Bufo 34	Ceratina flavopicta ..	76
himalayanus 17, 19, 29, 34, 35 ..	17, 19, 29, 34, 35	hieroglyphica ..	76
melanostictus .. 7, 19, 20, 34, 35 ..	7, 19, 20, 34, 35	Ceriagrion cerinorubellum ..	346
melanostictus himalayanus ..	19	coromandelianum ..	345, 346
penangensis 32	erubescens ..	346
Bufoidea 19	glabrum ..	346
†Burmoniscus ..	465, 466, 467	melanurum ..	346
†moulmeinensis 466	†olivaceum ..	345, 346
Buteo desertorum 286	Ceropalidae ..	444
Byrsax excisicollis 522	Ceropria induta ..	525
†tuberculatus 522	laticollis ..	525
		subocellata ..	525
		Certhia discolor ..	270
C		Certhiidae ..	270
Caecilius ceylonicus 351	Cervulus muntjac ..	92
Callophis maclellandi 37, 50, 54, 357 ..	37, 50, 54, 357	Cethosia biblis ..	63
Calobata 186	cyane ..	63
†trifascipennis ..	186	Cetoniinae ..	191
Calopsilia florella 64	Chaerilus tricostatus ..	131
Calopteryginae ..	335, 340	Chaetospania feae ..	142
Calotes jerdoni ..	37, 42, 52	Chaptia aenea ..	269
versicolor ..	37, 41, 51	Charadriidae ..	287
Calvisia 423	Chelifer ..	127
fusco-alata ..	415, 423	javanus ..	127
Calyptopogon albitarsis 150	superbus ..	127
Calyptrata 170	Chelisoche ..	143
Camponotinae 237	morio ..	143

	<i>Page</i>		<i>Page</i>
†Chelisoches tigris	143	Columbae	280
Chelisochoidae ..	143	Colydiidae	99, 104
Chelonia	37, 38, 54	Conosia irrorata	153
Cheritra freja ..	65	Copera vittata	343
Chibia hottentotta	269	Copera (Psilocneuis) annulata	342
Chilioperipatus ..	484, 488, 489	*Copera (Psilocnemis) annulata	
Chimarrhornis leucocephalus	276	stevensi ..	341
Chirixalus	33, 34	*Copera (Psilocnemis) vittata	
doriae	7, 18, 34	assamensis	342, 343
Chironomidae ..	150	vittata atomaria	343
Chironomus ..	151	imbricata	343
Chirotonetes	354	serapica	.. 343
Chloritis gabata merguensis	537	vittata	.. 343
Chloroclystes admixtaria	440	Coprinae	193
Chlorogomphinae	335	Copris magicus	194
Choloroperla (Isoteryx)	353	Coptotettix	313
†Chlorops maculipleura ..	188	indicus	311, 313
Chloropsis hardwickii	267	inflatus	313
Chrysididae	443, 447	†nullipennis	311, 313, 314
Chrysis durga	447	parvulus	311, 313, 314
Chrysoconops ..	152	†tumidus	311, 313
Chrysopa alcestris	355	Corbitidinae	.. 244
virgestes ..	355	Cordulegasterinae	.. 335
Chrysophlegma flavinucha	282	Corduliinae	339
Chrysopidae	354	Cordax	147
Chrysops? designata	157	forcipatus	147
Chrysoloxum sexfasciatum	169	Corrodentia	351
Cichlidae	491	Corvidae	260
Cidaria catenaria	440	Corvinae	260
Cirrhina reba	244, 249	Cotile riparia	280
Cirrhochroa aoris	63	sinensis	280
Cissa chinensis ..	260	†Cotyloplana pilleata	463
Cittocinclia macrura macrura ..	278	Crabro	446
Cladotonus eggerti	414	Crateropodidae	.. 262
Clarias assamensis	244, 252	Crateropodinae	.. 262
Clausilia	360, 601	Cremastocephalus ..	564, 565
†aborensis	618, 619	Cremastogaster biroi aitkeni	236
annandalei	618, 619	rogenhoferi	236
insignis	614, 618	Criniger	.. 268
ios	618	flaveolus burmanicus	267
†shimangensis	619	flaveolus flaveolus	267
Clausilia (Oospira) philippiana	539	Criotettix	.311, 312
†Clausilia (Phaedusa) annandalei	538	extremus	.. 311
insignis ..	538	grandis	.. 312
Clausiliidae	538	†maximus	311
Cleon	354	†orientalis	.. 311, 312
Clitumnus ..	417, 420	tricarinatus ..	312
ablutus	415, 417	Crocidura fuliginosa	87
†aboricus	415, 417	rubricosa ..	87
artemis	420	Crocisa decora	77
operculatus	415	elegans	.. 77
porrectus ..	420	emarginata ..	77
†Cuemocryptus epistomatus	326	Crocodilus palustris	38
Coelicia ..	339	Crocothemus servilia	.. 337
†bimaculata	341	Crustacea	4
Coelostoma horni	625	Crustacea Amphipoda	449
orbiculare ..	625	Crustacea Decapoda	289
stultum ..	625	Cryptaustenia	.360, 548
Coenosia	.. 173	†bicolor	.. 548
Coleoptera	16, 99, 119, 121, 191, 197, 199, 411, 515, 519, 615, 623	Cryptogirasia	.558, 559
Colias edusina	64	rubra	559
fieldii	64	Cryptogirasiae	558
Collembola	561, 567, 563	Cryptolopha affinis	.. 270
Coluber taeniurus ..	37, 48, 58	jerdoni	259
Colubridae	45, 286	xanthoschista jerdoni	270, 271
Colubrinae	45	xanthoschista xanthoschista	270
		Cryptoperla	354

	Page		Page
† <i>Cryptoperla divergens</i>	353	<i>Delias aglaia</i>	.. 63
Cryptophagidae ..	99, 112	<i>Dendrocitta frontalis</i>	.. 261
Cryptopinae	69	<i>Dendrocopus macii atrotus</i>	.. 283
Cryptops	69	<i>macii macii</i>	.. 283
<i>doriae</i>	.. 69	<i>Dendrophis gorei</i>	.. 37, 48, 53
Cryptus	328	<i>pictus</i>	.. 48
<i>filicornis</i>	.. 328	<i>Dercas virhuelfi doubledayi</i>	.. 63
<i>himalayensis</i>	.. 328	Dermaptera	3, 35
Cubaris	.. 465, 543	<i>Derosphaerus</i> (?) <i>Encyalesthus</i>	
† <i>caeruleus</i>	.. 467	<i>impresus</i>	.. 528
† <i>fragilis</i>	.. 468	† <i>Derosphaerus rugosus</i>	.. 528
† <i>marmoratus</i>	.. 545	<i>Desmogaster</i>	.. 366, 367
† <i>robusta</i>	.. 468	Diaparsis	326
Cucujidae 99, 107	<i>caudata</i>	.. 326
Cucujus	.. 107	† <i>Diaphanes simulator</i>	.. 119
† <i>kempi</i>	.. 107	<i>Diastramma unicolor</i>	.. 329
Cuculidae 285	<i>Dichraeosis bacillus</i>	.. 521
<i>Culicapa ceylonensis</i>	.. 274	<i>capucinus</i>	.. 520
Culicidae	.. 152	<i>Dicronomyia marmoripennis</i>	.. 152
<i>Cuniculina</i>	.. 418, 420	<i>saltans</i>	.. 152
† <i>perfida</i>	.. 415, 419, 420	Dicruridae 263
† <i>trotungina</i>	.. 415, 418	† <i>Dietyus filicornis</i>	.. 532, 533, 534
<i>detrectans</i>	.. 419	† <i>latifrons</i>	.. 534
Curvella	.. 615	† <i>nodicornis</i>	.. 533, 534
<i>sikkimensis</i>	.. 615	‡ <i>Dihangia</i>	.. 552
<i>sulcata</i>	.. 615	† <i>koboensis</i>	.. 552, 553
<i>Cutia nepalensis</i>	.. 266	<i>Dilemera arctata</i>	.. 437
<i>Cyaniris dilecta</i>	.. 64	<i>Diectes apostata</i>	.. 325
<i>limbata placida</i>	.. 64	Diopsinae	.. 184
<i>transpecta</i>	.. 64	<i>Dipsis indica</i>	.. 184
Cyanops 259	<i>Dioryx</i>	.. 494, 503, 569, 570
<i>asiatica asiatica</i>	.. 283	† <i>Diphyllus molestus</i>	.. 114
<i>asiatica rubescens</i>	.. 284	† <i>odiosus</i>	.. 112
<i>franklini franklini</i>	.. 284	<i>Diplacodes nebulosa</i>	.. 337
<i>franklini ramsayi</i>	.. 284	<i>trivialis</i>	.. 337
Cyclophoridae	.. 493, 540, 569	<i>Diplatyinae</i>	.. 135
Cyclophorinae 493, 569, 570	<i>Diplatys</i>	.. 135
Cyclophorus	.. 493, 569, 577, 585	<i>Diplommatina</i>	.. 494, 569
† <i>aborensis</i>	.. 494	† <i>acutulus</i>	.. 576, 577
† <i>oakesi</i>	.. 502	<i>exilis</i>	.. 577
† <i>Cyclophorus</i> (<i>Glossostylus</i>)		† <i>miriensis</i>	.. 576, 577
<i>koboensis</i>	.. 495	† <i>mucronatus</i>	.. 576
†(<i>Glossostylus</i>) <i>bapuensis</i>	.. 494	† <i>perobesa</i>	.. 537, 540, 541, 576
†(<i>Glossostylus</i>) <i>sidiensis</i>	.. 494, 495	<i>Diplommatininae</i>	.. 494, 569, 576
<i>Cyclopodia horsfieldii</i>	.. 97, 190	Diploptera 443, 446
Cycloryx 494, 503, 569	<i>Dipsadomorphinae</i>	.. 49
Cyclostomacea 575, 578	<i>Dipsadomorphus gokool</i>	.. 37, 49, 51
<i>Cynthia erota</i>	.. 63	<i>Dipsas gokool</i>	.. 49
<i>Cyornis hyperythrus</i>	.. 274	Diptera	.. 149
<i>Cyphoderini</i>	.. 563, 566	<i>Diptera Pupipara</i>	.. 97
‡ <i>Cyphoderopsis</i>	.. 566	<i>Discognathus</i>	.. 247
† <i>kempi</i>	.. 566, 567	<i>lamta</i>	.. 32, 243, 247
<i>Cyphoderus</i>	.. 561, 566	<i>lamta nasutus</i>	.. 247
<i>simulans</i>	.. 563, 566	<i>Discolomidae</i>	.. 99, 111
<i>Cyprinidae</i>	.. 32, 244, 250	<i>Dolichoderinae</i>	.. 237
<i>Cyprininae</i>	.. 247	<i>Dolichopidae</i>	.. 157
<i>Cyrestis thyodamas</i>	.. 62	<i>Dorylinae</i>	.. 233
		<i>Dorylus</i> (<i>Alaopone</i>) <i>orientalis</i>	.. 233, 234
		†(<i>Alaopone</i>) <i>vishnui</i>	.. 233
		<i>Draco</i>	.. 41, 54
		<i>blanfordii</i>	.. 41
		<i>maculatus</i>	.. 37, 40, 41, 52
		<i>norvillii</i>	.. 41
		<i>Drawida</i>	.. 365, 366, 367
		† <i>decourcyi</i>	.. 368, 373
		<i>grandis</i>	.. 375
		† <i>kempi</i>	.. 366, 368, 376

D

	<i>Page</i>		<i>Page</i>
Drawida pellucida	366, 368, 369, 372, 373	Epierus crenulatus	124
pellucida bournei	.. 369	foederatus	124
*pellucida stewarti	368, 369, 372	monticola	124, 125
†rotungana	368, 372	nemoralis	.. 124
Dremomys 90	planiceps	.. 124
lokriah	.. 90, 96	punctulipennis	124
pernyi	.. 85, 90, 96	purus	.. 124
†Drepanephora multimaculata	180	rufescens	.. 124
Drosophila	190	rusticus	.. 124
Drymochares	278	simplex	124
Dryonastes ruficollis	262	sylvanus	124
Dryophis prasinus	37, 50, 54	Epimys rattus	.. 90, 96
†Dryopidius montanus	116	†Epiphragma kempfi	155
Dryopidae	.. 99, 115	†Epipsocus conspersus	351
†Dryops monticola	115	costalis	352
Ducula insignis griseicapilla	.. 287	Epirhyssa flavobalteata	323
insignis insignis	286	Eristalis	169
Durgella	360, 555	Ethmostigmus	72
†aborense	.. 555	pygomegas	72
assamica	555	Euborellia	137
christianoe	555	†aborensis	137
dekhanensis	.. 555	Eucorymus crassicornis	198
levicula	556	Eucyrtus splendens	532
rogersi	.. 555	Eudermaptera	140
Durgellinae	360, 555, 559	Eulota	602, 613
Dynastinae	192	Eulotella similis	602
Dysantes elongatus	523	Eumelia rosalia	440
Dytiscidae	4	Eumenes arcuata	446
		Eumenidae	446
		Euparatettix	315
E		parvus	311, 315
Ecacanthothripidae	201	tenuis	.. 311, 315
Ecacanthothrips	.. 201	Eupatorus hardwickei cantori	.. 192
sanguineus	201	Euploea diocletianus	61
Echinorhynchus gigas	98	†Eupomatias	569, 579
Echinosoma	.. 136	†oakesi	579, 580
sumatranum	136	†sibbumensis	579
Echinomatinae	136	Euponera (Brachyponera) nigrita	233
Elapinae	.. 50	Eurypus	592
Elaunon	146	caseus	592
bipartitus	146	similis	592
Elis annulata 444	Eurytrachelus	517
iris 444	Eutermes ceylonicus	431
Empriinae 239	fletcheri	.. 431
Emydosauria	.. 37, 38	†moratus	431
Encyalesthus exularis	527, 528	†roboratus	430
†stevensi	.. 527	Eutochia lateralis	.. 526
Endomychidae	.. 3	Eutyphoeus	365, 366
Endoplon	.. 512	†aborianus	368, 406
Entomobryidae	561, 563	†kempfi	.. 368, 401
Entomobryinae	563	†koboensis	368, 404, 406, 407
Entomobryini	563, 566	magnus	368, 408
Entozoa 98	Exostoma davidi	243, 244, 252
Eoperipatus	471, 475, 476, 477, 478, 479, 482, 483, 484, 485, 486, 487, 489, 490, 491, 492	labiatum	243, 244, 252
horsti	.. 478		
weldoni	.. 484	F	
Eparchus	.. 146	Falconidae	.. 286
insignis	146	Falconinae	.. 286
Epedaninae	206	Fenienias albomaculatus	.. 327
Ephemera	354	erythropus	.. 327
Ephemeridae	351, 354	Ferussacidae	539
Ephestia cantella	440	Forcipula	139
Epierus	.. 124	pugnax	139
abrogatus 124	Forficulidae	145
amandus	124	Forficulinae	.. 145
		†Formicaleon bivittatum	355

	<i>Page</i>		<i>Page</i>
Formicidae	233	Grammatoptila striata austeni	262, 263
Fossores	443	Grandala	278
Fringillidae	279	Graptomyza brevirostris	167
Fringillinae	279	ventralis	167
Friona curvica rinata	327	*ventralis nigripes	167
†didymata	328	Graucalus layardi	272
rufipes	328	macii macii	272
variipes	328	Gryllacridae	329
Fruticola	606	Gryllacris genuali	330
		†gravelyi	329
G		Gymnodactylus himalayicus	.. 36
Gagrella feae	203	khasiensis	37, 39, 51
feae-humeralis	203	Gymnopleurus assamensis	193
lepida	203	sinuatus	193
Gagrellinae	203	Gynacantha	340
Gallinae	287	hyalina	340
Gallus ferrugineus	287	khasiaca	339, 340
‡Galongia	553	khasiaca nigripes	340
†kempi	553		
Gammaridea	453	H	
Garrulax leucolophus belangeri	262	Habropoda fulvipes	78
leucolophus leucolophus	262	Haemaphysalis	98
moniliger	262	bispinosa	98
Gastropoda	537	gigas	98
Gavialis gangeticus	37, 38, 51	koenigsberneri	97
Gecarcinucinae	291, 302	Haematospiza indica	279
Gecinulus grantia	282	Halictus	75, 83
Gecinus chlorolophus	282	funebri	75
Geckonidae	39	pulchriventris	75
Gelsana	543	rufo-zonatus	75
Geoemyda	54	Hamaxas	144
Geoplanidae	462	dohertyi	145
Geotelphusa	230, 290, 298, 301	feae	145
Geranomyia notatipennis	152	†kempi	144
Girasia 360, 550, 551, 554, 556, 557, 558		nigrorufa	145
cacharica	551, 552	semiluteus	145
†gladstonei	552	Hapalus	601, 615
hookeri	551, 553, 558	Haptoncus ocellaris	100
†maculosa	551	tetragonus	100
rubra	558	Harpactes	260
rubrum	558	erythrocephalus	285
Girasia (Ibycus) sikimensis	552	Hebomoia glaucippe	64
Glossula	359, 360, 585, 601	Hectarthrum trigeminum	111
†aborensis	618	Hedycryptus filicornis	328
botellus	539, 616, 617, 618	Helicarion gigas	558
†oakesi	615, 616, 617	helenae	592
ochracea	616, 617	idae	592
orophila	616	Helecarioninae	360, 548, 558
Globitelphusa	291	Helicidae	359, 360, 493, 505, 537, 569, 602, 604
†Globitermes audax	432, 433	<i>Helix</i>	537, 613
sulphureus	434	catostoma	601, 612, 613, 614
Glossostylus	569	damsangensis	610, 614
Glycyphana minima	191	huttoni	603, 604, 605, 606, 608, 613
nepalensis	191	huttoni radleyi	609
Glyphodes unionalis	441	merguiensis	537
Glyptosternum	32	orbicula	605, 606
Gnapholoryx velutinus	518	(Plectotropis) huttoni	607
Gomphinae	335, 340	Helochares	627
Gonocephalum depressum	520	Helochares (Hydrobaticus) lentus	626
pubeus	520	†kempi	626
ruficornis	520	minutissimus	627, 628
subspinosum	520	Hemerobiidae	354
Gonypeta irina	416	Hemicordulia asiatica	339
Goryphus fuscineris	327	Hemidactylus bowringii	37, 39, 51
maculiceps	327	brookii	37, 39, 40
Grammatoptila striata	262		

	Page		Page
Hemidactylus frenatus	37, 39, 51, 57	Hypoctonus dawnae	.. 128
<i>gleadowii</i>	39	wood-masoni	128
nepalensis	40	Hypolimnas bolina	62
†platyceps	56	Hypolycaena erylus	65
platyurus	37, 40, 52	Hypothymis azurea	275
reticulatus	57		
subtriedroides	40	Ianthia hyperythra	.. 277
Hemiplecta auriettae	582	rufilata	277
textrina	582	Icaria artifex	447
theodori	582	ornaticeps	.. 447
Hemiptera	82	variegata	447
Hemipus capitalis	271	Ichneumonidae	83, 323
Hemixus	268	†Idgia arbori	199
flavala	267	foveifrons	199
maclellandi maclellandi	268	Ilerda epicles	65
Henicurus leschenaulti	276	Ipidae	411
schistaceus	276	Ipidia variolosa	100
Hestina nama	62	Irdex	141
Heterolocha falconaria	439	nitidipennis	141
Heteromera	3	Iridopteryx	416
Heteroneurinae	175	irina	415, 416
Hierodula	416	Ischnura rufostigma	344
crassa	415, 416	Ixalus	.. 33, 34
Hinulia	43	anuandalei	7, 16, 33, 34
Hirudinidae	280	†argus	.. 7, 16, 33
Hister (Atholus) philippinensis	123	asper	7, 16, 27, 33, 34
Histeridae	121	assamensis	35
Holkeion auceps	537	horridus	25
Hololepta aequata	121	tuberculatus	7, 16, 17, 27, 29, 33
batchiana	121	Ixodes acutitarsis	98
elongata	121	angustus	97
indica	121	gigas	98
laevigata	121	Ixodidae	97, 98
procera	121	Ixulus flavicollis	.. 266
†Holophygus rugosus	111	occipitalis	266
Hoplia	192		
Hoplosternus	192	J	
nepalensis	192	Japalura	.. 54, 55
Hormurus australasiae	133	andersoniana	52, 58, 357
Humbertiella	416	†bengalensis	.. 57, 58
Hyalomma syriacum	97	planidorsata	.. 55, 58
†Hyberis similis	.. 106	variegata	.. 55, 58
Hydraena bihamata	624	yunmanensis	.. 57
cirrata	624	Juonia iphita	.. 62
†dimorpha	623, 624		
indica	624	K	
maculicollis	.. 624	Kachuga	38
Hydrochus annamita	624	intermedia	.. 38
binodosus	624	tectum	37, 38, 51
lacustris	624	Kallima inachus	62
latitans	624	Kalocrania	.. 135
opacus	624	picta	.. 136
violaceomicans	624	siamensis	135
Hydrophilidae	623	Khasiella	359, 593
Hydrophilus gibbus	623	†dinoensis	590, 593
Hydropsyche	356	Kolbea punctata	352
asiatica	356	Kosmetor	146
Hydropsychidae	355	brahma	146
Hydropsychodes	356		
Hydrophila	539	L	
Hylorana granulosa	9	Labeo angra	.. 244, 249
Hymenoptera	75, 81, 82, 83, 170, 233, 239, 323, 443, 444	dyochilus	.. 244, 249
Hymenoptera Anthophila	75	gonius	.. 244
Hypena occata	439		
Hyperythra lutea	.. 439		
Hypoctonus	127, 128		

	<i>Page</i>		<i>Page</i>
<i>Labia</i>	.. 142	<i>Lepus</i>	90
<i>curvicauda</i>	.. 142	<i>peguensis</i>	91
<i>mucronata</i>	142	<i>ruficaudatus</i>	90
<i>Labiduridae</i>	.. 137	<i>Lethe chandica</i>	61
<i>Labidurinae</i>	139	<i>Leuctra indica</i>	353
<i>Labiidae</i>	140	<i>Libellulidae</i>	.. 336
<i>Labiinae</i>	142	<i>Libellulinae</i>	336
<i>Labus humbertianus</i>	446	<i>Libnotes fuscinervis</i>	153
<i>Lacertidae</i>	.. 42	<i>punctipennis</i>	153
<i>Lacertilia</i>	37, 39	<i>Limicola</i>	287
<i>Lachesis gramineus</i>	51	<i>Liminites procris</i>	62
<i>monticola</i>	50	<i>Limnaea (Limnaea) acuminata</i>	539
<i>Lachnosterna</i>	192	<i>Limnaeidae</i>	539
<i>serricollis</i>	192	<i>Limnebius</i>	626
<i>sikkimensis</i>	.. 192	† <i>Limnophila claripennis</i>	153
• <i>Laemophloeus proximus</i>	110	† <i>quartarius</i>	154
† <i>semilaetaneus</i>	108	† <i>Limnophora kempii</i>	172
† <i>subturcius</i>	109	† <i>Limnosia magna</i>	173, 174
• <i>Lagocheilus</i>	494, 569	<i>subtinctipennis</i>	.. 174
† <i>daflaensis</i>	574, 575	<i>Lioptila annecteus</i>	265
† <i>oakesi</i>	574	<i>pulchella</i>	.. 265
† <i>sikhimensis</i>	574, 575	<i>Liopygus famelicus</i>	122
<i>tomotrema</i>	575	<i>Liothrix lutea</i>	266
<i>Lamellicornia</i>	.. 191	<i>Liotrichinae</i>	266
<i>Lampides bochus</i>	.. 64	<i>Liris aurata</i>	.. 444
<i>elpis</i>	.. 64	<i>ducalis</i>	445
<i>pura</i>	65	<i>Lispa ? pallitarsis</i>	172, 173
<i>Lampito</i>	389	<i>Litargus</i>	115
• <i>Lamprophorus</i>	.. 143	<i>Lithurgus dentipes</i>	76
<i>kervillei</i>	.. 143	<i>Litocerus khasianus</i>	197
<i>Landouria</i>	.. 604, 611	<i>paviei</i>	197
† <i>aborensis</i>	.. 605, 611, 614	<i>Lonchaea</i>	.. 179
† <i>damsangensis</i>	.. 605, 607, 609, 611, 614	<i>aenea</i>	179
<i>dawnaensis</i>	.. 614	† <i>montana</i>	178, 179
† <i>hengdanensis</i>	605, 608, 609, 610, 611, 614	<i>Lopaphus baucis</i>	422
<i>huttoni</i>	.. 605, 606, 614	<i>Loxura atymnus</i>	.. 65
<i>radleyi</i>	.. 605, 609, 610, 614	<i>Lucanidae</i>	515, 517
<i>Laniatores</i>	204	<i>Lucilia</i>	.. 171
<i>Laniidae</i>	.. 271	<i>Lumbricidae</i>	367
<i>Lanius tephronotus</i>	.. 271	<i>Luxiaria contigaria</i>	439
<i>Lariscus</i>	.. 90	<i>Lycaenesthes emolus</i>	64
<i>Lauxania</i>	176	† <i>Lychas graveleyi</i>	129, 130
† <i>flavicornis</i>	.. 179	<i>mucronatus</i>	128
<i>Lea arcuata</i>	.. 149	<i>rugosus</i>	130
<i>Lepidocephalichthys guntea</i>	243, 247	<i>scutilus</i>	129
<i>Lepidocyrtus</i>	.. 563	<i>Lygosoma albopunctatum</i>	.. 52
† <i>caudatus</i>	.. 563	† <i>courcyanum</i>	37, 43, 52
<i>imperialis</i>	.. 563	<i>indicum</i>	37, 43, 52
<i>maximus</i>	.. 563	<i>zebratum</i>	43
<i>pictus</i>	.. 563	<i>Lygosoma (Hinulia) cacharensis</i>	43
<i>Lepidoptera</i>	.. 61, 437	<i>Lyprops curticolis</i>	531
<i>Leptaulax</i>	.. 517	<i>Lyriothemis acigastra</i>	336
<i>bicolor</i>	.. 515, 517		
<i>cyclotaenius</i>	.. 517	M	
<i>cyclotaenius himalayae</i>	515, 517	<i>Mabuia macularia</i>	.. 37, 43, 51
<i>dentatus</i>	515, 517	<i>Macaca assamensis</i>	85
<i>roepstorfi</i>	.. 515, 517	<i>rhesus</i>	.. 85
<i>Leptocircus curius</i>	.. 63	<i>Macaria emersaria</i>	.. 439
<i>Leptogenys (Lobopetta) assamensis</i>	.. 233	† <i>Macrobunus aborensis</i>	.. 206
<i>Leptogomphus</i>	.. 340	† <i>longipes</i>	.. 205
<i>Leptoscapha</i>	.. 516	<i>singularis</i>	205, 206
<i>lignicola</i>	531	<i>Macrochlamyinae</i>	359, 581, 588
† <i>pulchra</i>	530, 531	<i>Macrochlamys</i>	359, 363, 547, 581, 584, 585, 586, 588, 590, 592
<i>spissicornis</i>	.. 530, 531	† <i>albulus</i>	.. 587
		† <i>bapuensis</i>	583, 584

	<i>Page</i>		<i>Page</i>
<i>Macrochlamys beata</i>	590	<i>Melittophagus swinhoei</i>	.. 284
† <i>burkilli</i>	586	<i>Melolontha argus</i>	.. 191
<i>causia</i>	582	<i>flabellata</i>	.. 192
<i>chaos</i>	582	<i>guttifera</i>	.. 192
<i>conseptata</i>	.. 581	<i>indica</i>	192
† <i>hardwickei</i>	582, 583	<i>Melolonthinae</i>	.. 192
<i>hardwickei politulus</i>	.. 582	† <i>Menophilus aborensis</i>	529
<i>hippocastaneum</i>	584	† <i>medius</i>	529
<i>hypoleuca</i>	582	<i>Menexenus</i>	421
<i>indica</i>	582, 583	† <i>rotunginus</i>	415, 421
† <i>luyorensis</i>	.. 587	<i>semiarmatus</i>	422
† <i>murdochi</i>	587	<i>Meranophus bicolor</i>	237
<i>uebulosa</i>	582	<i>Merganser castor</i>	288
<i>notha</i>	582	<i>Merginae</i>	.. 288
<i>noxia</i>	582	<i>Meromyza</i>	.. 188
† <i>psittacinus</i>	.. 584	<i>Meropes</i>	284
† <i>rotungensis</i>	584, 586, 589	<i>Meropidae</i>	.. 284
<i>rotungensis var.</i>	585	<i>Merula atrogularis</i>	278, 279
<i>salweenensis</i>	582	<i>ruficollis</i>	278, 279
† <i>shimangensis</i>	584	<i>Mesarmadillo</i>	543, 544
<i>Macrolinus</i>	516	<i>Mesogramma</i>	.. 163
<i>sikkimensis</i>	.. 515, 516	<i>Mesoperipatus</i>	486, 489, 496, 492
<i>Macromeris</i>	444	<i>Mesostenoides albomaculatus</i>	327
<i>honestata</i>	.. 444	<i>erythropus</i>	327
<i>Macrones affinis</i>	.. 253	<i>Metapodontus impressus</i>	517
† <i>merianiensis</i>	244, 253	† <i>Metathyreotus</i>	206
† <i>montanus dibrugarensis</i>	244, 254	† <i>aborensis</i>	206, 207
<i>tengara</i>	.. 244, 255	† <i>kempi</i>	.. 207
<i>vittatus</i>	244, 255	<i>Metisolabis</i>	.. 139
<i>Macronota nigricollis</i>	191	<i>caudelli</i>	139
<i>Macropeza</i>	150	<i>Metopius rufus</i>	324
<i>gibbosa</i>	150	<i>Micrixalus</i>	11, 33, 34, 54
<i>javanensis</i>	.. 150	† <i>borealis</i>	7, 10, 12, 29, 33
<i>Macrura</i>	.. 4	† <i>Microchrysa albitarsis</i>	156
<i>Malacodermidae</i>	119	<i>Microcichla scouleri</i>	276
<i>Mantidae</i>	.. 415	<i>Microgaster luteus</i>	.. 323
<i>Mantis aridifolia</i>	416	<i>Micropezinae</i>	186
<i>chloreudea</i>	.. 416	<i>Microserica</i>	193
<i>Mantis (Mantis) maculata</i>	416	† <i>viridicollis</i>	192
<i>servillei</i>	416	† <i>Mikiria</i>	.. 611
<i>Mazarredia aptera</i>	.. 314	† <i>diyungensis</i>	611, 612, 614
<i>Mecocerus allectus maculatus</i>	197	<i>Miltochrista perpallida</i>	.. 437
<i>Medaura austeni</i>	422	<i>undulosa</i>	438
<i>Megalaema</i>	.. 259, 260	† <i>Minyongia</i>	556, 559
<i>marshallorum</i>	283	† <i>kempi</i>	556, 559
<i>Megalophrys</i> 16, 20, 21, 27, 28, 29, 30, 34		† <i>Mixogaster vespiformis</i>	169
<i>hasseltii</i>	.. 29	<i>Mollusca</i>	209, 359, 493, 505, 537, 547, 569, 581, 601
<i>heteropus</i>	20, 29	<i>Molpastes haemorrhous bengal-</i>	
† <i>kempi</i>	7, 20, 29, 34	<i>ensis</i>	.. 268
<i>major</i>	7, 16, 29, 34	<i>haemorrhous burmanicus</i>	268
<i>montana</i>	.. 28, 29, 30	<i>Moniligaster</i>	.. 410
<i>parva</i>	.. 29, 31, 32	<i>grandis</i>	410
<i>Megascolecidae</i>	367	<i>Moniligastridae</i>	366, 367
<i>Megascolides</i>	365, 366, 367	<i>Monticola</i>	.. 279
† <i>oneilli</i>	368, 377	† <i>Moringua hodgarti</i>	244, 255
<i>Megauchenia angustata</i>	.. 100	<i>Moringuidae</i>	.. 255
<i>quadricollis</i>	.. 100	<i>Motacilla alba</i>	280
<i>Melanitis leda ismene</i>	.. 62	<i>alba ocularis</i>	.. 280
<i>phedima bela</i>	.. 62	<i>Motacillidae</i>	.. 280
<i>Melanochlora sultanea</i>	.. 267	<i>Muridae</i>	93, 94, 96
<i>Melanopa atrata</i>	.. 203	<i>Mus rattus</i>	.. 90
<i>varians</i>	.. 203	<i>Musca domestica</i>	.. 171
<i>Melanostoma mellinum</i>	.. 164	<i>Muscicapidae</i>	.. 274
<i>orientalis</i>	.. 164	<i>Muscicapula maculata</i>	.. 87
<i>scalare</i>	.. 164	<i>Muscidae</i>	.. 170
<i>univittatum</i>	.. 164	<i>Muscidae verae</i>	.. 170
<i>Melcha maculiceps</i>	327		

	<i>Page</i>		<i>Page</i>
Muscinae	170	Nodaria externalis ..	439
Mustela flavigula flavigula ..	86	innocens ..	439
<i>flavigula typica</i> ..	86	Nomia aurifrons ..	76
Mutilla ..	443	terminata ..	76
blanda ..	443	Nomioides ..	83
cassiope ..	443	Notodea ..	278
Mutillidae	443	Notogonia ..	444
Mycalesis meda	62	Notoscolex ..	365, 366, 367
perseoides khasia ..	62	†stewarti ..	366, 368, 382
visala	62	striata ..	383
Mycetophagidae	99, 115	striatus ..	365, 368, 380
Mycetophilidae	149	Nuria danrica ..	244, 251
Myiophoneus temminckii eugenii ..	278	Nyctemera coleta ..	437
temminckii temminckii ..	278	varians ..	437
Myophoneus	259	Nycticejus ornatus ..	87
Myopopone castanea maculata ..	233		
Myrmemorpha	188	O	
Myrmicinae	234	Odonata	4, 335, 336
Myronides	422	Odontomachus monticola ..	233
baucis	415, 422, 423	Odontotermes	428
†dawnanus	415, 422	assamensis ..	428
Myxostoma	493, 569	feae ..	428, 430
		†gravelyi ..	428, 429
N		microdens ..	428
Nacaduba bhutea	64	Odynerus fragilis ..	446
nora	64	icaroides ..	446
noreia	64	Oecophylla smaragdina ..	237
Nala	139	†Oedematopsis aborensis ..	324
nepalensis	139	apollos ..	324
Nandidae	256	ops ..	324
Nandus nandus	244, 259	Oegista	613
Nannophya pygmaea	337	Oligochaeta	365, 410
Naxa textilis	440	Oligodon erythrorachis ..	37, 48, 53
Neanias jacobsoni	333	Oncopodidae	207
†kempi	332	Oniticellus vertagus ..	194
pupulum	333	Onitis philemon	194
Neanura	561, 562	Onthophagus	195
muscorum	563	luzonicus	195
pudibunda	562	tarandus	194
Neanurinae	561	†triceratops	195
Nectariniidae	281	Onychargia vittigera ..	343
Nectariniinae	281	Oonychophora ..	471, 473, 487, 488, 490
Nemachilichthys	245	Ooperipatus	483, 488
Nemachilus	245	Opeas	360, 615
rupicola	31, 32	Ophidia	37, 44
Nemorhaedus bubalinus	91, 92	Ophiocephalidae	256
Nemoura	353	Ophilocephalus gachua ..	244, 256
Neoperla	354	gracilis	37, 42, 52, 357
Neopithecops zalmora	64	Ophyra anthrax	172
Neptis hordonia	62	caerula	171
hylas varmona	62	Opiliones	203
soma	62	Opisthocosmiinae	146
Nerius annulipes	184	Opisthopatus	484
†obscurus	183	blainvillei	484, 488
tibialis	184	cinctipes	488
Neurobasis sinensis	340	Orchestia	451, 452
Neuroptera	4, 354	parvispinosa	449, 451, 453
Neurothemis	338	Orchestoidea	451
fulvia	338	†Oreinus molesworthi ..	244, 247
intermedia	337, 338	plagiostomus	243, 244, 247
intermedia degener	337	richardsoni	243, 244, 247
tullia	337	sinuatus	243, 244, 247
Nicoria tricarinata	51, 54	Oriolidae	273
Niltava macgrigoriae	274	Oriolus melanocephalus ..	273
sundara	274	trailii	273
Nitidulidae	99, 100	Oromothrips sanguineus ..	201

	<i>Page</i>		<i>Page</i>
Orthetrum glaucum	336	Paraliris faceta	445
pruniosum neglectum ..	336	Paraperipatus	486, 488, 489
sabina	336	†Paraphytus hindu	193, 194
triangulare	337	ritsemae	194
Orthoptera	3:1	Paratelfhusa	302
Orthotomus sutorius	270	Paratelfhusa (Barytelfhusa) feae	301
Oscinella	190	harpax	289, 302
Oscininae	188	Paratenodera	416
Oscinis	190	†Parepierrez	124
obscuripes	190	amandus	124, 125
Osphromenidae ..	257	†corticicola ..	124, 125
Osphromenus nobilis	244, 257	Pareronia avator	64
olfax	257	Parinae	261
Ostariophysa ..	244	Parmarion ? rubrum.	557, 558
Otocompsa emerina emerina	268	Paromalus pardali	125
flaviventris	269	tcibodae	125
Otostigminae	69	vermiculatus	125
Otostigmus	69	Paronela	561
†burn-murdochi	69	Paronella	563, 564
insularis	70	borneri	566
rugulosus	71	†crassicornis	564
scaber	71	dahlii	566
Oxytes	547	†elongata	565
†aborensis	595, 597	†flava	565
blanfordi ..	595	Paronellini	563, 564, 565, 566
cycloplax	594	Parorchestia	.. 451
†oglei	594	Parus	261
oglei var.	594	major	261
oslei	359	major atriceps	261
oxytes	359, 594	major cinereus	261
pollux	595	Passalidae	3, 515
shanensis	595	Passandridae	99, 111
†siyomensis	595	Passer rutilans	280
		rutilans cinnamomea	280
		debilis	280
		Passeres	260
		Pectinibranchia ..	539
		Pedipalpi	127, 128
		†Pelitnus aborensis .	207
		annulipes ..	207
		Pellorneum mandellii	263
		mandellii minus	263
		Pelmatoplana aurantia	462
		sondaica	463
		Pelobatidae	4, 20, 28
		Percesoces	256
		Perciformes ..	256
		Pericrocotus fraterculus	272
		solaris ..	272
		speciosus	271, 272
		Perionyx	365, 366, 389, 398
		†aborensis	368, 392
		†annulatus ..	368, 372, 386
		arboricola 365
		†depressus ..	365, 368, 377, 394
		excavatus	368, 386, 399
		†foveatus	368, 396
		†kempi	368, 389, 399
		†koboensis ..	368, 389, 391, 396,
		sansibaricus	.. 399
		Peripatidae ..	472
		Peripatoides 483, 486, 488, 489, 490, 491,	492
		novae zealandiae	478, 487
		Peripatopsis ..	484, 489
		cinctipes ..	483

P

Pachycondyla (Ectomomyrmex)	
astutā ..	233
javana materna	233
Pachylophus	189
rufescens	188, 189
Palaemon ..	4, 303
dayanus	304
hendersoni	289, 290, 303, 304, 305
Palaemon (Parapalaemon ?) hen-	
dersoni ..	303
Palaemonidae ..	303
Palaeornis fasciatus.	286
Palpatores	203
Palpopleura sexmaculata	337
Paludetrinidae	540
Paludina	540
Paludomus	537
†rotungensis	537, 539, 540
Pandanus	4
Pandarus odoratissimus	102, 104
Pangshura cochinchinensis	38
Pantoporia zeroa	62
Papilio helenus	63
mennon agenor.	63
paris	63
philoxenus polyeuctes ..	63
polytes romulus ..	63
protenor euprotenor	63
Paradoxornithinae	261
Paragus rufiventris ..	157
serratus ..	158

	Page		Page
Peripatus	471, 472, 473, 474, 476, 478, 481, 483, 486, 488, 489, 490, 491, 492	Platysoma confucii	.. 123
ceramensis	488	†crassum	123
edwardsii	.. 483	silvestre	123
sedgwicki	.. 483	(Platylister) cambodjensis	122
Perlidae	.. 353	dissimile	122
Petaurista magnificus	88	†kempi	123
Petrophila	259, 279	marseuli	122
erythrogaster	279	odiosum	122
solitarius pandoo	279	Pleciomyia melanaspis	150
Phalangidae	203	Plecoptesa	353
Phalangodidae	206	Plectopylis	360, 493, 505, 508, 512
Phasianidae	287	brachyplecta	.. 512
Phasniidae	415, 417	brahma	509, 511
†Pheidologeton (Aneleus) abor-		emigrans	513
ensis	234, 236	fruhstorferi	513
Pheretima	365, 366, 367	hirasei	.. 513
anomala	.. 366	macromphalus	507
heterochaeta	366, 368, 399	muspratti	.. 508
lignicola	366, 368, 399	oglei	511
Philaematomyia insignis	.. 171	pinacis	506, 508
Philopotamus	355	plectostoma	.. 507, 509
Philoscia	455, 466	pulvinaris	.. 513
coeca	.. 465	sowerbyi	507, 509
Phloeopemon acuticorne	198	vulvivaga	602
Phoenicopharinae	285	†(Chersaecia) bedfordi	510
Phricotelphusa	.. 291	†williamsoni	509
Phrynoderma	.. 34	†(Endoplion) aborensis	511
asperum	.. 18	†(Endothyra) gregorsuni	.. 506
†moloch	7, 18, 21, 25, 29, 34	†miriensis	.. 507, 508
Phyllium	417	†oakesi	.. 505
celebicum	.. 415, 417	†(Sinicola) babbagei	512, 513
Phymatocerinae	240	Plectotropis	360, 601, 602, 606, 613
Pici	282	akoutongensis	.. 602
Picidae	282	austeni	.. 537, 538, 614
Picinae	.. 282	catostoma	.. 602
Picumninae	283	clarus	602
Pieris canidia	.. 63	emensus	602
Pileocera aegimiusalis	440	grumulus	.. 602
Pipistrellus abramus	87	huttoni	.. 601, 602
Pitta	.. 259	luttoni savadiensis	602
nepalensis nepalensis	282	mitanensis	.. 602
Pittidae	282	oldhami	.. 602
Placocephalus superbus	460, 462	orbicula	.. 602
Plagiolepis longipes	237	pudica	602
Plagiopholis	.. 45, 54	radleyi	.. 601
Planaria	317, 319	rotatoria	.. 602
†aborensis	317	sumatrana	.. 602
†kempi	319, 320	tapeina	602, 612
subtentaculata	.. 317	Plilona dunlopi	185
venusta	.. 318	Ploceidae	279
vitta	318	Plusia orichalcea	438
Planispira	.. 360	Plutellus	365, 366, 367
delibrata fasciata	.. 360	†aborensis	368, 384
Planorbis exustus	.. 539	Poduridae	.. 561
(Gyraulus) himalayanus	.. 539	Polistes hebraeus	447
Platanista gangetica	38	Polybia indica	.. 446
Platycnemis	.. 341	orientalis	446
†Platydemia alticornis	524	Polydonthophis collaris	37, 46, 53
annamitum	524	Polyommatus boeticus	65
†auremaculata	.. 523, 524	Polypedates afghana	9
subfascia	.. 524	Polyrhachis dives	237
Platylister	123	halidayi	237
Platypodidae	413	laevissima	237
Platyptilia taprobanes	441	mayri	237
Platypus	414	striatorugosa	237
†cupulifer	413	tibialis	237
		Pomatias	494, 569, 580

	Page		Page
Pomatiasinae ..	494, 569, 577	Pseudaustenia ater ..	592
Pomatorhinus ferruginosus ..	263	Pseudecheneis ..	32
Pompilidae ..	444	sulcatus ..	243, 244, 255
Pompilus ..	444	Pseudergolis wedah ..	63
Ponera confinis ..	233	Pseudeutropius atherinoides ..	244, 245
confinis wroughtoni ..	233	garua ..	244, 255
Ponerinae ..	233	Pseudoblaps javana ..	520
Porcellionides prunosus ..	465	†Pseudokaliella annandalei ..	596
Porifera ..	67	nevilli ..	597
Poritia hewitsoni tavoyana ..	65	†sadiyaensis ..	597
Potamiscus 230, 289, 290, 291, 296		Pseudomantis haanii ..	416
Potamon 229, 230, 290, 291, 297		maculata ..	416
adiatretum ..	229	Pseudominea castaneiceps ..	264
andersonianum ..	229	Pseudopomatias 569, 577, 579	
annandalei ..	291, 294	grandis ..	578
manii ..	229	himalayae ..	578
pealinum ..	293	†luyorensis ..	578
sikkimense 291, 292, 294, 295, 296,	297	pleurophorus ..	578
superciliosum ..	229	†siyomensis ..	578
tumidum ..	291	Pseudoxenodon ..	54
(Acanthotelphusa) feae 289, 301,	302	macrops ..	37, 49, 53
(Geotelphusa) adiatretum 289, 298,	299, 300, 301	Psila kempfi ..	187
*var. lophocarpus 289, 299, 301		Psilinae ..	187
†superciliosum 289, 300, 301		Psilopus ..	157
†(Potamiscus) aborense 289, 292,	294, 295, 296, 297	Psittaci ..	286
annandalei ..	293	Psittacidae ..	286
†decourcyi 289, 291, 292, 293,	294	Psocidae ..	351
†obliteratum 289, 292, 294,	295, 296, 297	†Psocus nirvanus ..	352
Potamonidae ..	290, 291	†Psychoda notatipennis ..	151
Potamoninae ..	290, 291	Psychodidae ..	151
Pratincola maura ..	276	Pterocyclos 494, 499, 502, 569, 570	
torquata przewalskii ..	276	†aborensis ..	498
Presbytes ..	85	†brahmakundensis ..	499
entellus ..	85	magnus ..	500
schistaceus ..	85	†miriensis ..	498, 500
†Pria diluticollis ..	102	†spiramentum ..	498, 500
†mirmidon ..	101	Pteromys magnificus ..	88
†Prisma aborense ..	209, 222	Pteropus giganteus leucocephalus	
Pristomyrmex brevispinosus ..	237	leucocephalus ..	81, 97
Prodenia littoralis ..	438	medius ..	87
Prosobranchia ..	493	..	87, 90
Prosopis ..	83	Pterutheus melanotis ..	267
Protaetia fusca ..	192	Ptyctolaemus ..	41, 54
inanis ..	191	gularis ..	37, 39, 41, 52
inanis cuprea ..	191	Pulmonata ..	537
Protanura ..	561, 562	Pupina ..	573
citronella ..	562	Pupipara ..	190
kraepelini ..	562	Pygidicranidae ..	135
spinifera ..	562	Pygidicraninae ..	135
Protodermaptera ..	135	Pygoplus trifasciatus ..	205
Psalinae ..	137	Pyrellia violacea ..	170
Psalis ..	138	Pyrocoelia lampyroides ..	119
dhorni ..	138	Pyrrhopicus pyrrhotes ..	283
femoralis ..	138		
Psammodynastes pulverulentus ..	37,	R	
..	50, 54	Rahula ..	599, 603
Psenulus ..	446	†aborensis ..	597, 598, 599
pulcherrimus ..	446	†burrailensis ..	598
Pseudagenia ..	444	†dihingensis ..	597, 598
tinctoria ..	444	†koboensis ..	598, 599, 600, 621
Pseudagrion ..	344	municipurensis ..	599
		Rana ..	11, 24, 31, 32
		afghana ..	7, 9, 21, 24, 31, 32, 33
		alticola ..	7, 8, 9, 21, 22, 33, 35
		blanfordii ..	8
		cyanophlyctis ..	7, 32, 35

	<i>Page</i>
<i>Rana erythraea</i> ..	9
† <i>gerbillus</i> ..	7, 10, 33
<i>granulosa</i> ..	7, 9, 33, 35
<i>jerboa</i> ..	10
<i>latopalmata</i> ..	9
<i>liebigii</i> ..	7, 8, 21, 28, 31, 32
<i>limnocharis</i> ..	7, 8, 29, 33
<i>tenasserimensis</i> ..	10, 11
<i>tigrina</i> ..	7, 8, 32, 35
<i>tylleri</i> ..	9
Ranidae ..	4, 7, 21, 32
<i>Rapala schistacea</i> ..	65
<i>Raparna digramma</i> ..	438
Raphaulus ..	494, 569, 570
† <i>aborensis</i> ..	571, 572, 573
<i>assamicus</i> ..	541
<i>assamica</i> ..	570, 571
<i>blanfordi</i> ..	573, 574
† <i>oakesi</i> ..	572
† <i>shimangensis</i> ..	572, 573
† <i>yamneyensis</i> ..	571
Raptomphalus ..	502, 503, 569
<i>Rasbora daniconia</i> ..	252
<i>rasbora</i> ..	252
* <i>rasbora kobonensis</i> ..	244, 251
Rathouisiidae ..	209
<i>Ratufa gigantea</i> ..	88, 93
<i>macruroides</i> ..	88
<i>melanopepla</i> ..	88
Reptilia ..	5, 37, 357
<i>Rhacophorus</i> ..	16, 27, 33
<i>bimaculatus</i> ..	7, 12, 33
<i>leucomystax</i> ..	14, 24
<i>maculatus</i> ..	14, 21, 24, 25
<i>maculatus himalayensis</i> ..	7, 14, 24, 33
<i>maculatus leucomystax</i> ..	14
<i>maximus</i> ..	7, 12, 33
† <i>microdiscus</i> ..	7, 13, 33
† <i>naso</i> ..	7, 12, 16, 27, 33
<i>reinwardtii</i> ..	24
<i>tuberculatus</i> ..	7, 14, 33
Rhamphidia ..	153
<i>Rhaphedophora brunneri</i> ..	329
<i>Rhingia binotata</i> ..	166
<i>sexmaculata</i> ..	166
<i>Rhinocypha quadrimaculata</i> ..	341
<i>Rhinia limbipennis</i> ..	170
<i>Rhinopalpa polynice birmana</i> ..	62
<i>Rhinosciurus</i> ..	90
<i>Rhipidura albicollis</i> ..	260
<i>albicollis albicollis</i> ..	276
* <i>albicollis kempii</i> ..	275, 276
<i>Rhopalocampta benjaminii</i> ..	65
<i>Rhopodytes</i> ..	260, 283
<i>tristis</i> ..	285
<i>Rhyacornis fuliginosus</i> ..	277
<i>Rhynchium argentatum</i> ..	446
<i>Rhynchomyia</i> ..	170
<i>Rhynchophora</i> ..	3
<i>Rhynchota</i> ..	16
<i>Rhyothemis</i> ..	383
<i>curiosa</i> ..	338, 339
<i>fulgeus</i> ..	339
<i>obsolescens</i> ..	339
<i>plutonia</i> ..	338
<i>pygmaea</i> ..	339

	<i>Page</i>
<i>Rhyothemis variegata</i> ..	338
<i>Rhysida</i> ..	71
<i>cuprea</i> ..	72
<i>immarginata</i> ..	71
<i>nuda</i> ..	71
<i>petersi</i> ..	72
<i>stuhlmanni</i> ..	72
† <i>Rhysodes alticola</i> ..	99
<i>intrusus</i> ..	100
<i>longiceps</i> ..	100
Rhysodidae ..	99
<i>Rioxa</i> ..	185
† <i>Rotungia</i> ..	590, 591
† <i>williamsoni</i> ..	590
† <i>Rotungus</i> ..	543
† <i>pictus</i> ..	544
Ruteinae ..	591
<i>Ruticilla aurea leucoptera</i> ..	277
<i>frontalis</i> ..	277
S	
<i>Saccobranchus fossilis</i> ..	244, 255
<i>Saedjahus</i> ..	543
<i>Salea</i> ..	34, 54
<i>austeniana</i> ..	52
<i>Salius praestabilis</i> ..	444
<i>Sapromyza</i> ..	176
<i>histrion</i> ..	176
<i>Sapromyzinae</i> ..	178
<i>Saprosites marchionalis</i> ..	196
<i>Sarama</i> ..	359, 363
<i>kala</i> ..	362, 363
† <i>kempii</i> ..	359, 362, 581, 591
<i>Sarcophaga</i> ..	170
<i>Sarika</i> ..	582
<i>concepta</i> ..	614, 618
<i>conseptata</i> ..	537
<i>Sasia ochracea</i> ..	283
<i>Saturnia pyretorum</i> ..	437
<i>Sauris abortivata</i> ..	440
<i>lineosa</i> ..	440
<i>Scaeorhynchus ruficeps ruficeps</i> ..	261
<i>Scelimena</i> ..	311, 312
<i>india</i> ..	311
<i>Sceliphron madraspatanam</i> ..	445
† <i>Schedorhinotermes magnificus</i> ..	425, 426
<i>malaccensis</i> ..	426
<i>Schizopygopsis stoliczkae</i> ..	249
<i>Sciara</i> ..	149
<i>orientalis</i> ..	149
Scincidae ..	43
<i>Sciomyza</i> ..	176
† <i>ocellata</i> ..	176
<i>quinquevittata</i> ..	176
† <i>septemlineata</i> ..	178
<i>strigata</i> ..	176, 178
<i>trypetoptera</i> ..	176
<i>Sciomyzinae</i> ..	176
<i>Sciuridae</i> ..	93, 94, 96
<i>Sciuropterus alboniger</i> ..	88
<i>Sciurus</i> ..	90
<i>atrodorsalis</i> ..	89
<i>bicolor</i> ..	88
<i>castaneoventris</i> ..	88
<i>castaneoventris griseopectus</i> ..	89
<i>erythraeus</i> ..	88

	Page		Page
<i>Sciurus erythraeus erythrogaster</i>	88	Sittidae	269
<i>erythraeus intermedius</i>	89, 93, 94,	<i>Siva cyanuroptera</i> ..	266
	97	<i>Sivella</i>	360, 585
<i>erythrogaster</i>	88	<i>castra</i>	603, 604
<i>giganteus</i> ..	88	* <i>castra kobonensis</i>	604
<i>gordoni intermedia</i>	89	<i>Solenosoma</i>	145
<i>griseopectus</i>	89	<i>birmanum</i>	145
<i>locria</i>	90	<i>Sorex fuliginosus</i>	87
<i>lokriah</i> ..	90	<i>Speculitermes cyclops</i>	431
<i>maccebellandi barbei</i>	89	<i>Sphaerophoria scutellaris</i>	164
<i>maccebellandi manipurensis</i>	89	<i>Sphecodes</i>	75
<i>maccebellandi novemlineatus</i>	89	<i>Sphegidae</i>	444
<i>maccebellandi typicus</i>	89	† <i>Sphegina tristriata</i>	165
<i>pernyi</i> ..	90	<i>Sphenisomyia sexmaculata</i>	185
<i>stevensi</i>	85, 89, 95	<i>Sphex umbrosus</i>	445
<i>Scleron ferrugineum</i>	520	<i>Spilogaster</i> ..	173
<i>Scolia</i>	443	<i>Spiraculum</i> 359, 494, 499, 500, 501, 502,	569, 570, 581
<i>floridula</i>	443	<i>andersoni</i>	501
<i>kirbyi</i>	443	† <i>kempi</i>	496, 467, 501
<i>sikkimensis</i>	443	† <i>luyorensis</i>	500, 502
<i>Scoliidae</i>	443	† <i>minimum</i>	501, 502, 581
<i>Scolopendra</i>	72	<i>nevilli</i>	497, 501
† <i>mazleii</i>	72	† <i>oakesi</i>	496
<i>morsitans</i>	73	† <i>planum</i>	497
<i>Scolopendridae</i>	69	† <i>putaoensis</i> 500, 501, 502, 581	
<i>Scolopendrinae</i>	72	<i>Spongilla crassissima</i>	67
<i>Scombresocidae</i>	256	<i>decipiens calcuttana</i>	67
<i>Scorpiops binghami</i>	133	<i>Spongilla (Eunapius) crassissima</i>	
<i>longimanus</i>	132, 133	<i>crassior</i> ..	67
<i>montanus</i>	133	<i>Spongilla (Euspongilla) proli-</i>	
<i>Selandriinae</i>	241	<i>ferens</i>	67
<i>Seopsis</i>	353	<i>Spongophorinae</i>	140
<i>Sepsinae</i>	185	<i>Spongovostox</i>	140
<i>Sepsis bicolor</i>	185	† <i>aborum</i>	140
<i>coprophila</i>	185	<i>kristenseni</i>	141
<i>indica</i>	185	<i>luteus</i>	140, 141
<i>viduata</i>	185	<i>Stachyris chrysaea</i>	264
<i>Serica</i>	193	<i>nigriceps</i>	264
<i>Sericomyia</i>	169	<i>Staphylinidae</i>	3
† <i>ristalioides</i>	167	<i>Statilia</i>	416
<i>Serranidae</i>	256	<i>haanii</i>	415, 416
† <i>Sesara globosa</i>	598, 599	<i>maculata</i>	416
<i>Setenis</i>	527	<i>Stenopelmatidae</i>	329
<i>crenatostrata</i>	526	<i>Stenopsyche griseipennis</i>	355
<i>dentipes</i>	527	<i>Stenopterina aenea</i>	183
<i>indosinica</i>	526	<i>eques</i> ..	183
† <i>kempi</i> ..	526	† <i>flavofemorata</i>	182
<i>laevis</i> ..	526	<i>Sternolophus (Neosternolophus)</i>	
<i>Sibia picaoides</i>	264, 265	<i>tenebricosus</i>	628
<i>Sibiinae</i>	264	<i>Stibocheona nicea</i>	62
<i>Silsila fulvipes</i>	327	<i>Stictolissonota foveata</i>	323
<i>Siluridae</i>	252	<i>Stegmus niger</i>	446
<i>Silvanus lateritius</i>	111	<i>Stoparola melanops.</i>	274
<i>Sima rufonigra</i>	234	<i>Stratiomyidae</i>	156
<i>Simotes albocinctus.</i>	37, 48, 53	<i>Streptaulus</i>	569, 570, 573
<i>juglandifer</i>	48	<i>blanfordi</i>	570
<i>Simplicia niphona</i>	439	† <i>luyorensis</i>	573
<i>Simuliidae</i> ..	150	† <i>miriensis</i>	573, 574
<i>Simulium</i>	150	<i>Streptoste (Elma) nevilli dubia</i>	593
<i>indicum</i> ..	150	<i>Stromboceros</i>	241
<i>Sinicola</i>	505, 573	<i>Stromboceros (Neostromboceros)</i>	
<i>Siphia</i>	274	<i>congener</i> ..	241
<i>albicilla</i>	274	* <i>congener tarsalis</i>	241
<i>parva</i>	274	<i>Strongylium cultellatum</i>	535
<i>Sirella</i>	359	† <i>curvicomis</i>	535
<i>Sitta cinnamoneoventris</i>	259, 269	<i>sobrinum</i>	534, 535
<i>neglecta</i>	259, 269		

	<i>Page</i>		<i>Page</i>
Strongylium †stevensi	534	Terias hecabe	64
westermanni ..	535, 536	silhetana	64
Sturnidae ..	273	†Termes annandalei	427
Sturnopaster ..	260	malaccensis	428
Sturnopasta contra ..	273	Termitidae ..	425
contra superciliaris	273	Tetriginae ..	311
Sufetula sunidesalis ..	441	Teucholabis biannulata	153
Sumniva ..	543	Themara maculipennis	185
Sus cristatus ..	92, 98	†Themus aborensis	119, 120
jubatus ..	92	Theopompa ..	416
Sylviidae ..	270	servillei	415, 416
Symbrenthia hippoclus khasi-		Thermesia rubricans	438
ana ..	62	Tholymis tillarga	339
Sympetrum orientale	338	Tho adonta	312
Syngamia floralis ..	441	spiculoba	311, 312
Sypna quadrisignata	438	Thraulius	354
Syrphidae ..	157, 164	Thymaris clotho	326
Syrphus ..	163, 164	Thyreotus ..	206
†aeneifrons	159, 160	bimaculatus	206
balteatus ..	159	Thysanoptera	201
cinctellus ..	163	Tiara (Melanoides) variabilis	539
cinctus ..	163	Tiara (Tarebia) lineata ..	539
†fulvifacies ..	161	Tiaridae	539
†maculipleura ..	162	Tiberioides	516
ribesii	162	austeni	515, 516
†transversus	160	Timeliinae ..	263
vitripennis ..	162	Timomenus ..	147
		aeris ..	147
T		Tinda indica ..	157
Tabanidae ..	157	Tinnunculus alandarius alanda-	
Tachininae ..	170	rius	286
Tachydromus sexlineatus	37, 42, 52	Tiphia intrudens ..	443
Tachysphex	444	Tipulidae	152
†Tadunia ..	588	Tomostethus	240
†muspratti ..	588	†assamensis	241
†oakesi ..	588, 589	barda	240
Taeniocerus ..	515	formosanus ..	241
bicuspis ..	515	†hirticornis	240
Taenioglossa ..	539	Tonkinus sculptilis	527
Talitridae	451, 452	striatipennis	527
Talitrus ..	451	Totaninae ..	287
Talorchestia ..	451, 452	Totanus ochropus ..	287
†kempi ..	449, 451, 452, 453	Toxicum ..	530
martensii ..	452	assamense	529
novaehollandiae ..	452	Toxorhynchites	152
parvispinosa ..	452, 453	Trachia delibrata	615
Talpa leucura	86	delibrata fasciata	614
micrura ..	86	Trachischium monticola	37, 45, 53, 357
Tamiops macclellandi	89, 95	Trachypholis decorata	107
Taphrospira	359	hispida	107
Tapinoma melanocephalum ..	237	signata ..	107
†Tarphiosoma kempi ..	104	Tribalus colombius ..	125
Tearchus annulipes ..	532	pluristriatus	125
Technomyrmex albipes	237	Trichogaster fasciatus	244, 257
Temnocephala ..	229, 230, 231, 232	Trichomma	325
chiliensis ..	232	decorum	325
fasciata ..	231	Trichoptera	355
madagascarensis ..	231	Trichopteryx sikkima	440
semperi ..	229, 230, 231, 290	Trichorypha ..	564
Temnocephalidae ..	229, 230	Trigonometopus frontalis	176
Temnocephaloidea ..	232	†trilineatus	175
Tenebrionidae	519, 526	Trigonophorus nepalensis	191
Tenebrioninae ..	530	Trimeresurus gramineus	37, 51, 54
Tenodera ..	416	monticola ..	37, 50, 51, 54
aridifolia	415, 416	Trirhinopholis	45, 54
Tenthredinidae	239	Trithemis aurora	338
		festiva	338

xxviii

	Page		Page
Trithemis pallidinervis	338	Vespa ducalis ..	447
Trochomorphoides acris	615	magnifica ..	447
Trogones	285	<i>Vesperugo abramus</i> ..	87
Trogonidae ..	285	Vespidae	446
Tropidonotus khasiensis	37, 49, 53	Vestalis amoena	340
piscator	37, 49, 53	gracilis ..	340
platyceps	37, 49, 53	Videna bicolor	604
†assamensis	58	conicoides ..	604
berdmorei	59	trilineatus ..	604
cochinensis	58	Viduiinae	279
yunnanensis	59	Vipera russeli	51
Trypeticus nemorivagus	121	Viperidae	50
Trypetinae	185	Viverra zibetha ..	85
Trypoxylon	445	Vivipara (<i>Idiopoma</i>) heliciformis	540
Tubulifera	201	Viviparidae ..	540
Tupaia belangeri	86, 97		
chinensis ..	86	X	
<i>ferruginea</i> ..	86	Xanthogramma ..	163
Turdidae	276, 278	Xanthopimpla nursei ..	323
†Typhloperipatus	479, 483, 484, 486, 487, 489, 490, 491	punctata	323
†williamsoni	472, 476	Xenocerus khasianus	198
Typhlopidae	.. 44	rectilineatus ..	198
Typhlops beddomii	44	Xiphocaridina	491
braminus	44, 52	curvirostris	290, 491
diardi	37, 44, 53	Xuthia parallela	104
†diversiceps	37, 44, 53	Xyleborus	411, 413
tephrosoma	37, 44, 53	†aplanatus	412
		capucinus ..	413
U		†gravelyi ..	411
Uleiota atratula ..	108	perforans	413
Uloma javana 525	Xylinales annulipes	198
orientalis 525	plagiatus ..	198
orientalis minor 525	Xylocopa ..	82
Ulominae	530	collaris	79
Upupa epops indica.	284	dejeanii	79
Upupae	284	latipes	79
Upupidae ..	284		
Uroloncha acuticauda	279	Y	
Uroproctus	127	Ypthima affectata	61
assamensis	127, 128	baldus	61
Ursus torquatus	86	philomela indecora	61
V		Z	
Vanessa indica ..	62	Zaleptus sulphureus ..	204
Varanidae	42	Zamenis mucosus ..	37, 48, 53
Varanus bengalensis	37, 42	Zeltus etolus	65
Vespa basalis ..	447	Zemerus confucius ..	63
bicolor ..	447	flegyas ..	63
cincta ..	447	Zonitidae ..	359, 537, 548, 559, 569, 603, 604
dorylloides	447		

ZOOLOGICAL RESULTS OF THE ABOR EXPEDITION, 1911-1912.

INTRODUCTION.

In accordance with a resolution passed at the Conference on Museums and Archæology held at Simla in July, 1911, it was arranged by the Trustees in consultation with the Military and Education Departments of the Government of India that I should accompany the Abor Expeditionary Force as Zoologist and Anthropologist, and that Mr. R. Hodgart, Zoological Collector in the Indian Museum, should go with me as assistant.

The present volume is devoted to the zoological results and parts will be issued from time to time as reports become available for publication. The issue of the usual annual volume of the "Records" will continue as heretofore.

In addition to results obtained with the Abor Expeditionary Force, in several groups the determinations of animals obtained by Mr. F. H. Gravely in November and December, 1911, between Moulmein and the Siamese frontier will be included. The fauna of this area, at least of the Thaungyin Valley, is not very dissimilar from that of the Abor country, and notice of specimens from this area will be of interest from a comparative point of view.

Mr. Hodgart and I left Calcutta on November 13th and arrived back on April 3rd after having spent about four months in the Abor country.

On the outward journey we were detained for about a fortnight before we could proceed to Kobo, the base camp of the expedition, and this period was spent in making observations on the fauna at Dibrugarh on the left bank of the Brahmaputra and at Sadiya which lies on the right bank at the foot of the Mishmi country.

We reached Kobo on November 29th and remained until December 13th; on the 14th we left by boat-convoy up the Dihang River and arrived at Pasighat and Janakmukh on the evening of the 16th, and leaving the latter camp on the 19th we marched to Renging and Rotung, arriving on December 21st.

I was unfortunate in being unable to join the party which left Yembung, the head-quarter camp, on December 27th to explore the course of the Dihang river and survey the country to the north: shortage of transport compelled Major-General Bower to refuse my application. My main object when I joined the expedition was to explore the practically unknown Himalayan fauna of the N.-E. Frontier, but it now became

evident that this idea would have to be abandoned. As events turned out this was not so disadvantageous as was feared, for the fauna of the foot-hills yielded material of very considerable interest and in investigating this region it was not necessary to make any drastic reduction in the apparatus required to carry on the work. But the country visited must be regarded purely as the foot-hill region of the north-east Himalayas and this fact must not be lost sight of in any comparisons which may be drawn between the Abor fauna and that of the ranges further to the west.

We remained at Rotung until January 12th, when we proceeded to the head-quarter camp at Yembung, and a week later I received permission to join a small party with an Indian surveyor under Capt. A. L. M. Molesworth which was going up the right bank of the Dihang to explore the courses of the Siyom and Shimang rivers, two large tributaries of the Dihang. I left with Capt. Molesworth on January 23rd. We reached Parong, a village two marches below Riga, and Damda which is some ten miles up the Siyom river, and returned to Yembung, sooner than was anticipated, on February 3rd. Triangulation did not form part of the survey work on this occasion; there were consequently no halts of any considerable length and the greater part of the time was occupied in hard marching which afforded but little facility for zoological work. Subsequently I proceeded to the Komsing camp, close to the Abor village of that name, and remained there until March 3rd, but the work accomplished during this period was almost entirely anthropological.

On my return to Yembung I proceeded as soon as possible to Rotung, where I purposed staying a few days to make further observations on the zoology of the district. But Capt. Sir George Duff Dunbar, who was then in command of Rotung post, suggested that I should make a short expedition to the west beyond Kalek and the Sireng valley towards Misshing, offering to provide an escort and Abor cooly transport for the purpose. I accepted this offer gladly, and in company with Mr. J. Coggin Brown of the Geological Survey and Dr. Falkiner of the Assam Valley Light Horse, who was then attached to the Lakhimpore Military Police, I left Rotung on March 15th, marching two days towards the west, halting the third day and returning on the evening of the 19th. This expedition gave me an opportunity of visiting fresh country with high tree-jungle of a type not elsewhere seen, a change which was of course accompanied by a corresponding change in the fauna. My thanks are due to Sir George Dunbar for suggesting this expedition and for making every arrangement necessary for its success.

On the morning of March 20th we left Rotung and marched beyond Renging to the camp of the 32nd Pioneers in the Sirpo valley and after a day's halt proceeded to Pasighat, from which place we reached Balek, where three days were spent in anthropological work. Returning to Pasighat we proceeded on March 27th by boat-convoy to Kobo, and leaving there a few days later arrived in Calcutta on the evening of April 3rd.

The Abor country lies in the N. E. corner of the Indian Empire and is bounded on the east by Mishmi country, on the north by Tibet,

on the west by the land inhabited by the Miris and on the south by the Brahmaputra River. Plate I, which is reproduced by permission from the maps made during the expedition by officers of the Survey of India under Capt. O. H. B. Trenchard, R.E., shows the routes which I traversed and the principal places at which collections were made.

In a north and south direction the Abor country extends for about 80 miles.¹ Between the base camp at Kobo and Janakmukh and Balek it consists of an alluvial plain situated at a height of 400 to 600 feet above sea-level and clad with dense jungle interspersed with open patches of long grass or *chapris*. In some places the jungle has at one time or another been cleared for cultivation, but the older clearings are for the most part so thickly grown with scrub and creepers as to be almost impenetrable. To the north of this the country is a maze of hills, often precipitous, and intersected by the boulder-strewn courses of the numberless small streams that drain into the Dihang river. All the hills were originally clothed from foot to summit with tree-jungle, but frequently, more particularly in the vicinity of the Abor villages, large tracts of country have been cleared. Many such areas have, after a period of years, been discarded for cultivation purposes and soon develop thick scrub-jungle.

The majority of the specimens obtained were found at comparatively low altitudes between 400 and 2,500 ft., but some were taken at greater heights up to about 5,000 ft. and part of a small collection of insects, made for me by Mr. J. Coggin Brown of the Geological Survey, was obtained on Geku hill at a height of about 10,000 ft. Papu, the highest of the foot-hills proper, reaches an altitude of 6,390 ft.

As regards invertebrates, the best results were obtained by searching under bark and in rotten wood and large collections were made by this means alone. In particular some dead and partially decayed jack-fruit trees, which were being cut by the 32nd Sikh Pioneers in the vicinity of Rotung, afforded admirable material; many of the branches were hollow and bored and when split open were found to be literally filled with a varied assortment of Carabidae, Staphylinidae, Passalidae, Endomychidae, Heteromera, Rhynchophora, Dermaptera and Blattoidea along with representatives of many other groups. Considerable collections were also made under stones and numerous interesting species were found in this situation.

In the Abor country the cold weather season is also the driest season of the year and numbers of invertebrates, to which an abundance of moisture is a necessity, take refuge during this period in plantain trees, living behind the great ensheathing leaf-stems. Water is invariably found in the leaf-base and the atmosphere in the almost completely closed chamber behind the leaf-stems is probably always at or near saturation point. A considerable amount of rain fell during February and March, and in the latter month it was noticed that the numbers of animals found in such situations had greatly diminished.

¹ Only the southern portion of the Abor country is shown on the map (Pl. I.); the great ranges further to the north were not visited.

Small Dytiscidae, Odonata larvae and Oligochaete worms were found in the tops of plantains and screw-pines (*Pandanus*), living in a collection of water at the leaf-base at from 20 to 40 ft. above the surface of the ground.

No lakes or pools of standing water were met with in the country and practically all the aquatic animals obtained were taken in small streams and rivers draining into the Dihang or its larger tributaries: in the Dihang itself, which in the rains appears to rise in some places at least sixty feet above its winter level, practically nothing could be found. The smaller rivers and streams of the hills teem with the larvae of Neuroptera and Odonata and in the warm weather the perfect insects must occur in very great numbers. Brachyurous Crustacea were not uncommon, but Macrura appeared to be represented only by a single species of *Palaemon*. Fish were plentiful and some, along with tadpoles belonging to the families Ranidae and Pelobatidae, show interesting adaptations to life in hill-streams subject to sudden spate.

The different groups of animals are unfortunately very unevenly represented in the collection, but this was to a large extent unavoidable. The cold weather season is not the best time of year for zoological work and the poverty of the collection in several respects, notably in some sections of the Insecta, must be attributed to this fact. Other groups, again, should have received far more attention than I was able to give them, for only by the devotion of a specialist's whole time to the subject could satisfactory results have been obtained among the birds and mammals.

My activities and opportunities for work were, as was only to be expected, somewhat limited by the restrictions necessary in the case of a military expedition carried out in a hostile country, but it was only in a few instances that these restrictions were severely felt. Work with an escort must of necessity have some drawbacks, though in many cases, more especially with Gurkhas, the escort was keenly interested and showed itself adept in the capture of the more elusive species.

To express my gratitude for all the help I received in the course of the expedition is a difficult task. I have in the first place to thank Major-General Bower, C.B., General Officer commanding the Force, for the interest which he took in my work and for the facilities extended to me.

To the 32nd Sikh Pioneers I am under very great obligation for the large collections of Reptilia and other animals which they made on my behalf. During the winter season most of the reptiles were in hiding beneath the ground or in roots of trees, and in these situations they were found by the Pioneers when cutting roads along the hill-sides. The extent of the help they gave me may be judged from the fact that while snakes are now represented by twenty-six species, three hitherto unknown and one belonging to an undescribed genus, my own individual efforts only resulted in the acquisition of three common forms. Through the instrumentality of the British officers of the regiment and in particular of Capt. the Hon. M. de Courcy each non-commissioned officer on works was provided with a large section of bamboo fitted with a plug

and every day a valuable collection of snakes, lizards, frogs, centipedes, scorpions, etc., was obtained.

Numerous specimens were also contributed by Col. (now Brigadier-General) D. C. F. Macintyre, Capt. A. L. M. Molesworth, by Capt. J. S. O'Neill, Capt. F. H. Stewart and Capt. R. S. Kennedy of the Indian Medical Service, by Capt. H. W. Price and Mr. G. F. T. Oakes. We are indebted to Major Sweet, Capt. B. R. Nicholl, Capt. J. Masters and Mr. I. Burn Murdoch for the loan of various mammals and to Dr. J. Falkner for the loan of a collection of birds.


Opportunity will I hope be found of making fuller individual acknowledgments in the course of publication of the different reports, though it will still be impossible for me to express my thanks in an adequate manner for all the kindness and hospitality I received from officers of different services attached to the expedition.

I was fortunate in having Mr. R. Hodgart with me as my assistant. Throughout the period during which we were attached to the expedition his energy was unremitting and he spared no efforts to render the work a success.

STANLEY KEMP,

June 1st, 1912.

Offg. Superintendent, Indian Museum.



Map of the southern part of the Abor country.

I. BATRACHIA.

(Plates ii-iv.)

By N. ANNANDALE, D.Sc., F.A.S.B., Superintendent, Indian Museum.

Mr. Kemp's collection of Batrachia from the Abor country and the frontiers of Assam comprises 57 specimens of frogs and toads and a considerable number of tadpoles. With those obtained on the Expedition of 1911-1912 I have included two interesting specimens taken by Mr. Kemp on a previous visit to the frontier of eastern Bhutan. In all at least 25 species are represented, of which about one-third are new to science, while several have only been recorded hitherto from Burma or from Assam south of the Brahmaputra. The collection, therefore, affords the opportunity of making important additions to the fauna of the Himalayas, the animals of the extreme eastern region of which have up to the present been almost unknown.

PART I.—SYSTEMATIC.

List of species represented in Mr. Kemp's collection :—

- | | |
|--------------------------------------|---|
| 1. <i>Rana cyanophlyctis</i> . | 14. <i>Rhacophorus maculatus him-</i> |
| 2. <i>R. liebighi</i> . | <i>alayensis</i> , nov. |
| 3. <i>R. tigrina</i> . | 15. <i>Rh. tuberculatus</i> . |
| 4. <i>R. limnocharis</i> . | 16. <i>Ixalus asper</i> . |
| 5. <i>R. alticola</i> . | 17. <i>I. annandalei</i> . |
| 6. <i>R. granulosa</i> . | 18. <i>I. argus</i> , nov. |
| 7. <i>R. afghana</i> . | 19. <i>I. tuberculatus</i> . |
| 8. <i>R. gerbillus</i> , nov. | 20. <i>Chirixalus doriae</i> . |
| 9. <i>Micrixalus borealis</i> , nov. | 21. <i>Phrynoderma moloch</i> , nov. |
| 10. <i>Rhacophorus maximus</i> . | 22. <i>Bufo melanostictus</i> . |
| 11. <i>Rh. bimaculatus</i> . | 23. <i>B. himalayanus</i> . |
| 12. <i>Rh. naso</i> , nov. | 24. <i>Megalophrys</i> (?) <i>major</i> . |
| 13. <i>Rh. microdiscus</i> , nov. | 25. <i>M. kempii</i> , nov. |

(a) ADULTS.

Fam. RANIDAE.

1. *Rana cyanophlyctis*, Schneid.

Boulenger, *Fauna*, p. 442.

This frog occurs all over the plains of India and ascends the Himalayas to altitudes of at least 5,000 ft. In Kumaon in the Western Hima-

layas it is the common frog at 4,500 ft., but at 6,000 ft. is entirely replaced by *R. blanfordii*. It is very largely aquatic in habits and is not as a rule found in dense jungle.

Mr. Kemp obtained a single specimen at Dibrugarh.

2. *Rana liebighii*, Günth.

Boulenger, *Fauna*, p. 445.

R. liebighii is very common in the Darjiling Himalayas at altitudes between 4,000 and 10,000 feet; it also occurs, though rarely, in the Western Himalayas (Simla and Kashmir), while to the south-east its range extends to northern Tenasserim. It is essentially a jungle frog and is usually found among dense undergrowth or at the edge of streams.

No adults of this species were obtained on the expedition, but two tadpoles were taken in a small stream near Yembung (alt. 1,100 ft.) on the east side of the Dihang River.

3. *Rana tigrina*, Daud.

Boulenger, *Fauna*, p. 449.

Although it is apparently rare in the Himalayas, *R. tigrina* occurs all over the plains of India. In different parts of India, however, its habits differ considerably and certain structural differences also seem to occur, so that it is probable that several different races will ultimately have to be recognized as distinct. In Bengal *R. tigrina* is essentially a "tank" frog, inhabiting moderately large masses of water.

Two specimens were obtained by Mr. Kemp at Sadiya under a log at the edge of the Dikrang River.

4. *Rana limnocharis*, Wiegman.

Boulenger, *Fauna*, p. 450.

Being much more adaptable in its habits than *R. tigrina*, *R. limnocharis* has an even wider range than that species. It is equally at home in flooded rice-fields and at the edge of rocky streamlets in the densest jungle. Both in the Himalayas and in the mountains of Burma it ascends to an altitude of at least 6,000 ft.

Mr. Kemp obtained specimens at Sadiya, Kobo and Rotung and in the Siyom valley below Damda.

5. *Rana alticola*, Boulgr.

Boulenger, *Cat. Batr. Sal. Brit. Mus.* (2nd ed.), p. 63, fig. (1882).

The range of this frog is considerable both in longitude and latitude and also in altitude. Colonel Alcock obtained specimens of the charac-

teristic tadpole at an altitude of 8,500 ft. in the Hundur Yarm Valley in northern Kashmir, while Capt. R. B. Seymour Sewell has recently taken precisely similar larvae a few feet above sea-level and a few yards from the shore, in a small stream running into Heinze Basin on the coast of Tavoy. I have also examined larvae from other parts of Tenasserim and Assam and from Little Andaman I., and adults from Tenasserim and Assam, Bengal and Orissa. I have not, however, seen specimens from any place in the plains west of Calcutta. In Lower Burma and Orissa the frog is usually found amongst dense vegetation at the edge of ponds and lakes. On the shores of the Sar Lake near Puri it is abundant on the leaves of plants that grow out of the water and also on ledges in the sides of old wells.

The back of the young frog is brown, often mottled with black, but in the adult it becomes of a bright leaf-green. I have redescribed the tadpole below (p. 22). *R. alticola* is of much more slender habit and usually of smaller size than *R. erythraea*, which resembles it in habits, but apparently is not found west of the Bay of Bengal.

Mr. Kemp took a small specimen of *R. alticola* on the Assam-Bhutan frontier in the north-east of the Mangaldai division of the Darrang district in January, 1911.

6. *Rana granulosa* (Anderson).

Hylorana granulosa, Anderson, *J.A.S.B.*, (2), xl, p. 23 (1871).

?*Rana tyleri*, Boulenger, *Fauna*, p. 458 (*partim*).

Rana granulosa, *id.*, *Ann. Mus. Genova* (2nd ser.), xiii, p. 333, pl. viii, fig. 2 (1893).

This frog is known from the Karin Hills and Pegu in Burma, from Yunnan and from north-eastern Assam (Sibsagar). It is apparently arboreal in habits.

A specimen was taken on a tree-trunk near Dibrugarh.

7. *Rana afghana* (Günth.).

Polypedates afghana, Günther, *Rept. Brit. Ind.*, p. 432.

Rana latopalmata, Boulenger, *Fauna*, p. 462.

Rana afghana, *id.*, *Ann. Mus. Genova* (2nd ser.), v, p. 420 (1888).

It is very unfortunate that this frog should have to bear the inappropriate specific name "*afghana*," for there can be little doubt that its range does not extend west of Nepal. It is not uncommon in the Darjiling Himalayas at moderate altitudes and has also been found in Assam and Burma.

An adult was taken by Mr. Kemp at Yembung (alt. 1,100 ft.), and also two tadpoles, which are noticed below (p. 24).

8. *Rana gerbillus*, sp. nov.

(Plate ii, fig. 1.)

Allied to *R. jerboa* (Günther), from which it differs in its much smaller tympanum and in other particulars.

Habit slender. Length from snout to vent 33 mm.

Head broad, triangular; snout bluntly pointed, somewhat depressed at the tip, a trifle longer than the diameter of the orbit; nostril a little nearer the tip of the snout than the eye, which is large and prominent; interorbital space flat, as wide as the upper eyelid; canthus rostralis bluntly angular; loreal region concave; tympanum not very distinct, small, about $\frac{1}{3}$ as wide as eye.

Mouth.—A distinct tooth at the tip of the lower jaw; no free papilla on the tongue; vomerine teeth ill-developed, in two small roundish patches situated close together in the middle of the palate between the choanae.

Limbs slender. Fingers slender, with well-developed disks; that on the third rather larger than the tympanum; that on the first small; a rudiment of a web between the third and fourth fingers; others quite free; first finger shorter than second. Hind limbs very long, the tarsotibial articulation reaching far beyond the edge of the snout. Toes almost completely webbed; their disks subequal, smaller than the tympanum; subarticular tubercles large but not prominent, oval; a low oval inner metatarsal tubercle; no outer one; no tarsal fold.

Skin.—A distinct glandular latero-dorsal fold and another, less distinct, extending from the eye above the tympanum almost to the shoulder. Dorsal surface of the head minutely pitted, of the back obscurely granular with large compressed longitudinal tubercles scattered more especially on the sides. Ventral surface and limbs smooth.

Colouration.—Dorsal surface very dark grey obscurely mottled with a paler shade; lips with pale vertical stripes; sides pale, spotted with dark grey; limbs, especially the thighs, conspicuously barred; ventral surface dull greenish yellow with large round or oval brown spots on the chin, throat and chest.

Habitat.—Yembung, Abor foot-hills at an altitude of 1,100 ft. A single specimen was found at the edge of a small stream in February.

Type.—No. 16925 of Indian Museum register of Reptiles and Batrachia.

Rana jerboa (Günther), to which this new species is most closely allied, is found in Burma, the Malay Peninsula, Java, Borneo, etc.

9. *Micrixalus borealis*, sp. nov.

(Plate ii, fig. 2.)

Closely allied to *Rana tenasserimensis*,¹ Sclater, from which it may be distinguished by the small size of the disks on its fingers, its hidden tympanum and its almost completely webbed toes.

¹ P. Z. S., 1892, p. 345, pl. xxiv, fig. 4, and *List Batrachia Ind. Mus.*, p. 8 (1892).

Habit stout. Length from snout to vent 25 mm.

Head short, broad, triangular, by no means depressed; snout bluntly pointed, convex at the tip, a little shorter than the orbit; canthus rostralis rounded; loreal region not concave, sloping outwards and downwards; eye large, not very prominent; interorbital region slightly convex, a little broader than the upper eyelid; tympanum concealed; a prominent fold running backwards and downwards from the eye to a point a little behind the gape.

Mouth.—A prominent tooth at the apex of the lower jaw; no trace of vomerine teeth; the choanae situated far forward; no papilla on the tongue. An internal vocal pouch in the male.

Skin.—No dorso-lateral fold. Dorsal surface minutely warty; ventral surface almost smooth but with a rather indistinct reticulation of grooves; a transverse fold running across the posterior end of the throat in the male.

Limbs stout. Fore limbs short; fingers short; the first a little shorter than the second; disks very small but distinct; a rudiment of a web between all the fingers; subarticular tubercles inconspicuous but of fairly large size. Tibio-tarsal articulation reaching the eye; toes moderately slender, with very small disks; web almost complete but not quite reaching or barely reaching the disks of the first and fifth toes; subarticular tubercles inconspicuous; an elongate but not very prominent internal metatarsal tubercle followed by a fold of skin on the tarsus; another fold on the external margin of the foot; no outer metatarsal tubercle.

Colouration.—Back dark brown mottled with purplish black; a dark interorbital cross-bar usually present; limbs obscurely barred and digits more conspicuously so; ventral surface yellowish, powdered on the throat and chin with purple-brown, as a rule so densely that the skin appears almost black to the naked eye; ventral surface of hands, feet and thighs powdered in a similar manner but not so densely; belly sometimes spotted.

Localities.—Rotung (alt. 1,300 ft.) and about 3 miles S. of Yembung.

Type.—No. 16932 in Indian Museum register of Reptiles and Batrachia.

I think I am right in referring this species, of which I have examined eleven specimens, to Boulenger's genus *Micrixalus* rather than to *Rana*; for I can find no trace of vomerine teeth, while the small size of the frog and its general facies are features in which it resembles the south Indian species included in the former genus. Sclater's *Rana tenasserimensis*, of which three type-specimens are now in our collection, has apparently vomerine teeth in some individuals and none in others, but these teeth are never well developed. It should also, in my opinion, rank as *Micrixalus*.¹

¹ See Boulenger, *P.Z.S.*, 1883, p. 205, and *Fauna*, p. 464; also W. L. Sclater, *P.Z.S.*, 1892, p. 345, pl. xxiv, fig. 4, 4a, and Boulenger, *Ann. Mus. Genova* (2nd ser.), xiii, p. 331.

Ten specimens of *M. borealis* were taken under stones in a stream at Rotung and one a few miles S. of Yembung.

10. *Rhacophorus maximus*, Günth.

Boulenger, *Fauna*, p. 472.

A common species on the lower slopes of the Darjiling Himalayas and the Khasi Hills. Colonel Godwin-Austen obtained specimens in the Dafla country.

Mr. Kemp obtained a half-grown specimen at Upper Rotung at an altitude of about 2,000 ft.

11. *Rhacophorus bimaculatus*, Boulgr.

Boulenger, *Fauna*, p. 472.

Not uncommon in the Khasi Hills. Mr. Kemp's only specimen, which was taken at Rotung (alt. 1,300 ft.) in January, has the web of the feet of a bright scarlet colour and lacks the dark spots on the sides usually so characteristic of the species. In the former particular it agrees with other specimens in our collection.

12. *Rhacophorus naso*, sp. nov.

(Plate ii, fig. 3.)

This peculiar species can be distinguished from any other of the genus that occurs in Assam or Burma by the dermal appendage on its snout.

Habit moderately stout. Length from snout to vent 43 mm.

Head rather broad, triangular, with convex sides; snout much longer than orbit, pointed, convex above, nostril much nearer tip of snout than eye; canthus rostralis indistinct; loreal region concave, almost horizontal. Diameter of tympanum, which is distinct, about $\frac{2}{3}$ that of eye.

Mouth.—No prominent tooth at apex of lower jaw; no papilla on the tongue; vomerine teeth forming two small, almost circular patches, one close to the inner margin of each choana; choanae small.

Skin of dorsal surface rugose with many prominent and irregular tubercles; a small subquadrangular dermal projection on the snout; rounded tubercles scattered on the basal part of the thighs; ventral surface coarsely granular, more so on the chest and throat than on the belly. A fairly distinct dorso-lateral fold and a more prominent one running from the eye above the tympanum to the shoulder; serrated cutaneous fringes on the outer edges of the forearm and the shin.

Limbs stout, not very long. Fingers with a rudimentary web; their disks well developed, transversely oval; that on the third finger almost as large as the tympanum; subarticular tubercles well developed. The tibio-tarsal articulation reaches the eye; toes almost completely

webbed ; their disks like those of the fingers ; subarticular tubercles well developed ; a somewhat elongate but not very prominent inner metatarsal tubercle ; no corresponding outer tubercle.

Colouration.—Dorsal surface purplish-brown irregularly marked with dark slate-colour ; irregular paler markings on the sides ; limbs irregularly marked ; fingers and toes barred with alternate brown and grey stripes. Ventral surface dirty white, shaded posteriorly with dark grey ; dark grey spots on throat and an irregular reticulation of the same shade on the chest. Ventral surface of hands and feet grey, with the tubercles white ; a white spot on the soles in the place where an external metatarsal tubercle would be.

Habitat.—Egar stream between Renging and Rotung (9-i-12).

Type (a unique specimen). No. 16929 in the Indian Museum register of Reptiles and Batrachia.

13. *Rhacophorus microdiscus*, sp. nov.

(Plate ii, fig. 4.)

This species is easily recognized by the small size of its digital disks.

Habit slender. Length from snout to vent 29 mm.

Head large, flat, broadly ovoid ; snout rounded in front, by no means prominent, rather shorter than the orbit ; nostril much nearer tip of snout than eye ; canthus rostralis indistinct ; loreal region concave, oblique ; tympanum distinct, close to eye ; its diameter about $\frac{1}{3}$ of that of eye ; interorbital space about as wide as upper eye-lid, flat.

Mouth.—No tooth at apex of lower jaw ; no papilla on the tongue ; choanae large ; vomerine teeth in two short, ridge-like series running obliquely backwards and inwards from a point near the anterior inner margin of the choanae but widely separated from one another.

Limbs slender but short. Fingers with a slight rudiment of a web ; their disks very small, that on the third being much less than $\frac{1}{2}$ as wide as the tympanum ; subarticular tubercles well developed ; a large rounded inner metacarpal tubercle. Tibio-tarsal articulation barely reaching the eye ; disks of toes like those of fingers ; feet less than half webbed ; subarticular tubercles well developed ; an elongate but by no means prominent inner metatarsal tubercle.

Skin of dorsal surface, throat and chest smooth ; that of belly separated into polygonal areas by a very distinct network of grooves ; on the posterior part these areas gradually take the form of low conical tubercles ; isolated rounded tubercles on basal part of thighs. No dorso-lateral fold ; an indistinct fold running from above tympanum to shoulder.

Colouration.—Dorsal surface pale slate-grey irregularly marked with darker grey and powdered with black ; dorsal surface of limbs

indistinctly barred; outer margin of thigh reddish; ventral surface dirty white.

Habitat.—Kobo, at base of Abor foot-hills (alt. 400 ft.) : 29-30-iii-12.

Type (a unique specimen). No. 16924 in Indian Museum register of Reptiles and Batrachia.

14. *Rhacophorus maculatus* (Günther).

Rhacophorus maculatus and *R. leucomystax*, Boulenger, *Fauna*, pp. 474, 475.

I have little doubt that what may be called the "Common Tree-frog" of Peninsular India, the Himalayas and the Malay Peninsula really represents three local races of a single species. These three races or subspecies may be distinguished as follows :—

1. *Rhacophorus maculatus* (Günther) (*forma typica*). No parieto-squamosal arch; dorsal surface of skull smooth; skin of dorsal surface of head free. *Distribution*.—Peninsular India and Ceylon.

2. *Rhacophorus maculatus himalayensis*, subsp. nov. A well-developed parieto-squamosal arch; dorsal surface of skull smooth; skin of dorsal surface of head free. *Distribution*.—The Eastern Himalayas, Assam, western China.

3. *Rhacophorus maculatus leucomystax* (Gravenhagen). A well-developed parieto-squamosal arch; dorsal surface of skull rugose; skin of dorsal surface of head adhering to the skull. *Distribution*.—Lower Burma, the Malay Peninsula and many of the Malay islands.

Mr. Kemp's specimens belong to the second race. The larvae of the three races are discussed below (p. 24).

Two small specimens were taken at Kobo and a very large one on the east side of the Dihang R. at an altitude of 1,100 feet.

15. *Rhacophorus tuberculatus*, Anderson.

(Plate ii, fig. 5.)

Anderson, *J.A.S.B.*, (2), xl, p. 26; Boulenger, *Fauna*, p. 474.

The specimens in Mr. Kemp's collection agree well as regards structure and dimensions with the late Dr. Anderson's description and with his type specimens, which are in the same condition as they were when the species was described. The colours of the latter specimens had, however, already faded at that date and the fresh ones now before me are particularly interesting in this respect. No two of the four.

brought back from the Abor country and taken together in circumstances which I will describe immediately, are precisely identical in colouration ; but in all the colours are so blended that those commonly found on the stems of bamboos growing in thickets in damp jungle are accurately reproduced. The back and the dorsal surface of the head and limbs are in all dull clay-colour sparingly powdered with black and suffused more or less definitely with yellow, the canthus rostralis is outlined in black, the ventral surface is pale yellow and the inner surface of the thighs wholly or partly scarlet. In one individual there is a narrow black, white-edged line running backwards along each side from the eye to the base of the hind limb, while the back of the head and the back are ornamented with two large irregular marks outlined by similar lines. In this specimen, which appears to be an adult male, there are also dark longitudinal lines on the limbs and on the inner margin of the fifth toe and the web that intervenes between that toe and the fourth. In the second individual, a female, the sides of the back are so strongly suffused with yellow that they may be described as ochraceous, the webs of the feet are almost black and the red colour of the inner surface of the thigh extends down that of the shin. In another male, rather smaller than the first, there are small black spots on the back and a dark-edge white line running transversely on the dorsal surface above the vent. The fourth specimen, probably a young male, has no very definite markings. The colouration of the dorsal surface in all the specimens bears a close resemblance to that of a bamboo stem overgrown with minute fungi and lichens such as are usually found on bamboo-stems in a very damp atmosphere. The scarlet of the thighs would be completely concealed in the attitude of rest.

It is interesting to have the opportunity of comparing Anderson's types with fresh specimens of the species, but it is still more interesting to be able to put on record the peculiar circumstances in which these specimens were taken. They were found in an internode of bamboo which was intact as regards both its sides and its two nodes, except that there was in one side a small hole apparently made by some insect, less than a quarter of an inch in diameter. When the bamboo was split open in preparation for bridge-making the four frogs were seen seated on the inner surface near one end (the upper at the time), while two earthworms and a land planarian occupied the other. There can be little doubt, therefore, that they had entered the bamboo as small frogs and had been supplied with food by the intrusion of worms and other small animals through the hole by which they had originally entered, and from which their increase in bulk rendered it impossible for them to emerge. In spite of the fact that they must have lived for some considerable time practically in the dark they had preserved their colouration, which was of a distinctly protective type.

Four specimens from Upper Rotung (alt. *ca.* 2,000 ft.) : 22nd January 1912, collected by Capt. the Hon. M. de Courcy. Anderson's specimens were from Sibsagar in N.-E. Assam. This place is situated on the northern bank of one of the smaller tributaries of the Brahmaputra.

16. *Ixalus asper*, Boulgr.

Boulenger, *P.Z.S.*, 1886, p. 415, pl. xxxix, and *Ann. Mus. Genova* (2nd ser.), xiii, p. 340, 1893 ; Sclater, *P.Z.S.*, 1892, p. 347 ; Robinson, *Journ. F.M.S. Mus.*, i, p. 24.

This species appears to be the most widely distributed of the Indian *Ixali*. It was originally described from the mountains of Perak in the Malay Peninsula and was found by the late Signor Fea in the Karin Hills and by one of our collectors in the hills between Burma and Siam. The Indian Museum possesses a specimen labelled as being from Kolasi in the Purnea district of Bihar, but this specimen very possibly came actually from the foot-hills of eastern Nepal.

The irregular white or greyish-white markings on the posterior part of the body of *I. asper* give it exactly the appearance of being overgrown with a mould or fungus. Whether this is of any protective advantage to the frog may perhaps be doubted, but I have noticed a similar phenomenon in the case of several Malayan insects belonging to the orders Coleoptera and Rhynchota and it is well known that in tropical jungles insects are frequently attacked by fungi which produce a white mycelium and finally, having killed them, fasten their dead bodies by means of this mycelium to tree-trunks or other inanimate objects.

A specimen of *Ixalus asper* was taken on a tree-trunk at the edge of Egar stream between Renging and Rotung on 9th January, 1912. With it were taken the type specimen of *Rhacophorus naso*, a specimen of *Ixalus tuberculatus* and also tadpoles of two species, a *Megalophrys* (*M. ? major*) and a Ranid which cannot be identified.

17. *Ixalus annandalei*, Boulgr.

(Plate iii, fig. 2.)

Boulenger, *J.A.S.B.*, 1906 (2), p. 335.

This species was not taken on the Abor Expedition, but a single specimen was obtained by Mr. Kemp in December, 1910, on the Bhutan frontier of Assam in the Mangaldai division of the Darrang district. It is common in the Darjiling district between 4,000 and 5,000 ft. and occurs both among dead leaves and low herbage in the jungle and on tea-bushes.

18. *Ixalus argus*, sp. nov.

(Plate iii, fig. 3.)

Habit moderately-slender, *Rhacophorus*-like. Length from snout to vent 27 mm.

Head short and broad, triangular ; snout blunt, somewhat depressed at tip, obliquely truncate vertically, projecting, a little longer than the orbit ; nostril nearer tip of snout than eye ; canthus rostralis fairly distinct ; loreal region vertical, concave ; tympanum distinct, small, about $\frac{1}{3}$ as broad as eye, interorbital space broader than upper eye-lid.

Mouth.—A small tooth at apex of lower jaw ; no papilla on the tongue ; choanae small, widely separated, situated far forwards.

Skin.—No latero-dorsal or supratympanic folds. Skin of head and neck smooth, of back beset with small scattered tubercles ; ventral surface smooth.

Limbs slender. Fingers slender, free, with large disks ; that on third finger as large as tympanum ; first finger a little shorter than second ; subarticular tubercles large, rounded, a little prominent ; no metacarpal tubercles. Tibio-tarsal articulation reaching nostril ; toes fully webbed ; disks a little smaller than those on fingers ; subarticular tubercles distinct but by no means prominent ; a very distinct inner, but no outer, metatarsal tubercle ; no tarsal fold.

Colouration.—Dorsal surface dark slate-grey with a paler reticulation on the back that gives it the appearance of being faintly ocellated ; limbs and fingers conspicuously barred with dark grey and white ; ventral surface dirty white faintly spotted on the throat and chest with grey ; soles of feet and palms of hands dark grey.

Habitat.—Upper Renging, alt. 2,150 feet.

Type (a unique specimen). No. 16950 in the Indian Museum register of Reptiles and Batrachia.

19. *Ixalus tuberculatus*, Anderson.

(Plate iii, fig. 1.)

Anderson, *Anat. Zool. Res. Yunnan Exp.*, p. 855, pl. lxxviii, fig. 7 (1878).

I think I am right in identifying a series of specimens in Mr. Kemp's collection with this species, but Anderson's figure is poor and his specimens are not forthcoming for comparison. It is evidently variable in several characters, notably in stoutness of habit, in colouration, in roughness of skin and in the relative size of the disks of the fingers and toes. In some of the Abor specimens the skin of the dorsal surface is smooth except for small scattered tubercles, in others the tubercles are so much larger and closer together that it is quite rough ; in some the finger-disks are much smaller than the toe-disks, but in others they are of almost exactly the same size, while some individuals are much stouter than others. In all the ground-colour of the dorsal surface is very dark slate-grey rather than olive, but this may be due to the fact that the specimens had been hardened in formalin before being preserved in spirit. A pale cross-bar between the eyes can usually be detected and there is always a pale patch on the sides near the groin marked diversely with black ; the corresponding surface of the outer margin of the thighs is similarly marked ; in some individuals there is a large dark x-shaped mark on the back.

Specimens were taken at Janakmukh (600 ft.), in Egar stream between Renging and Rotung, at Rotung (1,300 ft.) and at Kalek (3,800 ft.). At the two last-named localities several individuals were found between the 24th and 29th of December hiding under the leaf-stems

of banana-trees. Probably they were hibernating. In all seven specimens were found.

20. *Chirixalus doriae*, Boulgr.

Boulenger, *Ann. Mus. Genova* (2nd ser.), xiii, p. 341, pl. x, fig. 5 (1893).

A single male taken under a log at Kobo (alt. 400 ft.) agrees well with Mr. Boulenger's figure and description. *Ch. doriae*, which is the only known species of the genus, was originally found by the late Signor Fea in the Karin Hills.

21. *Phrynoderma moloch*, sp. nov.

(Plate iii, fig. 4.)

This species differs from *Phrynoderma asperum*, Boulenger,¹ the only one hitherto known, in several important characters, notably in the much more pronounced nature of the asperities on its back.

Habit slender. Length from snout to vent 41 mm.

Head short, broad, triangular, depressed; snout about as long as orbit, sinuously truncate transversely, obliquely truncate vertically; nostril close to tip of snout, very prominent; eye large, prominent; canthus rostralis indistinct; loreal region almost vertical, slightly concave; inter-orbital region slightly concave, broader than upper eyelid; tympanum fairly distinct, about $\frac{2}{3}$ as wide as eye.

Skin.—No supratympanic or dorso-lateral folds; back bearing very prominent ridge-like, more or less serrated, warts which run longitudinally and obliquely; these warts larger on neck and across shoulders, on which they form V-shaped figures; shorter warts on head and dorsal surface of limbs; throat and ventral surface of limbs smooth; belly and sides coarsely granular.

Limbs slender but rather short. Fingers free, slender, flattened; disks very large, that on the 3rd finger nearly equalling the tympanum; subarticular tubercles small; external and internal subcarpal tubercles indistinct. Tibio-tarsal articulation reaching tympanum; toes $\frac{1}{2}$ webbed, the web reaching the disks of all but the 4th, up which it extends as a narrow fringe almost to the disk; subarticular tubercles small and by no means prominent; a very small and indistinct internal metatarsal tubercle; no external metatarsal tubercle; a narrow serrated fringe on the 5th toe and a less distinct serrated ridge running along middle of ventral surface of shin.

Colouration.—Back grey with black spots; the larger warts buff; the tympanum black; a large black and white diversified patch on each side between the two fore and hind limbs and a white patch in the axilla; external surface of thigh irregularly banded and marbled with black, white and grey; ventral surface black with a faint vermicular

¹ *Ann. Mus. Genova* (2nd ser.), xiii, p. 342, pl. xi, fig. 1.

reticulation running all over the body and onto the ventral surface of the thighs.

Habitat.—Upper Renging (alt. 2,150 ft.) : 5-10-ii-12.

Types (two adults). Nos. 16951 and 16952 : presented by Capt. the Hon. M. de Courcy.

The circumstances in which these frogs and their tadpoles were found are of considerable interest. Capt. de Courcy writes about them as follows :—

“ [The frogs were taken] between Upper Renging and the Yernu Kotal. It was a few yards this side of Prospect Col that some of my men found the new *Phrynoderma* frogs—3 of them, under a log—and kept them, trying to make me see them among some lumps of earth—almost an impossibility. One escaped a few minutes after I had taken them over, and while I was standing there, the men cut off a big bit of the same log, the usual old felled tree on a *jhoom* [clearing], and rolled it down on to the road. Some water gushed out of a hole and I saw the tadpoles wriggling about on the ground and collected all I could.”

The tadpoles, which are described below (p. 25), evidently belong to the same species as the adult frogs, for one of them has progressed far in its metamorphosis and has begun to develop the characteristic ridge-like warts on the back.

Fam. BUFONIDAE.

22. *Bufo melanostictus*, Schneid.

A typical specimen of this toad was taken by Mr. Kemp at Dibrugarh in the middle of November and on the same date he found a number of tadpoles in which the hind limbs were not developed.

23. *Bufo himalayanus*, Günth.

Bufo melanostictus var. *himalayanus*, Günther, *Rept. Brit. Ind.* p. 442.

Bufo himalayanus, Boulenger, *Fauna*, p. 505.

I am inclined to agree with Dr. Günther in regarding this form merely as an Alpine race of *B. melanostictus*. The greatest difficulty is often experienced in separating specimens and quite typical individuals of *B. melanostictus* are often found at considerable altitudes in the Himalayas. Almost every gradation between the two forms can be found. Tadpoles (plate iv, fig. 7), however, from above 4,000 ft. in the E. Himalayas can, so far as my experience goes, be distinguished from those found in the plains of India by the fact that the eyes are not prominent but rather sunken. Tadpoles from the plains agree well with one from the Malay Peninsula figured by Flower (*P.Z.S.*, 1896, p. 911, pl. xlv, fig. 3), and I have found similar specimens at an altitude of over 7,000 ft. in the W. Himalayas near Naini Tal.

Mr. Kemp obtained four toads at Kobo in November and December, which I assign to Günther's "variety" with some doubt. Their tympana

are smaller than is usually the case in *B. melanostictus* and in two of them are also somewhat obscured. The parietal ridges are absent in one specimen and in the others, although they are present, they are very indistinct.

Fam. PELOBATIDÆ.

The genus *Megalophrys* is represented in the collection by tadpoles which seem to belong to at least two species and also by a single small frog, which I have accepted as the type of a new species. The tadpoles are discussed below (p. 28).

25. *Megalophrys kempii*, sp. nov.

(Plate iii, fig. 5.)

Although the only specimen obtained is very small, I am inclined to think from its general appearance that it is at least sub-adult. The species is clearly related to *M. heteropus* (Boulenger),¹ but the snout is rather longer, the tympanum larger and the hind legs longer.

Habit slender. Length (of type) from snout to vent 15 mm.

Head moderate, little depressed; snout longer than eye, sloping forwards above, rounded at the tip, truncate vertically, projecting very little beyond lower jaw; nostril much nearer tip of snout than eye; canthus rostralis distinct; loreal region concave, nearly vertical; inter-orbital distance much greater than width of upper eye-lid; tympanum fully exposed, nearly as large as eye, close to which it is situated.

Mouth.—No vomerine teeth; tongue slightly cleft behind.

Skin of back, limbs, throat and chest smooth, of belly obscurely granular; some irregular tubercles on the base of the thighs. No dorso-lateral or supratympanic folds.

Limbs slender. Fingers short, free, bearing (except the first, which is merely blunt) very small but distinct disks; first finger shorter than second; lower surface of hand smooth; subarticular tubercles poorly developed; no metacarpal tubercles. Hind limb long; the tibio-metatarsal articulation reaches the tip of the snout; toes slender, the fourth very long, with small disks and the rudiments of a web; sub-articular tubercles poorly developed; no metatarsal tubercles; no tarsal fold.

Colouration.—Dorsal surface dark olive with more or less symmetrically arranged greyish-green marks; upper surface of snout of latter shade; hind limbs obscurely banded with dark olive; ventral surface yellowish, suffused with dark olive; throat densely powdered with that shade but ornamented with clear yellowish spots.

Habitat.—Upper Rotung (2,000 ft.): i-i-12.

Type.—No. 17013 in Indian Museum register of Reptiles and Batrachia.

¹ *Ann. Mag. Nat. Hist.*, (7), vi, p. 186 (1900), and *P.Z.S.*, 1908 (1), p. 422, fig. 2.

(b) TADPOLES.

The tadpoles here discussed are those of :—

- | | |
|-------------------------------------|-----------------------------------|
| A. <i>Rana liebigii</i> , | D. <i>Rhacophorus maculatus</i> , |
| B. <i>Rana alticola</i> , | E. <i>Phrynoderma moloch</i> , |
| C. <i>Rana afghana</i> , | F. An unidentified Ranid, |
| G. and H. <i>Megalophrys</i> , spp. | |

Fam. RANIDAE.

A. Larva of RANA LIEBIGII, Günth. (Plate iv, fig. 2.)

Annandale, *J.A.S.B.*, 1906 (2), p. 290.

The tadpole of this species is very abundant in small jungle streams in the neighbourhood of Kurseong (4,000—5,000 ft.), E. Himalayas, both in April and May and in July, August and September. Mr. Kemp obtained two specimens in a small stream on the east side of the Dihang R. near Yembung between the 13th and the 17th of January. Their hind legs were just beginning to appear.

The larva of *R. liebigii* may be redescribed as follows :—

Head and body feebly arched above but not quite flat, sloping slightly from behind forwards, convex on ventral surface, broadly ovoid as seen from above ; the tip of the snout rounded.

Mouth ventral ; lips broad but not excessively so, directed inwards and enclosing a considerable cavity ; posterior lip directed backwards, anterior lip forwards and inwards, thus forming a vestibule to the mouth ; lower lip with a complete double row of rather elongate tubercles ; upper lip fringed with a single row of rather smaller tubercles which is widely interrupted in the middle ; dental formula usually $2 : 5 + 5 \quad 1 + 1 : 2$,¹ sometimes three uninterrupted rows of teeth present on the upper lip ; neither upper nor lower beak divided ; both horseshoe-shaped and minutely serrated ; upper lip very narrow.

Nostril and eye small, by no means prominent ; the former situated midway between the eye and the tip of the snout ; eye directed obliquely upwards, situated at the junction of the dorsal and the ventral surfaces, much nearer the tip of the snout than the base of the tail.

Glands.—None apparent.

Spiracle sinistral, pointing backwards and slightly upwards, small, circular, not markedly tubular.

Vent dextral.

Tail gradually pointed, tapering, twice as long as head and body ; its greatest depth $\frac{1}{5}$ of total maximum length ; fin-membranes well

¹ For convenience in printing this and other dental formulae in this paper are printed in a somewhat novel form. The figures to the right of the central dark vertical line represent the tooth-rows of the upper, those to the left of this line the tooth-rows of the lower lip. The colons merely separate the number of undivided from that of divided rows.

developed for its whole length both above and below the muscular portion ; the greatest depth twice that of the muscular portion.

Colouration somewhat variable ; dorsal surface brownish, marked in some individuals with yellow ; fin-membrane pale, with large dark pigment-cells which in some individuals tend to be arranged in vertical bars ; a dull yellowish mid-dorsal streak sometimes present at base of tail ; spiracle surrounded by a white ring.

Dimensions of an individual in which the hind legs are beginning to appear :—

Total length	..	44 mm.
Length of head and body		15 „
Length of tail	..	29 „
Maximum breadth of body		12 „
Maximum depth of body		10 „
Maximum depth of tail		12 „
Full-grown tadpoles measure about 56 mm. in length.		

B. Larva of RANA ALTICOLA, Boulgr. (Plate iv, fig. 1.)

Boulenger, *Cat. Batr. Sal. Brit. Mus.*, p. 62, fig. ; Annandale *P.Z.S.*, 1905 (I), p. 58, pl. vi.

The tadpoles figured and described by Mr. Boulenger were evidently badly preserved and faded. I have, therefore, re-described this larva from well-preserved and recently captured specimens.

Head and body flattened both above and also on the anterior part of the ventral surface, oval, truncate anteriorly.

Mouth ventral ; lips well developed, the posterior lip directed backwards, not excessively broad, with a single row of rather small tubercles running all along its margin ; anterior lip directed forwards and inwards, fringed for rather less than a third of its length on either side and bearing on its ventral surface in the same region numerous small tubercles, bare in the middle. Dental formula 2 : 5+5 1+1 : 8, the outer row of teeth on the posterior lip feebly developed ; beak in two parts, an upper and a lower ; both parts roughened on the surface but not serrated at the margin ; upper part crescentic, projecting slightly in the middle ; lower part broadly V-shaped.

Nostril and eye small, by no means prominent ; the former in well-preserved specimens rather nearer the eye than the tip of the snout ; eye directed obliquely upwards, situated near the dorsal surface, much nearer the snout than the base of the tail.

Spiracle sinistral, tubular, directed outwards and a little upwards.

Vent dextral.

Glands.—A large and prominent oval parotoid gland¹ present on each side and a small and less conspicuous single gland (in large tadpoles) on the dorsum at the base of the tail.

Tail bluntly pointed, about $1\frac{1}{3}$ times as long as head and body, very shallow at its base owing to poor development of both fin-membranes; these become deep shortly afterwards and then diminish again somewhat abruptly, so that the outline of the tail is strongly sinuous.

Colouration.—In the young tadpole the head and body are boldly diversified with dark and pale markings not of a symmetrical nature, while the muscular part of the tail bears numerous large and small ocelli, which are replaced on the fin-membranes by small black spots. As the tadpole grows and the limbs begin to develop the colours darken and the caudal ocelli become indistinct or disappear, with the exception, as a rule, of one large ocellus on each side at the base of the tail. Sometimes a second smaller and more distal ocellus also persists and occasionally there is a row of ocelli all along the tail of even full-grown larvae, gradually diminishing in size from in front backwards. The central spot of each ocellus is black, the outer ring yellow. The number of ocelli is not always the same on both sides of the body.

Dimensions.—The tadpole reaches a length of at least 57 mm. The following are the measurements of two specimens in one of which (A) the hind limbs appear as minute buds, while in the other (B) the toes can just be detected :—

	A (tail injured).	B.
Total length	— mm.	42 mm.
Length of head and body	27 „	18 „
Length of tail	— „	24 „
Maximum breadth of body	16 „	10 „
Maximum depth of body	14 „	9 „
Maximum depth of tail	16 „	9 „

I was at first inclined to adopt the opinion that the large series of ocellate Ranid larvae in our collection represented two distinct species, one with a distinct supra-caudal gland, prominent parotoids and not more than two ocelli on each side of the tail, the other with no supra-caudal gland, much less prominent parotoids, the head and body spotted and mottled, and numerous caudal ocelli. Specimens, however, recently obtained by Mr. F. H. Gravely and Capt. R. B. Seymour Sewell, I.M.S., in Lower Burma, show that there is a complete gradation between the two forms, the latter being merely a younger stage of the former. Mr. Gravely's specimens, in none of which were the hind limbs developed, were taken in a small pond near Kawkarcik in November and Capt.

¹ This gland is not nearly so conspicuous in fresh specimens as it is in old and faded ones.

Sewell's, which were in a slightly more advanced stage of development, on the coast of Tavoy in spring. A young frog which had already assumed the green back of the adult was taken with Mr. Gravely's tadpoles.

C. Larva of *RANA AFGHANA* (Gunth.) (Plate iv, fig. 3.)

? *Rhacophorus reinwardtii* (Larva), Boulenger, *Cat. Batr. Sal. Brit. Mus.*, p. 89, fig.

Rana afghana, *id.*, *Ann. Mus. Genova* (2nd ser.), v, p. 420 (1883), and *P.Z.S.*, 1893, pp. 526, 527.

Mr. Boulenger has given an excellent description of this larva in the "Annali" of the Genoa Museum and has also published a key whereby those tadpoles of the genus *Rana* which are provided with large ventral suckers may be distinguished from one another. This key is in the *P.Z.S.* for 1893.

Mr. Kemp obtained two tadpoles that agree well with Mr. Boulenger's description in a small stream running into the Dihang R. near Yembung in January. Their hind limbs have not yet appeared. An adult *R. afghana* was taken at the same place in the same month.

D. Larva of *RHACOPHORUS MACULATUS* (Gray).

(Plate iv, figs. 4, 5.)

Rhacophorus leucomystax, Flower *P.Z.S.* 1896, p. 906, pl. xlv, fig. 2, and 1899, p. 898, pl. lix, figs. 3, 3a, Butler, *Journ. Bombay Nat. Hist. Soc.*, xv, p. 202.

Rh. maculatus, Ferguson, *ibid.*, p. 504, pl. B, fig. 4.

Captain Flower, in the first of the two papers cited after his name, describes and figures the tadpoles of the Malay race (*leucomystax*) of this species, his specimens being from Singapore. In his second paper he does the same for tadpoles from Bangkok, which seem to differ in more than one particular from those of the Malay race. I propose here to describe those of the Himalayan and the Peninsular races—the latter from specimens taken in September in Orissa, the former from tadpoles caught in the neighbourhood of Darjiling and identified after breeding out several individuals from the same lot in Calcutta. In both cases the tadpoles were taken in pools of rain-water.

1. Tadpoles of Himalayan race (*himalayensis*).

Head and body moderately flat above, ovoid, rounded in front, convex on ventral surface.

Mouth nearly terminal, comparatively small; lips relatively narrow, both directed forwards; upper lip smooth except at the corners, which bear numerous rounded papillae; lower lip with a fringe interrupted in the middle, and consisting of similar papillae about three deep; dental formula 1: 3+3 1+1: 2 or 1: 3+3 3; beak in two parts; the upper beak not hooked, the lower crescentic; both parts massive, both serrated.

Eye and nostril.—Eye lateral, directed outwards ; nostril nearer tip of snout than eye.

Glands.—There is a large gland in front of and slightly below each eye.

Spiracle sinistral, pointing backwards and a little upwards, flap-like, large.

Anus dextral.

Tail long and slender, about twice as long as head and body, sharply pointed ; its outline not strongly sinuous ; fin-membranes deep throughout its length.

Colouration.—Mottled with dark brown on dorsal surface and sides ; fin-membranes minutely spotted ; ventral surface white.

Dimensions.—The following are the measurements of a tadpole with well developed hind legs (A) and of one in which the hind legs are just about to appear (B) :—

	A.	B.
Length	62 mm.	33 mm.
Length of head and body	20 "	12 "
Length of tail	42 "	21 "
Maximum width of body	13 "	8 "
Maximum depth of body	10 "	7 "
Maximum depth of tail	14 "	8 "

2. Tadpole of Peninsular Indian race (*maculatus*).

The larvae of this race differ from those of the Himalayan race in the following points :—

- i. The head and body are flatter above.
- ii. The lips are even narrower.
- iii. The habit is more slender and the dimensions less.
- iv. The colour is very pale brown or green with scattered pigment-cells of a darker shade on the back and sides.

Both races apparently differ from the Malay one in having the outline of the tail less strongly sinuous and in colouration. In all essential structural characters, however, they are identical. Ferguson's Travancore tadpoles differed slightly in colouration from my Orissa ones. The breeding season in Travancore lasts, according to that author, from June to November ; both in the E. Himalayas and the plains of Lower Bengal, it is in progress in July, August and September. I have never seen the tadpoles of either race in running water. The eggs in their frothy covering are usually laid at the edge of small pools.

E. Larva of *PHRYNODERMA MOLOCH*, sp. nov.

(Plate iv, fig. 6.)

The tadpoles are very like those of *Ixalus horridus*, Blgr.,¹ but have a longer and more pointed tail, and when full-grown develop the

¹ Report on the Reptiles in Annandale and Robinson's *Fasciculi Malayenses*, i. p. 139, pl. vi, figs. 2a, 2b (1902)

characteristic dorsal rugosities of the adult before the fore-legs break through.

The head and body are flattened and broadly oval, being only a little longer than broad. The tail as a whole tapers gradually and is nearly twice as long as the head and body. The colour is an almost uniform black or dark grey, except that the margins of the lips are white and the lower surface slightly paler than the dorsal. At the time at which the hind limbs begin to sprout out the skin is still quite smooth, but as they develop ridge-like warts appear on the dorsal surface. There are numerous little sensory pits arranged in lines on the head and body.

Mouth subterminal, small ; its lips by no means strongly developed. The margin of the upper lip for the most part smooth but with a few conical tubercles where it joins the lower lip, the margin of which is completely fringed with a double row of similar tubercles. The dental formula is 1 : 3+3 3, the first interrupted row of teeth being much longer than the other two. The beak consists of an upper and a lower part each of which is undivided ; both are rather narrow ; the anterior is narrowly crescentic in form, while the posterior is deeply excavated in the middle ; neither has a concavity on the exposed surface and both are minutely serrated.

Nostril and eye small, dorsal, directed upwards, by no means prominent. The internasal space is about $\frac{2}{3}$ the interorbital and $\frac{1}{2}$ that between the nostril and the eye ; the nostril is equidistant from the tip of the snout and the eye.

Sensory pits.—On the dorsal surface of the head and body four longitudinal lines of minute white pits can be distinguished, two on either side of the middle line. Two on each side start close together on the snout and after diverging in a sinuous manner so as to include between them a nostril and an eye meet together behind the latter. On the ventral surface there are three backwardly concave transverse lines, one just behind the mouth and two, the corner of which is somewhat sinuous, posterior to it. There is also a lateral line of pits along the muscular portion of the tail and a row of larger, isolated ones along the upper fin-membrane.

Spiracle sinistral, large and patent, directed backwards and situated somewhat low down on the side.

Vent in the middle line.

Tail.—The muscular portion is slender and tapers to a fine point. In the middle part of the tail the membrane equals it in depth both above and below, but at the base the membrane is shallow on both sides.

Dimensions of an individual with well-developed hind limbs (A) and of one (B) in which they have just begun to sprout :—

	A.	B.
Total length	58 mm.	55 mm.
Length of head and body	20 „	17 „
Length of tail	38 „	38 „

	A.	B.
Maximum breadth of body	16 mm.	13 mm.
Maximum depth of body	.. 11 „	9 „
Maximum depth of tail	10 „	12 „

Habitat, etc.—Four tadpoles were taken by Capt. de Courcy near Upper Renging between the 5th and the 10th of January, 1912. They were in a small pool of rain-water inside a log of rotten wood and were accompanied by two adults, the types of the species. There can be no doubt as to their identity, for the largest tadpole has already begun to develop the characteristic features of the adult. It has the hind limbs well developed.

F. UNDETERMINED RANID LARVA.

(Plate iv, fig. 11.)

A number of large Ranid larvae were taken on the 9th and 16th of January in the Egar stream between Renging and Rotung by Mr. Kemp and Capt. de Courcy. Specimens of *Rhacophorus naso*, *Ixalus asper* and *I. tuberculatus* were taken with one of them, and also several *Megalophrys* tadpoles. Mr. Kemp suggests that they may be the larvae of one of the adults captured on the same occasion and they may well be that of a *Rhacophorus*. There is, however, no direct evidence as to their parentage and none of them have reached a stage in their metamorphosis that would render it possible to throw any definite light on the subject. It may be as well to describe them, however, as they present certain features of biological interest. There can be no doubt that they are specifically identical.

Head and body convex above, flattened on the ventral surface, ovoid, snout bluntly rounded, projecting.

Mouth ventral, very large with both lips highly developed and the lower lip directed backwards and of unusual size ; upper lip not fringed except at the corners ; lower lip extending backwards for some distance behind the last row of teeth and covered in this region with relatively large rounded tubercles, also completely fringed with smaller and more elongate tubercles ; dental formula $2+2 : 1 : 5+5$; $1+1 : 2$; the outermost row of teeth on each side of the upper lip curving upwards and inwards to the margin of the lip which it reaches near the middle without meeting its fellow of the opposite side ; beak stout, in two parts ; the upper part having a flattened, nearly triangular area in the middle of the free margin ; this area separated from the rest of the upper beak by a minutely serrated ridge which is in continuity with the edge of the beak on either side ; the lower beak broadly V-shaped, minutely serrated on the margin.

Nosil and eye by no means prominent, of moderate size ; eyes lateral but near the dorsal surface, directed outwards and upwards ; nostril nearer eye than tip of snout.

Glands, etc.—No very definite glands can be detected but there are numerous little pits arranged along the upper lip, in a line extending

from near the tip of the snout upwards outside each nostril to the eye, round the eye and in two patches behind it.

Spiracle sinistral, large and tubular, pointing directly backwards.

Tail rather more than twice the length of the head and body, stout, with both fin-membranes well developed throughout its length.

Dimensions of tadpoles without hind limbs :—

Total length	76 mm.
Length of head and body	25 „
Length of tail	51 „
Maximum width of body	16 „
Maximum depth of body	15 „
Maximum depth of tail	18 „

Colour an almost uniform dark brown ; fin-membranes and ventral surface a little paler.

There can be no doubt that the lips form in this species a powerful sucker, as they do even in such tadpoles as that of *Rana liebigii*. It is the largest Indian tadpole with which I am acquainted.

Fam. PELOBATIDAE.

G. H. Larvae of MEGALOPHRYS spp. (Plate iv, figs. 8, 9, 10.)

Weber, *Ann. Jard. Bot. Buitenzorg*, xv, suppl. ii, p. 5, 1898 ; Laidlaw, *P.Z.S.*, 1900, p. 889 ; Gadow, *Camb. Nat. Hist., Amphib. and Rept.*, p. 60, fig. 11 ; Boulenger, "Report on the Reptiles," *Fascic. Malay., Zool.*, i, p. 131 ; Annardale, *ibid.*, p. 275, fig. I ; van Kampen, *Natuurk. Tijd. v. Ned.-Ind.*, lxix (1), p. 27 ; Boulenger, *P.Z.S.*, 1908 (1), pp. 413, 426.

It has been pointed out by Mr. Boulenger and other authors that the larvae of the Oriental genus *Megalophrys* fall into two groups one of which resembles the larvae of European Pelobatidae and is in no way remarkable, while the other is distinguished from all other known tadpoles by the peculiar structure of the mouth. It is with the latter group that I am at present concerned.

The first representative of this group to be described was *M. montana* (Kuhl), a species not uncommon in hilly districts of Java and the Malay Peninsula but not known to occur in the Indian Empire. This larva has frequently been described and I need only refer to its colouration. The whole of the body and tail are dark brown with paler markings along the sides. At altitudes of from 2,000 to 5,000 ft. in the E. Himalayas a tadpole of precisely similar structure is abundant in small jungle streams, but it differs in colouration in two particulars, firstly in that the ventral surface is much paler than the sides and secondly that there are dark

instead of pale markings on the latter. These markings take the form of irregular spots and veinings. I have found every stage in the metamorphosis of this tadpole into *M. parva*, Boulenger.

A second tadpole of identical structure but slightly different colouration, larger size and more slender form also occurs in the E. Himalayas and was taken in considerable numbers in the Abor foot-hills. It differs from that of *M. parva* in lacking the dark markings on the sides or at any rate in having them much less conspicuous and consisting only of minute spots. I believe that this larva, which I have seen from the N. Shan States, is that of *M. major*, Boulenger, but have only circumstantial evidence in support of this belief.

A fourth *Megalophrys* tadpole was taken by Mr. Kemp in the Abor foot-hills. It differs from the one I have assigned provisionally to *M. major* in being of an almost uniform dark brown colour both above and below. There is no evidence that this is the tadpole of *M. kempii*, but it may be stated that a very similar larva was taken by Mr. F. H. Gravely on the Siamese frontier of Tenasserim near the Dawna Hills in company with a young frog that apparently represents *M. heteropus* (Boulenger), a species hitherto known from the hills of the Malay Peninsula.

It is a remarkable fact that whereas the tadpoles of *M. hasseltii* (Tshudi), a Malayan species, are of typical Pelobatid structure, those of other members of the genus from the same region are not only extremely different from that form but also extremely close to one another in structure. Were it not proved that the common Darjiling tadpole is the larva of *M. parva*, it would be difficult to believe that it was specifically distinct from that of *M. montana*, although there is no difficulty in distinguishing the adults. Seeing that these two larvae belong to different species, it is evident that the larva of other species must only be identified with great caution.

PART II.—BIOLOGICAL.

Of all the frogs and toads taken in the Abor foot-hills only two (*Rana limnocharis* and *Bufo himalayanus*) entirely lack adhesive digital disks, and we may take it that the majority of the other species are normally arboreal in habits. Some few of them, however, seem rather to be rupicolous and to live among stones at the edge of streams. It is probable that *Micrixalus borealis* belongs to the latter category, but *Ixalus tuberculatus*, although individuals were actually found at the edge of a stream, is known also to frequent tree-trunks. *Phrynoderma moloch* apparently affects dead tree-trunks and lays its eggs in or near small masses of water that accumulate in them. In the dense jungles of the Malayo-Himalayan tract most of the *Batrachia* are as a rule arboreal and Mr. Kemp's collection in this respect is no exception. It may be noted, further, that the great majority of his specimens are protectively coloured and resemble either the lichen-clad bark of jungle trees or the green leaves of their foliage.

A very interesting set of biological phenomena is illustrated by the tadpoles of the Abor foot-hills, which agree precisely in this respect with those of the Darjiling district and of hilly districts in the Malay Peninsula. In the great majority of cases their structure is adapted for life in rapid-running mountain streams subject to sudden floods, and they are provided with special apparatus with which to contend against the dangers incidental to such a life. In other words, they possess special structural facilities either for clinging to fixed objects or else for floating away lightly on the surface of the flood.

There has been considerable dispute as to the function of the lozenge-shaped structure surrounding the mouth of the tadpole of *Megalophrys montana* and similar species. Dr. Gadow suggests that the peculiar teeth on the inner surface may be used for scraping the leaves of water-plants and Dr. van Kampen has observed the tadpoles rasping algae from the sides of a glass aquarium with them ; but observations made in the Malay Peninsula, Burma and the Himalayas confirm me in the



FIG. 1.—Tadpoles of *Megalophrys montana*.

opinion, which was originally advanced by Prof. Max Weber, that the chief function of the whole structure is to act as a float whereby the tadpole can be carried along on the surface, and also convince me that this function is correlated with the fact that floods are one of the chief dangers which tadpoles living in mountain streamlets have to fear. I reproduce above a photograph of two living tadpoles of *M. montana* taken some years ago in the Malay Peninsula. One of these tadpoles is lying at the bottom of the basin of water in which they were photographed and has the float folded, the other has it expanded on the surface. The water was too shallow for this tadpole's tail to hang vertically downwards as it usually does. Dr. Gadow's figure in the *Cambridge Natural History* is therefore more correct in this respect.

Although to act as a float is the main function of the oral apparatus of these *Megalophrys* tadpoles, and although the teeth on this apparatus are used for scraping off algae from stones (and, I think from an examination of the contents of their stomachs, also fungi from rotting wood), the apparatus has at least two other uses. In the first place it facilitates breathing when the tadpole is lying among vegetable débris in the corner of little pools, and in the second it assists it to crawl over wet rocks and thus, perhaps, to reach a fresh pool if the one in which it is confined dries up during a period of drought. The first of these secondary func-

tions only comes into play when the funnel is folded. As is shown in my figure in *Fasciculi Malayenses*, the mouth is well protected by the manner in which the lower part of the apparatus folds upwards over the upper part, but the lateral parts do not completely close together. Each margin bends inwards towards the other in such a way that on each side a narrow tube is formed down which water free from all but very minute fragments of débris is drawn to the mouth by the respiratory movements.

I have often observed the larvae of *M. parva* adhering to rocks at the edge of small streams in the E. Himalayas and even making their way up the faces of rocks in this position until at least the anterior part of the body was out of water. Adhesion was brought about very largely by the application of the outer surface of the lower part of the apparatus, which only bears teeth on the inner surface, to the face of the rock ; but the ventral surface of the body was applied in the same manner, while the movements of the powerful tail thrust the animal upwards. A small fish (*Nemachilus rupicola*) common in the same little streams actually, as I have myself observed, makes its way over rocks from pool to pool in a very similar manner, adhering partly by means of the smooth skin of its belly and partly by means of its lips.

Other tadpoles which frequent the streams of the Himalayas, instead of allowing themselves to be borne away lightly on the surface of the flood, adhere to rocks at the bottom or sides of the stream and have been found firmly fixed even in the immediate vicinity of waterfalls. The majority of the species both of the Darjiling district and of the Abor foot-hills adhere mainly by means of their lips, the inner, tooth-bearing surface of which is applied to the rocks. It is for this reason that in these species the position of the mouth is much more nearly horizontal than it is in many other tadpoles and the lips themselves are more or less enlarged. In such forms as the larva of *Rana liebigii* the lips, although they are distinctly broader than in the typical *Rana* tadpole and are directed forwards and backwards respectively instead of both forwards, are not monstrously developed ; but in other forms, such as the large unidentified Ranid larva described above, they attain enormous dimensions and form together a powerful sucker possibly rendered necessary by the large size and heavy build of their possessor, for even the lips of the tadpole of *R. liebigii* enable it to cling tightly.

In some other species a sucker quite separate from the lips and not homologous with the sucker that many young *Batrachia* larvae possess, is found on the ventral surface, doubtless for the same purpose. *Rana afghana* and several allied Himalayo-Malayan frogs have this sucker in their larval stage.

It is noteworthy that the means whereby the fishes of the streams of the Himalayas adhere to rocks are analogous to those adopted by the tadpoles which live in the same environment. *Nemachilus rupicola*, as has already been stated, clings to rocks by means of the smooth skin of its belly and of its enlarged lips, just as the tadpole of *Megalophrys parva* does ; several other mountain Cyprinidae have their mouths

modified in a manner that would suggest their being used in the same way as the lips of *Rana liebigii*, while *Discognathus lamta* certainly adheres to the bottom in rapid water very largely by means of the lips, which recall those of the unidentified Ranid tadpole of the Abor foot-hills. The Silurid genera *Pseudecheneis* and *Glyptosternum*, on the other hand, cling by means of a separate abdominal sucker as the tadpole of *R. afghana* does.

There can be no doubt that these are instances of convergence, and there is some evidence that even in the case of the tadpoles of mountain streams, the same method of adhering to fixed bodies in rapid-running water has been acquired independently in some instances by different species. The lips of the tadpole of *Bufo penangensis*,¹ for example, appear to resemble closely those of the unidentified Abor larva, which must certainly be assigned to the Ranidae, and enlarged lips like those of *Rana liebigii* are found in different species the adults of which are by no means closely allied. It can hardly be that the adaptation of such tadpoles, striking as it is, has been brought about by genetic relationship between the different species that possess it. The adults of most of the Abor frogs and toads have developed adhesive disks on their fingers without being in all cases closely related to one another, and we must suppose that the development of special abdominal suckers or of greatly enlarged lips is a similar phenomenon due, directly or indirectly, to environment rather than to the possession of a common ancestry by animals which have undergone parallel evolution in one particular.

PART III.—GEOGRAPHICAL.

GEOGRAPHICAL LIST OF THE SPECIES IN THE COLLECTION.

NAME.	SIKHM.	ASSAM.	BURMA.	REMARKS.
RANA ..	x	x	x	Cosmopolitan.
1. <i>R. cyanophlyctis</i> ..	x	x	x	Throughout Oriental Region.
2. <i>R. liebigii</i> ..	x	x	x	..
3. <i>R. tigrina</i> ..	x	x	x	Almost all over Oriental Region.
4. <i>R. limnocharis</i> ..	x	x	x	Even more widely distributed.
5. <i>R. alticola</i>	x	x	Also in N. W. Himalayas, Bengal and Orissa.

¹Flower, *P.Z.S.*, 1899, p. 909.

NAME.	SIKKIM.	ASSAM.	BURMA.	REMARKS.
6. <i>R. granulosa</i>	x	x	Also in Yunnan; in Assam only N. of Brahmaputra.
7. <i>R. afghana</i> ..	x	x	x	Not in Afganistan.
8. <i>R. gerbillus</i> *
MICRIXALUS	x	Hitherto regarded as peculiar to S. India and Ceylon.
9. <i>M. borealis</i> *
RHACOPHORUS ..	x	x	x	Oriental Region, China, Japan, Madagascar.
10. <i>Rh. maximus</i> ..	x	x
11. <i>Rh. bimaculatus</i>	x	..	Only known hitherto from the Khasi Hills.
12. <i>Rh. naso</i> *
13. <i>Rh. microdiscus</i> *
14. <i>Rh. maculatus himalayensis.</i>	x	x	x	Also in Yunnan.
15. <i>Rh. tuberculatus</i>	x	..	Not known S. of Brahmaputra.
IXALUS ..	x	x	x	S. India, Ceylon, E. Himalayas, Assam, Burma, Indo-China, Malaysia.
16. <i>I. asper</i>	x	Also in Malay Peninsula and islands.
17. <i>I. annandalei</i> ..	x	x	..	Not known S. of Brahmaputra.
18. <i>I. argus</i> *
19. <i>I. tuberculatus</i>	x	Described from the Chinese frontier of Burma.
CHIRIXALUS	x	Hitherto only known from Burma.
20. <i>Ch. doriae</i>	x	..
PHRYNODERMA	x	x	Hitherto only known from Burma.
21. <i>Ph. moloch</i> *

NAME.	SIKHEM.	ASSAM.	BURMA.	REMARKS.
Bufo ..	x	x	x	Almost cosmopolitan; not in Australia or Madagascar.
22. <i>B. melanostictus</i> ..	x	x	x	Throughout Oriental Region.
23. <i>B. himalayanus</i> ..	x	E. Himalayas only.
MEGALOPHRYS ..	x	x	x	Damper parts of Oriental Region; not in S. India or Ceylon.
24. <i>M. major</i> ..	x	x
25. <i>M. kempfi</i> *

So far as the genera mentioned in the foregoing list are concerned, its most striking features are the number of species of *Ixalus* included and the fact that the allied Burmese genera *Chirixalus* and *Phrynoderma* are also represented. Until recently the genus *Ixalus*, which abounds in Burmese and Malay forms, was not known from any part of the Himalayas or Assam, and even now only one species has been proved to exist in the Himalayas, west of Bhutan, namely *Ixalus annandalei*, Boulenger.¹ It is probable that another species (*I. asper*) also occurs in the Nepal foot-hills, but, even so, the genus is characteristically non-Himalayan.² *Chirixalus* and *Phrynoderma* have hitherto been monotypic genera, as the former still remains, and have only been recorded from the Karin Hills.

The occurrence of the S. Indian genus *Micrixalus* in the Abor foot-hills is exactly parallel to that of the lacertilian genus *Salea*, a species of which has recently been described from the adjacent Dafia country.

We may analyse the list further as regards species in the following manner, dividing them into seven geographical groups:—

- | | |
|---|-------|
| A. Species apparently endemic in the foot-hills E. of Bhutan | 7=28% |
| B. Forms found in the Himalayas, Assam and Burma | 5=20% |
| C. Species of very wide distribution | 4=16% |
| D. Species known from Sikkim and from Assam S. of the Brahmaputra | 2= 8% |
| E. Species hitherto only known from Assam | 2= 8% |
| F. Species hitherto only known from Burma or Burma and Malaysia | 3=12% |
| G. Species only known from the Himalayas | 2= 8% |

¹ J.A.S.B., 1906 (2), p. 385.

² Cf. Blanford, *Phil. Trans. Roy. Soc.*, cxciv (B), p. 421 (1901).

Stated in this manner the figures show at a glance how very distinct the Batrachian fauna of the N. E. corner of Assam and the Abor foot-hills is from that both of the Himalayan foot-hills W. of Bhutan and from that of the district S. of the Brahmaputra. They also show, however, that there is considerably greater affinity in the latter direction than in the former, and it is by no means improbable that further research among the Batrachia of southern Assam will increase rather than diminish the resemblance. It is, moreover, noteworthy that one of the two purely Himalayan forms (*Bufo himalayanus*) represented in Mr. Kemp's collection of frogs and toads is of doubtful validity as a species and that specimens from the Abor hills do not agree in every respect with those from Darjiling.

Several of the species whose names occupy a place in Mr. Kemp's list were not taken actually in the foot-hills, but in the plains at their base. These species are the following :—*Ixalus assamensis* from group G ; *Rana alticola* and *R. granulosa* from group B, and *Rana cyanophlyctis*, *R. tigrina* and *Bufo melanostictus* from group C. If we eliminate these names, we get the following numbers and percentages in the different groups :—

A. Species apparently endemic in the Himalayan foot-hills E. of Bhutan	7=37%
B. Species from the Himalayas, Assam and Burma .	3=16%
C. Species of very wide distribution	1=5.25%
D. Species from Sikhim and from Assam S. of the Brahmaputra	2=10.5%
E. Species hitherto known only from Assam	2=10.5%
F. Species hitherto known only from Burma or from Burma and Malaysia	3=16%
G. Species known only from the Himalayas	.. 1=5.25%

These percentages, which are calculated roughly, show that the Batrachian fauna of the Abor foot-hills, in so far as it is illustrated by Mr. Kemp's collection, includes about 37% of apparently endemic species, about 16% of species that also occur both in Sikhim on the one hand and in Burma and Assam on the other and also of species hitherto known from Burma but not from Assam, about 10½% of species not known from Burma but common to the E. Himalayas and Assam, and the same percentage of species only known hitherto from Assam and about 5¼% both of widely distributed species and of exclusively Himalayan forms. A true jungle fauna, if the jungle be of the damp equatorial kind, rarely includes many species of Batrachia of a very wide distribution, and it is no exception to this rule to find that such forms are scarce in the Abor foot-hills ; the large number of endemic species is a correlated fact. It is well known to be the case that the reptiles and Batrachia of the E. Himalayas, Assam and Burma have a strong affinity, and it seems probable that a rich Malayan element has made its way northwards and westwards into the damp evergreen jungles of these countries, gradually becoming more and more attenuated as the climate grows drier and less equable towards the west.

All the evidence at present available, therefore, supports the view that the fauna of the extreme eastern part of the Himalayan foot-hills is not, at any rate so far as the frogs and toads are concerned, Himalayan in the sense in which the term has hitherto been understood, but allied rather to the fauna of Assam south of the Brahmaputra or even to that of Burma. In other words, Blanford's "Eastern Himalayan Tract" does not extend, so far as the Batrachia are concerned, nearly so far to the east as he believed, while his "Assam Tract" extends northwards to include the foot-hills north of the Brahmaputra as well as its upper valley and the mountains lying south of it. It is probable also that no very clear line of division exists between his "Assam Tract" and his "Upper Burma Tract."

In conclusion I must thank Mr. Kemp for the very careful way in which his collection was preserved and labelled, and also those who helped him, especially Capt. the Hon. M. de Courcy, for the interesting specimens they contributed.


EXPLANATION OF PLATE II.

ABOR BATRACHIA.


- | | |
|--|----------------------------------|
| FIG. 1.— <i>Rana gerbillus</i> ,
sp. nov. | 1a. Profile of head (From type). |
| „ 2.— <i>Micrixalus borealis</i> ,
sp. nov. | 2a. „ „ (From co-type). |
| „ 3.— <i>Rhacophorus naso</i> ,
sp. nov. | 3a. „ „ (From type). |
| „ 4.— <i>Rhacophorus microdiscus</i> ,
sp. nov. | 4a. „ „ (From type). |
| „ 5.— <i>Rhacophorus tuberculatus</i> ,
Anderson. | 5a. „ „ |




1



4.



3.



5




2.




EXPLANATION OF PLATE III.

ABOR BATRACHIA.

- | | |
|--|---------------------------|
| FIG. 1.— <i>Ixalus tuberculatus</i> ,
Anderson. | 1a. Profile of head. |
| „ 2.— <i>Ixalus annandalei</i> ,
Boulgr. | 2a. „ „ (× 2). |
| „ 3.— <i>Ixalus argus</i> , sp. nov. | 3a. „ „ (From type). |
| „ 4.— <i>Pharynoderma moloch</i> ,
sp. nov. | 4a. „ „ (From co-type). |
| „ 5.— <i>Megalophrys kempii</i> ,
sp. nov. | 5a. „ „ (From type, × 2). |



3.



1



4.



2.



5




EXPLANATION OF PLATE IV

HIMALAYAN TADPOLES.

- FIG. 1.—Tadpole of *Rana alticola*, Boulgr. 1a. Head of younger tadpole from above ($\times 2$).
- „ 2.—Tadpole of *Rana liebighii*, Günth. 2a. Mouth of same ($\times 2$).
- „ 3.—Tadpole of *Rana afghana* (Günth.) ($\times 1\frac{1}{2}$). 3a. Mouth and ventral sucker of same ($\times 2$).
- „ 4.—Tadpole of *Rhacophorus maculatus* (Gray), (typical form).
- „ 5.—Tadpole of *Rhacophorus maculatus himalayensis*, subsp. nov.
- „ 6.—Tadpole of *Phrynoderma moloch*, sp. nov. 6a. Head of larger tadpole from above.
- „ 7.—Tadpole of *Bufo himalayanus*, Günth. ($\times 2$).
- „ 8.—Tadpole of *Megalophrys parva*, Boulgr. ($\times 1\frac{1}{2}$). 8a. Head of same from above ($\times 2$). 8b. Ventral surface of same ($\times 2$) 8c. Mouth of same from in front ($\times 4$).
- „ 9.—Tadpole of *Megalophrys* (?) *major*, Boulgr. ($\times 1\frac{1}{2}$).
- „ 10.—Tadpole of *Megalophrys*, sp. indet. from Abor country ($\times 1\frac{1}{2}$).
- „ 11.—Tadpole of undetermined Ranid from Abor country. 11a. Mouth of same ($\times 2$).



11a x 2



11



2a x 2



1.



1a x 2



2.



3 x 1/2




3a x 2



4.



10



5




9.



6



8 x 1/2



8c x 4.



6a.



8a x 2.



8b x 2.



7 x 2.

A. C. Chowdhary, del.

HIMALAYAN TADPOLES.

II. REPTILIA.

(Plate v.)

By N. ANNANDALE, D.Sc., F.A.S.B., Superintendent,
Indian Museum.

Although the reptiles brought back by Mr. Kemp are perhaps less interesting than the Batrachia, this is due solely to the fact that the reptiles of the eastern parts of the Indian Empire have been far more carefully studied than the toads and frogs. The collection consists of 83 specimens of snakes, representing 26 species (of which 3 have not previously been described); 40 specimens of lizards, representing 16 species (of which 1 is new), and a single tortoise; that is to say, 124 specimens in all, representing 43 species.

PART I.—SYSTEMATIC.

LIST OF SPECIES COLLECTED OR OBSERVED.

- | | |
|---------------------------------------|---|
| EMYDOSAURIA. | |
| 1. <i>Gavialis gangeticus</i> . | 20. <i>Typhlops tephrosoma</i> . |
| | 21. <i>Typhlops diversiceps</i> , nov. |
| | 22. <i>Trachischium monticola</i> . |
| CHELONIA. | |
| 2. <i>Kachuga tectum</i> . | 23. <i>Blythia reticulata</i> . |
| | 24. <i>Aproaspidops antecursorum</i> ,
gen. nov., sp. nov. |
| LACERTILIA. | |
| 3. <i>Gymnodactylus khasiensis</i> . | 25. <i>Polydontophis collaris</i> . |
| 4. <i>Hemidactylus frenatus</i> . | 26. <i>Ablabes porphyraceus</i> . |
| 5. <i>Hemidactylus bowringii</i> . | 27. <i>Ablabes pavo</i> , nov. |
| 6. <i>Hemidactylus brookii</i> . | 28. <i>Ablabes frenatus</i> . |
| 7. <i>Hemidactylus platyurus</i> . | 29. <i>Simotes albocinctus</i> . |
| 8. <i>Draco maculatus</i> . | 30. <i>Oligodon erythrorachis</i> . |
| 9. <i>Ptyctolaemus gularis</i> . | 31. <i>Zamenis mucosus</i> . |
| 10. <i>Acanthosaura minor</i> . | 32. <i>Coluber taeniurus</i> . |
| 11. <i>Calotes versicolor</i> . | 33. <i>Dendrophis gorei</i> . |
| 12. <i>Calotes jerdoni</i> . | 34. <i>Pseudoxenodon macrops</i> . |
| 13. <i>Ophisaurus gracilis</i> . | 35. <i>Tropidonotus platyceps</i> . |
| 14. <i>Varanus bengalensis</i> . | 36. <i>Tropidonotus khasiensis</i> . |
| 15. <i>Tachydromus sexlineatus</i> . | 37. <i>Tropidonotus piscator</i> . |
| 16. <i>Mabuia macularia</i> . | 38. <i>Dipsadomorphus gokool</i> . |
| 17. <i>Lygosoma indicum</i> . | 39. <i>Psammodynastes pulverulentus</i> . |
| 18. <i>Lygosoma courcyanum</i> , nov. | 40. <i>Dryophis prasinus</i> . |
| | 41. <i>Callophis maccllellandii</i> . |
| OPHIDIA. | |
| 19. <i>Typhlops diardi</i> . | 42. <i>Amblycephalus monticola</i> . |
| | 43. <i>Trimeresurus monticola</i> . |
| | 44. <i>Trimeresurus gramineus</i> . |

EMYDOSAURIA.

1. *Gavialis gangeticus* (Gmel.).

Boulenger, *Fauna*, p. 3.

Although no specimens of the Gharial were obtained, Mr. Kemp tells me that it is not uncommon at Kobo together with the Gangetic Porpoise (*Platanista gangetica*). Apparently the shortnosed Crocodile (*Crocodilus palustris*) does not make its way so far up the Brahmaputra, although it occurs between Mangaldai in the Darrang district and Gauhati.

CHELONIA.

2. *Kachuga tectum* (Gray).

Boulenger, *Fauna*, p. 43 ; *Cat. Chelonia Brit. Mus.*, p. 58 ; Siebenrock, *Zool. Jahrb. Jena*, 1909, suppl. x, p. 454.

A shell and skull of the genus *Kachuga* were obtained from the Dihang R. below Pasighat and must be referred to this species. The shell, however, which measures 21.7 cm. in length, is narrower than is usually the case, and Mr. Kemp tells me that the soft parts were deep olive in life without reddish marks of any kind. Possibly the specimen represents a local race, but I have examined apparently typical individuals of *K. tectum* from N. E. Assam.

The limits of the range of *K. tectum* are very imperfectly known. It has been recorded from several localities in central and western India and certainly occurs in many different parts of the river-systems of the Indus, Ganges and Brahmaputra. All specimens, however, that I have examined from central India, although several were labelled *K. tectum*, actually represented *K. intermedia*, a very closely allied and somewhat variable form that is common all over the Mahanaddi river-system and also in the lower reaches of the Godavari. I have recently seen large numbers of this form from Cuttack and Sambalpur in Orissa and find that the proportions and outlines of the neural plates are so variable that little reliance can be placed on them in separating the "species" from *K. tectum*. The skulls of the two forms are identical. The only constant feature of difference is therefore colouration, and I am inclined to think that the late Dr. Blanford¹ was right in regarding *K. intermedia* as being only a "variety" (or, as I would prefer to call it, a subspecies) of *K. tectum*. The true *K. tectum* also occurs, according to Siebenrock, in Cochin China, *Pangshura cochinchinensis*, Tirant,² being synonymous.

Mr. Kemp tells me that he could hear of only one land-tortoise having been seen during the Expedition and that it was not secured. Terrestrial Chelonia hibernate in northern India, but no species has as yet been recorded from the Himalayas.

¹ *J.A.S.B.*, (2) xxxix (1870), p. 339, and xlvi (1879), p. 110.

² *Etudes Div. Miss. Pavie*, iii, p. 494 (1904).

LACERTILIA.

Fam. GECKONIDAE.

3. **Gymnodactylus khasiensis** (Jerdon).

Boulenger, *Fauna*, p. 68.

Originally described from the Khasi Hills, this lizard appears to have a fairly wide range in the mountains of Assam and northern Burma.

Two specimens were taken by Mr. Kemp at Kobo in December under the bark of a tree, together with a young *Ptyctolaemus gularis*. Others were obtained by the 32nd Sikh Pioneers at an altitude of about 2,000 feet at Upper Rotung.

My own *G. himalayicus*¹ is a very closely related species, differing in its slighter build, more obscure colouration and less compressed digits and also in having a well-defined triangular patch of enlarged scales just behind the praeanal pores. This last point, however, is not a very good character as in some specimens of *G. khasiensis*, of which I have examined a large series, there is a single enlarged scale, or even a pair of such scales, in the same position, although in others the scales are uniformly small.

4. **Hemidactylus frenatus**, D. & B.

A very common lizard at low altitudes in the E. Himalayas, Assam, Burma and the Malay Peninsula ; often found in houses.

A specimen was taken on a lamp-post in the streets of Dibrugarh in November.

5. **Hemidactylus bowringii** (Gray).

Not uncommon at low altitudes in the E. Himalayas and in Assam and Burma, this species is occasionally found in houses. It is, however, more often taken in the jungle.

There is a specimen in the collection from Sadiya.

6. **Hemidactylus brookii**, Gray.

Hemidactylus gleadowii, Boulenger, *Fauna*, p. 86, fig. 27.

Hemidactylus brookii, *id.*, *Ann. Mag. Nat. Hist.*, (7) i, p. 123 (1898), and Annandale, *Rec. Ind. Mus.*, vii, p. 45 (1912).

This species, which is one of the commonest house-lizards all over the plains of India (in northern Madras certainly the commonest), has a very wide distribution in the tropics but apparently avoids mountainous regions. The highest altitude from which I have seen a specimen is 4,500 ft., and this was on the isolated mountain Parésnath in Chota Nagpur, now in the new Province of Bihar and Orissa. Several specimens were obtained at Sadiya.

¹ *J.A.S.B.*, 1906, p. 287, and *Rec. Ind. Mus.*, I, p. 152, pl. vi, figs. 1, 1a, 1b, 1c, 1d (1907).

H. brookii is by no means always domestic in its habits, but individuals found living wild among rocks or in jungle are as a rule darker in colour and somewhat stouter in form than those which inhabit human dwellings. In either case the species is mainly nocturnal.

*H. subtriedroides*¹ from Upper Burma only differs from *H. brookii*, of which it should probably be regarded as a variety, in its rather stouter habit and much larger dorsal tubercles. The tubercles are extremely variable in size even within the strict limits of *H. brookii*, and less frequently vary also in number. There is in the Indian Museum a specimen from Bangalore on which there are only two dorso-lateral rows of tubercles on each side, the middle region of the back being perfectly smooth. This individual was taken by myself, together with a normal one, on the post of a railing.

7. *Hemidactylus platyurus* (Schneid.).

This gecko is common in the E. Himalayas, in which it is found as a rule at rather higher altitudes than the two preceding species. It is widely distributed in the Indo-Malayan and Malayan countries. So far as I am aware the western limit of its range is situated in Nepal, my own *H. nepalensis*² being undoubtedly synonymous. In the Darjiling district *H. platyurus* is often found in houses, but it is naturally an inhabitant of tree-trunks.

An unusually dark specimen was taken by Mr. Kemp at Pasighat under the bark of a tree.

Fam. AGAMIDAE.

8. *Draco maculatus* (Gray).

Boulenger, *Fauna*, p. 112.

This species is widely distributed in Assam, Burma, the Malay Peninsula and Yunnan, the western limit of its range being situated near the point in Assam north of the Brahmaputra at which that river bends southwards. There are three specimens in Mr. Kemp's collection :—

- | | | |
|-------|----------------------------|----------------------------|
| 16881 | Janakmukh (alt. 600 ft.) | Capt C. E. Edward-Collins. |
| 16882 | Yembung (alt. 1,100 ft.) : | |
| | “ found in a hut in camp.” | S. W. Kemp. |
| 16997 | Pasighat (alt. 500 ft.) | W. Cave-Brown. |

Mr. Kemp describes the colouration of No. 16882, an immature male, as follows :—“ Colour of back mottled warm brown, dark brown and black. Neck above with two elliptical black markings and a pair of black spots. ‘ Wings ’ above orange-brown. Belly dull greenish yellow ;

¹ Annandale, *Ann. Mag. Nat. Hist.*, (7) xv, p. 29 (1905), and *J.A.S.B.*, 1905, pl. ii, fig. 1.

² *Rec. Ind. Mus.*, i, p. 151, pl. vi, figs. 2, 2a, 2b, 2c (1907).

bright yellow beneath and on either side of pouch. 'Wings' beneath dull yellowish green, the orange colouring having a tendency to show through. Lateral neck-flaps orange beneath. Predominant headcolour very dark brown."

The only other species of *Draco* as yet found in Assam is *D. norvillii*, Alcock,¹ a single specimen of which, now in the Indian Museum, was taken at Dum Duma in N.-E. Assam some years ago. This species is closely related to *D. blanfordii*, Boulenger, from which it differs in having the tympanum completely covered with small scales. From *D. maculatus* it is easily distinguished by its longer snout and by the broad patches of enlarged scales scattered on each side of the back along the base of the alar membrane.

9. *Ptyctolaemus gularis*, Boulgr.

Boulenger, *Fauna*, p. 117 ; Annandale, *J.A.S.B.*, 1905, p. 85 ; Wall, *Journ. Bomb. Nat. Hist. Soc.*, xviii, p. 505.

This is a very rare lizard in collections. I have only seen three specimens hitherto, two of them from Assam N. of the Brahmaputra. Major F. Wall, however, states that the species is common at Shillong in the Khasi Hills (4,900 ft.) and our third specimen is from that locality. The western limit of its range is apparently the same as that of *Draco maculatus*. It is possible that both will be found in the Buxa duars ; we have both in the Indian Museum from Goalpara (Dhubri). *Ptyctolaemus* is certainly not indigenous in Calcutta.

Mr. Kemp found a very young specimen at Kobo in November ; although probably not long hatched (or born), it bore the characteristic grooves on the side of the neck. Adults were taken at Kobo by Capt. de Courcy and at Rotung by Mr. Kemp, who did not obtain the species at altitudes of over 1,300 ft.

10. *Acanthosaura minor* (Gray).

Boulenger, *Fauna*, p. 127.

This species is known from both Sikkim and the Khasi Hills. It is, however, rare in the Darjiling district. A small specimen was obtained at Upper Rotung (alt. ca. 2,000 ft.) in January.

11. *Calotes versicolor* (Daud.).

Boulenger, *Fauna*, p. 135, fig. 42 ; Annandale, *Rec. Ind. Mus.*, vii, p. 46.

The only specimen in the collection is a very young one obtained at Sadiya.

¹ *J.A.S.B.*, (2) lxiv, p. 14, pl. iii (1895).

12. **Calotes jerdonii**, Günth.

Boulenger, *Fauna*, p. 137.

Common in the Khasi Hills ; Col. Godwin-Austen obtained specimens in the Daffa Hills to the west of the Abor country, in which Mr. Kemp collected it at Komsing, Yembung and Balek. Both his specimens and Col. Godwin-Austen's are quite typical.

Fam. ANGUIDAE.

13. **Ophisaurus gracilis** (Gray).

Boulenger, *Fauna*, p. 159, fig. 47.

A common species in the E. Himalayas at altitudes of between 4,000 and 5,000 ft.; it also occurs in the Khasi Hills, in Upper Burma and Yunnan and probably in the hills of Pegu.

A number of individuals of different ages were taken in the neighbourhood of Upper Rotung and Upper Renging (2,000—2,150 ft.) by the 32nd Pioneers while road-making. The large specimens have the back of a bright brick-red with very conspicuous blue cross-bars ; on the tail the red fades to dull brown. The young are white with two blue-black stripes down each side and a somewhat indistinct and broken mid-dorsal stripe of the same colour ; the latter is crossed at intervals by transverse rows of very small black spots, and there are rather larger black spots scattered on the lips and snout.

Fam. VARANIDAE.

14. **Varanus bengalensis** (Daud.).

All over India and Ceylon except at high altitudes ; also in Upper Burma.

A skin was obtained at Kobo by Col. (now Brigadier-General) D.C.F. Macintyre and presented by him to the Indian Museum.

Fam. LACERTIDAE.

15. **Tachydromus sexlineatus**, Daud.

Boulenger, *Fauna*, p. 169 ; *Fasciculi Malayenses*, i, p. 158 ; Annandale, *J.A.S.B.*, 1905, p. 140.

This lizard is widely distributed in the damper parts of the Oriental Region. It is not found in Peninsular India.

A specimen was taken at Janakmukh (alt. 600 ft.) under the bark of a tree in December. It was probably hibernating, as the species is usually found among long grass.

Fam. SCINCIDAE.

16. *Mabuia macularia* (Blyth).

A very common skink all over the plains of India, Burma and Ceylon. Two specimens were taken at Sadiya in November, "under chips of wood."

17. *Lygosoma indicum* (Gray).

Lygosoma indicum and *L. zebratum*, Boulenger, *Fauna*, p. 195.

Lygosoma indicum, *id.*, *Ann. Mus. Genova* (2nd ser.), xiii, p. 319.

A very common species in the foot-hills of the E. Himalayas and also in hilly country in Assam and Burma.

Specimens were taken under stones in the neighbourhood of Rotung and Upper Rotung (1,300 to 2,000 ft.) in December, January and March.

18. *Lygosoma courcyanum*, sp. nov.

(Plate v, fig. 5.)

Subgenus *Hinulia*. Allied to *L. (Hinulia) cacharensis*¹ from Assam but much more slender and with shorter limbs.

Habit slender, lacertiform; the distance between the axilla and the groin nearly twice that between the tip of the snout and the fore limb. Tail nearly twice as long as head and body. Total length 100 mm.

Head small, narrow, triangular; snout pointed; loreal region vertical; lower eye-lid scaly; ear-opening subcircular, much smaller than eye, without lobules; nostril pierced in a single nasal. Rostral much broader than deep, forming an extensive suture with fronto-nasal; fronto-nasal undivided, much longer than praefrontals; no supranasals; praefrontals separate short; frontal shorter than its distance from tip of snout, a little shorter than the parietals; its greatest breadth to its length as 7 to 9; parietals not meeting behind interparietal, forming a lengthy suture; interparietal small; a single pair of enlarged nuchals; 4 large, subequal supraoculars; 7 upper labials, 4th, 5th and 6th under eye.

Scales of body smooth, imbricate, in 24 rows; the ventrals slightly larger than the dorsals; two large praeanales.

Limbs short but well-formed, separated by a considerable distance where adpressed. Toes not compressed, of moderate length; 11 smooth lamellae under 4th toe.

Colouration.—Dorsal surface olive-brown minutely speckled with black; tail rather darker than back; a narrow pale band extending on each side from above the eye to the base of the tail; a rather broader black band running immediately below it; sides yellowish speckled with black; lateral surface of tail suffused with slate grey; ventral surface yellowish, speckled with slate-grey on tail.

¹ Annandale, *J.A.S.B.*, 1905, p. 145.

Dimensions of type :—

Total length	.	.	100 mm.
Length of head and body	.	.	35 „
Length of head			8 „
Breadth of head	.	..	4 „
Fore limb			9 „
Hind limb			13 „

Type.—No. 16900 in the Indian Museum register of Reptiles and Batrachia.

Habitat.—Two specimens were taken by Capt. the Hon. M. de Courcy, one at Rotung (1,300 ft.), the other at Upper Rotung (ca. 2,000 ft.).

OPHIDIA.

Mr. Kemp asks me to state that a very large proportion of the snakes in his collection were captured by the officers and men of the 32nd Sikh Pioneers at the instance of Capt. the Hon. M. de Courcy. They were found while road-making, chiefly in December and January, and were probably hibernating at the time. Other snakes were presented by Capt. J. S. O'Neill, Capt. F. H. Stewart and Capt. R. S. Kennedy of the Indian Medical Service.

Fam. TYPHLOPIDAE.

19. *Typhlops diardi*, Schleg.

Boulenger, *Fauna*, p. 238, fig. 70.

A considerable number of specimens of this common Himalayo-Burmese species were taken at Kobo, Pasighat, Janakmukh and Balek, several of them having been found crawling about on the surface in camp after rain.

T. diardi occurs all over Assam and Burma and is also found in the Himalayas, Siam and other adjacent countries.

20. *Typhlops tephrosoma*, Wall.

Wall, *Journ. Bombay Nat. Hist. Soc.*, xviii, p. 314.

A small specimen taken by the 32nd Sikh Pioneers at Janakmukh agrees well with Major Wall's description of the type from the Khasi Hills.

21. *Typhlops diversiceps*, sp. nov.

(Plate v, fig. 1.)

This species belongs to the same group as *T. braminus* and *T. beddomii*, but the anterior nasal is not in contact with the praeocular and the posterior nasals do not meet behind the rostral.

Snout rounded, projecting. Nostril lateral, nasal completely divided; rostral barely reaching the level of the eyes, not half as wide as snout; upper head-scales about twice as large as body-scales, trans-

verse; anterior nasal widely separated from praeocular, in contact with first labial below; posterior nasal much larger, in contact with second labial, not meeting its fellow on the top of the head; eyes moderately distinct; praeocular larger than ocular, in contact with second and third labials; ocular in contact with third and fourth labials; 4 upper labials. Diameter of body 40 times in total length; tail longer than broad, ending in a minute spine; 18 scales round body.

Colour dark olive-brown, slightly paler on ventral surface. Dorsal surface of head chestnut, ventral surface pale yellow; the latter shade extending upwards on either side to the level of the eyes in the form of a narrow triangle, very clearly defined posteriorly on the throat.

Length 160 mm. : length of tail 3 mm.

Type.—No. 16864, Ind. Mus.

Locality.—Pasighat (500 ft.) 25-iii-12. (Capt. R. S. Kennedy, I.M.S.)

Fam. COLUBRIDAE.

Subfamily COLUBRINAE.

22. *Trachischium monticola* (Cantor)

Boulenger, *Fauna*, p. 286.

A common snake in the hills of Assam, less abundant in the E. Himalayas. Ten specimens were taken by the 32nd Sikh Pioneers while road-making in the neighbourhood of Upper Rotung (alt. ca. 2,000 ft.) in January. Capt. de Courcy took another in the Sirpo valley near Renging.

23. *Blythia reticulata* (Blyth).

Boulenger, *op. cit.*, p. 287, fig. 92.

Three specimens were taken at Upper Renging, at Upper Rotung and in the Sirpo valley near Renging. The species, which is the only one in the genus, is characteristically Assamese. Fresh adult specimens are almost black in colour with a beautiful deep-blue iridescence, the pale markings becoming inconspicuous with age.

Aprospidops, gen. nov.

The new genus is allied to *Trirhinopholis* and *Plagiopholis*, Boulenger, both of which are only known from Burma. It thus belongs to a peculiar little group of monotypic genera that includes *Blythia* and the two just mentioned and inhabits hilly country in Burma and Assam. *Aprospidops* can be recognized easily by the fact that there is an azygous shield between the rostral and the supranasals and also a small postnasal on each side. There is no praeocular and no loreal, unless the latter name should be applied to the small scale I have called the postnasal.

Maxillary moderate, with about 20 teeth, which decrease slightly in size from before backwards; mandibular teeth similar to maxillary. Head not distinct from neck; eye small, with round pupil; nostril pierced between two nasals, the posterior of which is followed by a small scale (postnasal) in contact with the supranasal, prae-frontal and first upper labial. Prae-frontal entering the eye and in contact with upper labials; no praecular; rostral separated from supranasals by a triangular azygous shield. Scales smooth, without apical pits, imbricate, in 12 straight rows; ventrals rounded. Tail short; subcaudals in two rows.

24. *Aproaspidops antecursorum*, sp. nov.

(Plate v, fig. 2.)

Snout moderate, rounded. Rostral much wider than deep, just visible from above, much deeper than the shield which separates it from the supranasals; the latter completely divided, about half as long as the prae-frontals, which are also completely divided; frontal about $1\frac{1}{3}$ times as long as broad, about as long as its distance from the snout, much shorter than the parietals. Nostril between two small, deeply concave scales, separated from the first labial; postnasal triangular, smaller than the two nasals of one side together. Prae-frontal in contact with second and third upper labials, the latter and the fourth entering the eye; no subocular; a single large postocular; supraocular much longer than deep; 6 upper and 6 lower labials; loreals 1+2. Two pairs of chin-shields, posterior pair very short, in contact with the fourth pair of labials; the first pair in contact with three pairs of labials, first pair of labials forming a long suture behind the mental. Ventrals 136; subcaudals 16; anal divided; tail ending in a sharp spine.

Colour dark olive, each body-scale with a slightly darker border; ventrals and subcaudals with pale borders; an incomplete white collar extending over the greater part of the neck on each side some distance behind the gape.

Length 162 mm. : length of tail 12 mm.

Type.—No. 16844, Ind. Mus.

Locality.—Janakmukh, 600 ft. : 13-xii-11.

The single small, perhaps immature, specimen was taken by the 32nd Sikh Pioneers while road-making. It closely resembles young *Blythia reticulata* in appearance but can easily be distinguished by its circular pupil and by the extra scales behind the rostral and the nasals.

25. *Polydontophis collaris* (Gray).

Boulenger, *op. cit.*, p. 302.

A common snake all over the Himalayas up to 10,000 feet, in Assam, Upper Burma, etc. Specimens were taken in the Sirpo valley near Renging and at Kobo by Capt. de Courcy. They represent the typical form.

26. **Ablabes porphyraceus** (Cantor).

Boulenger, *op. cit.*, p. 308.

A common Malayo-Himalayan snake probably not found at great altitudes. Specimens were taken at Balek and between Kalek and Misshing by Capt. Wilson and Mr. Kemp respectively.

27. **Ablabes pavo**, sp. nov.

(Plate v, fig. 3.)

A magnificent species easily recognized by the large black and yellow ocelli on its back, but also to be distinguished by numerous scale-characters, notably by the large number of ventrals.

Rostral much wider than deep, visible from above; supranasals distinct, about half as long as praefrontals, which are also distinct; frontal $1\frac{1}{2}$ times as long as broad, a little shorter than its distance from the tip of the snout, almost as long as the parietals; nasal completely divided, extending backwards as far as the suture between the second and third labial; no distinct loreal; a single praeocular; two post-oculars, only the upper one in contact with the parietal; 7 upper labials, the third and fourth entering the eye; temporals 2+2; the shields on the sides of the head minutely pitted; two pairs of chin-shields, the anterior pair in contact with three pairs of labials, the posterior pair in contact with only one pair. Scales in 19 rows. Ventrals 233; subcaudals 80; anal and subcaudals divided.

Colour.—Sides and back pale bluish grey, each scale bearing an irregular patch of peach colour; back ornamented with a row of large black longitudinally oval rings, each with a yellow centre and separated one from another by only a short interspace; on the tail the yellow centres of the rings break up into small spots and finally at the tip, disappear altogether; about 50 rings in all; sides with irregular zig-zag black, yellow-edged vertical bars; ventral surface yellowish with numerous black cross-bars which are usually interrupted in the middle line; head black with a broad yellow bar across the snout, a second across the vertex behind the eyes and a third across the nape, the two latter being Λ -shaped; lips, chin and throat yellow with large black spots.

Length 640 mm. : length of tail 108 mm.

Type.—No. 16797, Ind. Mus.

Locality.—Upper Rotung; taken by 32nd Sikh Pioneers while road-making, 13-xii-II.

28. **Ablabes frenatus** (Günth.).

Boulenger, *Fauna*, p. 306.

A characteristic Assamese species. Three specimens were taken at Upper Rotung (2,000 ft.) in January. Two were found while road-making, while one was sitting coiled up in the middle of a path.

29. *Simotes albocinctus* (Cantor).

Boulenger, *Fauna*, p. 312.

Not uncommon in the hills of Assam and Burma. Wall¹ has described, under the name *juglandifer*, a peculiar variety distinguished mainly by colouration but now regarded by him as a distinct species. It is from the E. Himalayas and Assam. Specimens of the typical form were taken by Capt. de Courcy at Kobo and in the Sirpo valley near Renging.

30. *Oligodon erythrorhachis*, Wall.

Wall, *Journ. Bombay Nat. Hist. Soc.*, xix, p. 923, pl.

Two specimens from Upper Rotung (alt. ca. 2,000 ft.) taken in December by the 32nd Sikh Pioneers must be referred to this species. Both, however, differ in colouration from the type. The smaller specimen measures 300 mm. in length and is of a brick-red colour with numerous white, black-edged cross-bars on the body and tail. There are faint traces of a dark mid-dorsal line and the head and ventral surface are marked as in Major Wall's figures. The second specimen is larger, measuring 510 mm., and differs from the smaller one in being of a deep crimson colour and having the cross-bars on the body and tail relatively broader and slate-grey instead of white.

31. *Zamenis mucosus* (Linn.).

Boulenger, *Fauna*, p. 324.

A small specimen of the Common Rat-Snake was taken at Janakmukh by Capt. O'Neill.

32. *Coluber taeniurus* (Cope).

Boulenger, *Fauna*, p. 333 ; *Fascic. Malay.*, i, p. 162.

Two specimens were taken in January at Upper Rotung by the 32nd Sikh Pioneers while cutting a road. The species has a somewhat curious distribution, ranging from Darjiling into not only south-western but also northern China. In the Malay Peninsula it is usually found in caves feeding on bats. Cavernicolous individuals are always very pale in colour, but it is by no means certain that this is not due to the direct effect of lack of light on the organism.

33. *Dendrophis gorei*, Wall.

Wall, *Journ. Bombay Nat. Hist. Soc.*, xix, p. 829, pl., fi. 1-3 (1910).

I doubt whether this is more than a local race of *D. pictus* peculiar to the north-east corner of Assam and the neighbouring foot-hills. There

¹ *Journ. Bombay Nat. Hist. Soc.*, xix, pp. 3, 8, and xx, p. 1162, fig.

are three specimens in the Abor collection which agree well with Major Wall's specimen from Dibrugarh now in the collection of the Indian Museum. They are from Kobo (400 ft.), from between Janakmukh and Balek and from the Siyom valley below Damda (ca. 1,400 ft.).

34. ***Pseudoxenodon macrops*** (Blyth).

Boulenger, *Fauna*, p. 340.

A specimen was taken near Sidi stream (alt. ca. 2,000 ft.) by the 32nd Sikh Pioneers. The species is very common in the Darjiling district from the base of the foot-hills up to 5,000 ft., occurring also in the hills of Assam and Burma but apparently in smaller numbers.

35. ***Tropidonotus platyceps***, Blyth.

Boulenger, *Fauna*, p. 344.

A very variable species common in the Himalayas up to 10,000 ft. ; also occurs in the Khasi Hills and the mountains of Burma. A specimen was taken at Upper Rotung.

36. ***Tropidonotus khasiensis***, Boulgr.

Boulenger, *Fauna*, p. 344, and *Ann. Mus. Genova* (2nd ser.), xiii, p. 322.

A scarce species hitherto only found in the Khasi and Karin Hills. One was taken at Rotung by Capt. de Courcy.

37. ***Tropidonotus piscator*** (Schneid.).

Boulenger, *Fauna*, p. 349.

Young specimens of this very common and widely distributed species were taken at Rotung and Upper Rotung (1,300 and ca. 2,000 ft.). It occurs in the W. Himalayas up to at least 4,500 ft.

Subfamily DIPSADOMORPHINAE.

38. ***Dipsadomorphus gokool*** (Gray).

Dipsas gokool, Boulenger, *Fauna*, p. 360.

Dipsadomorphus gokool, *id.*, *Cat. Snakes Brit. Mus.*, iii, p. 64 (1895).

A specimen was taken at Dibrugarh. It is not improbable that this snake is actually restricted to Assam, for Cantor's localities are notoriously inaccurate and the record of this species from Penang apparently rests on a specimen from his collection in the British Museum. Bengal and Assam were not clearly distinguished by many of the older naturalists who wrote on Indian reptiles.

39. **Psammodynastes pulverulentus** (Boie).

Boulenger, *Fauna*, p. 363, and *Cat. Snakes Brit. Mus.*, iii, p. 173.

Specimens, 8 in all, were taken at the following places during the Expedition:—Kobo, Balek, Rotung, the Sirpo valley near Renging and Renging. The species is widely distributed in the damper parts of the Oriental Region but does not occur in Peninsular India. Mr. Kemp's series exhibits a remarkable range of colour-variation, no two individuals being precisely alike in colouration.

40. **Dryophis prasinus**, Boie.

Boulenger, *Fauna*, p. 369.

A widely distributed species in the E. Himalayas, Assam, Burma, Indo-China and Malaysia. Specimens were taken at Janakmukh by Capt. O'Neill and Capt. de Courcy, at Rotung by Capt. F. H. Stewart and at Balek by Capt. Wilson. All belong to the typical leaf-green form.

Subfamily ELAPINÆ.

41. **Callophis maccllellandii** (Reinh.).

Boulenger, *Fauna*, p. 385, and *Cat. Snakes Brit. Mus.*, iii, p. 398.

A very common snake in the hills of Assam, occurring also in the E. Himalayas, Burma, S. China, etc. Two specimens of the typical form were taken at Upper Rotung (2,000 ft.) by the 32nd Sikh Pioneers.

Fam. AMBLYCEPHALIDÆ.

42. **Amblycephalus monticola** (Cantor).

Boulenger, *Fauna*, p. 415.

This snake, which occurs in the E. Himalayas, the hills of Assam and the Nicobar Is., is evidently very common in the Abor foot-hills. The 32nd Sikh Pioneers took 12 specimens of different sizes while road-making in the neighbourhood of Upper Rotung in January. They also caught specimens at Rotung and in the Sirpo valley.

Fam. VIPERIDÆ.

43. **Lachesis monticola** (Günth.).

Trimeresurus monticola, Boulenger, *Fauna*, p. 426.

Lachesis monticola, *id.*, *Cat. Snakes Brit. Mus.*, iii, p. 548.

Widely distributed in the E. Himalayas, the mountains of Assam, Burma and Yunnan and in hilly districts of the Malay Peninsula. Two specimens were obtained, one at Rotung, the other at Upper Rotung.

Mr. Kemp tells me that he heard it stated on several occasions that Russel's Viper (*Vipera russeli*) occurs in the Abor country but that these statements probably referred to *Trimeresurus monticola*.

44. *Lachesis gramineus* (Shaw).

Trimeresurus gramineus, Boulenger, *Fauna*, p. 429.

Lachesis gramineus, *id.*, *Cat. Snakes Brit. Mus.*, iii, p. 554.

A small specimen of the typical green form was taken at Kobo by Capt. Mitchell. The species is common in the hills of Assam, Burma, Malaysia, etc. and also occurs in the Himalayas.

PART II.—GEOGRAPHICAL.

The following species are only represented in Mr. Kemp's collection by specimens obtained at Dibrugarh or Sadiya or observed or caught in the Dihong R. :—*Gavialis gangeticus*, *Kachuga tectum*, *Hemidactylus frenatus*, *H. bowringii*, *H. brookii*, *Calotes versicolor*, *Mabuia macularia* and *Dipsadomorphus gokool*. These 8 species cannot, therefore, be regarded as having been proved to be inhabitants of the Abor foot-hills. The Indian Museum is, however, fortunate in possessing the collection of reptiles made in the Daffa foot-hills, which are situated a short distance to the west of the Abor country, by Col. Godwin-Austen many years ago, and it will add interest to geographical speculations about the fauna of the latter country if we combine the list of Mr. Kemp's collection with one of that obtained by Col. Godwin-Austen.

GEOGRAPHICAL LIST OF THE REPTILES KNOWN FROM THE HIMALAYAN FOOT-HILLS E. OF BHUTAN.

Name.	Assam.	Sikhim.	Burma.	REMARKS.
<i>Nicoria tricarinata</i> ..	x	Taken by Col. Godwin-Austen in the Daffa Hills; also known from Assam north of the Brahmaputra and from Chota Nagpur in Peninsular India.
<i>Gymnodactylus khasiensis</i>	x	..	x	Khasi Hills and Upper Burma.
<i>Hemidactylus frenatus</i> ..	x	x	x	Common in the Malay Peninsula; taken in the Daffa Hills; very widely distributed.

Name.	Assam.	Sikhim.	Burma.	REMARKS.
<i>Hemidactylus platyurus</i> ..	x	x	x	The Himalayas from the Nepal Valley eastwards ; also Ceylon, Malayasia, etc.
<i>Draco maculatus</i> ..	x	..	x	Malaya Peninsula, W. China, etc.
<i>Ptyctolaemus gularis</i> ..	x	Assam N. of Brahmaputra and Khasi Hills.
<i>Acanthosaura minor</i> ..	x	x	..	E. Himalayas and Khasi Hills.
<i>Calotes jerdonii</i> ..	x	Khasi Hills; both Dafa and Abor Hills.
<i>Japalura andersoniana</i> * ¹	Only known from the Dafa Hills.
<i>Salea austeniana</i> * ²	Ditto.
<i>Ophisaurus gracilis</i> ..	x	x	x	Perhaps only in Upper Burma.
<i>Varanus bengalensis</i> ..	x	..	x	I cannot find any record of the occurrence of this common Indian species in Sikhim, but it probably does occur there.
<i>Tachydromus sexlineatus</i>	x	x	x	I have not seen this species in the Himalayas, but it is said to occur in Sikhim: it is also found in S. China, the Malay Peninsula, etc.
<i>Lygosoma indicum</i> ..	x	x	x	A common species in Burma, not known from the Malay Peninsula.
<i>Lygosoma courcyanum</i> ,* sp. nov.	Only known from the Abor Hills.
<i>Lygosoma albopunctatum</i>	x	..	x	Also from Peninsular India, Malay Peninsula, etc.
<i>Typhlops braminus</i> ..	x	x	x	Very widely distributed; known from the Dafa Hills.

¹ Annandale, *J.A.S.B.*, 1905, p. 85.² *Id.*, *Rec. Ind. Mus.*, II, p. 37 (1908).

Name.	Assam.	Sikhim.	Burma.	REMARKS.
<i>Typhlops tephrosoma</i> ..	X	Only known from the Khasi Hills.
<i>Typhlops diardi</i> ..	X	X	X	Also in Indo-China, Siam, etc.
<i>Typhlops diversiceps</i> ,* sp. nov.	Only known from the Abor Hills.
<i>Trachischium monticola</i> ..	X	X	..	Common in the E. Himalayas and the hills of Assam.
<i>Aproaspidops antecursum</i> ,* gen. nov.	Genus only known from the Abor Hills.
<i>Blythia reticulata</i> ..	X	Common in the hills of Assam.
<i>Polydontophis collaris</i> ..	X	X	X	Also in W. Himalayas and S. W. China.
<i>Ablabes porphyraceus</i> ..	X	X	X	Also in Yunnan, Malay Peninsula, Sumatra, etc.
<i>Ablabes pavo</i> , sp. nov.	Only known from the Abor Hills.
<i>Ablabes frenatus</i> ..	X	Hills of Assam.
<i>Simotes albocinctus</i> ..	X	X	X	Common in Assam.
<i>Oligodon erythrorhachis</i> ..	X	Hitherto only known from hills S. of Brahmaputra.
<i>Zamenis mucosus</i> ..	X	X	X	A widely distributed Oriental species.
<i>Coluber taeniurus</i> ..	X	X	X	From the E. Himalayas to Yunnan; the Malay Peninsula; also Manchuria and Indo-China.
<i>Dendrophis gorei</i> ..	X	Only known from the N. E. corner of Assam.
<i>Pseudorenodon macrops</i> ..	X	X	X	Hill species but not found at great altitudes.
<i>Tropidonotus platyceps</i> ..	X	X	X	Found up to 10,000 ft.
<i>Tropidonotus khasiensis</i> ..	X	..	X	Found in Karim and Khasi Hills.
<i>Tropidonotus piscator</i> ..	X	X	X	Widely distributed in Oriental Region.

Name.	Assam.	Sikhim.	Burma.	REMARKS.
<i>Psammodynastes pulverulentus.</i>	x	x	x	Also in Indo-China, Siam, the Malay Peninsula and Archipelago, etc.
<i>Dryophis prasinus</i> ..	x	x	x	Ditto.
<i>Callophis macclellandii</i> ..	x	x	x	Also in Nepal, S. China, etc.
<i>Amblycephalus monticola</i>	x	x	..	Also in Nicobars.
<i>Trimeresurus monticola</i> ..	x	x	x	Also in Yunnan, Malay Peninsula, etc.
<i>Trimeresurus gramineus</i>	x	x	x	Also in Siam, S. China, Indo-China, the Malay Peninsula and Archipelago.

The only tortoise in this list (*Nicoria*¹ *tricarinata*) is one of a small group of more or less terrestrial Chelonia that occur both in Chota Nagpur in Peninsular India and in the northern part of Assam but apparently not in any intermediate locality. It is, however, evident that we know as yet very little about the distribution of the Indian land-tortoises.

The genera of lizards and snakes that are known to occur in the foot-hills N. of the Brahmaputra and E. of Bhutan are for the most part widely distributed. Only one appears to be endemic, viz., the new genus *Aproaspidops*, which is allied to the Burmese genera *Trirhinopholis*² and *Plagiopholis*³. The genera *Blythia* and *Ptyctolaemus* are characteristically Assamese, while *Pseudozenodon*, *Japalura* and *Draco* are restricted to the damper parts of the Oriental Region, the last-named being a characteristically Malaysian genus also found in S. India, while the two first are essentially continental in distribution. The genus *Salea* only occurs in the foot-hills E. of Bhutan, in Lower Burma and in S. India, being thus almost analogous in its range to the Ranid genus *Micrixalus*.

If we separate out the 42 species in the list into geographical groups as was done in the case of the Batrachia (p. 34 *antea*) we find that the following percentages can be calculated :—

I. Species of wide distribution in the Himalayas, Assam, Burma and Indo-China or Malaysia	16 = 38%
II. Species of very wide general distribution	6 = 14%

¹ Stejneger (*Proc. Biol. Soc. Wash.*, xv, p. 238 (1902), and Siebenrock (*Zool. Jahrb. Jena*, 1909, suppl. x, p. 494) point out that by the strict letter of the law of priority the name of this genus should be *Geoemyda*.

² Boulenger, *Cat. Snakes Brit. Mus.*, i, p. 419.

³ *Id.*, *ibid.*, p. 301.

III. Apparently endemic species	6=14%
IV. Species only known hitherto from Assam	7=16.5%
V. Assamo-Burmese species	3= 7%
VI. Himalayo-Assamese species	3= 7%
VII. Species only known from Assam and Peninsular India	1=2.5%

There are no exclusively Himalayan species in the list and none that have hitherto been known only from Burma. If we compare this analysis with that of the Batrachian fauna of the Abor Hills published on p. 35 of this volume, the chief apparent difference is that the endemic forms appear to be fewer and the representatives of what I have called elsewhere the Malayo-Himalayan fauna much more numerous. This may be due in part to the fact that the lizards and snakes of Assam are much better known than the frogs and toads, and in part to the wider distribution of species in the former groups. In the main the results are strictly comparable in what may be regarded as their essential feature, *viz.*, in illustrating the non-Himalayan nature of the Abor fauna. Unfortunately we know almost nothing of the reptiles and Batrachia of Bhutan, but the little that we do know would suggest that the eastern boundary of the true Himalayan fauna is formed by the R. Tista, which flows down south through the Himalayas to the west of Bhutan. This river, at any rate in its present course, is apparently a much more ancient one than the existing Brahmaputra.

The reptiles of the extreme east of the Himalayas, although they have strong Assamese affinities, are by no means identical with those of the Khasi Hills. It is particularly noteworthy that the species of *Japalura* which occur in the Daffa Hills is not nearly so closely related to the common *J. variegata*¹ of Sikkim as that species is to *J. planidorsata* of the Khasi Hills, and none of the six apparently endemic species have, so far as we are aware, close allies in the other mountains of Assam.

We may say therefore that the reptiles of the Abor foot-hills agree with the Batrachia in differing considerably from those of the foot-hills immediately to the west of Bhutan and in including a well-marked endemic element, but that they appear to be more closely connected with the fauna characteristic of the damp jungles of the E. Himalayas, Assam, Burma, Indo-China and the Malay Peninsula. It is to this fauna that it is convenient to apply the term "Malayo-Himalayan." Probably the comparative dryness of the forests on the foot-hills west of Nepal has prevented many damp-loving animals of Malayan origin from penetrating further afield in a westerly or north-westerly direction, while a smaller contingent has been stayed by the course of the R. Tista.

¹ I was wrong in stating that this species occurs in Assam (J.A.S.B., 1905, p. 92), having been misled by badly preserved specimens of *J. planidorsata*.

APPENDIX.

DESCRIPTIONS OF THREE NEW INDIAN LIZARDS.

As a matter of convenience I take this opportunity to describe three lizards, one of which is of particular interest in connection with the Abor fauna in that it comes from the Bengal frontier of Bhutan. The other two are from the Bombay Presidency and Sylhet respectively.

Hemidactylus platyceps, sp. nov.

Habit slender ; depressed ; size small.

Head narrowly ovoid, strongly depressed and very shallow ; snout bluntly pointed, about as long as distance between eye and ear ; eye small ; ear-opening minute, longitudinally oval. Rostral much more than twice as broad as deep, feebly cleft above ; nostril between rostral, first labial and three small scales ; 9 upper, 7 lower labials ; snout covered with strongly keeled granules, rest of head with smaller and slightly irregular convex granules ; two pairs of chin-shields, followed on each side by several irregular scales ; first pair of chin-shields forming a long suture behind mental ; throat-scales small, smooth, imbricate.

Scales.—Back covered with small convex granules of somewhat unequal size, with 12 longitudinal rows of much larger strongly keeled tubercles ; these much larger than ear-opening ; dorsal surface of limbs covered with unequal keeled granules ; ventral scales rather large, smooth, imbricate ; tail covered above and below with smooth imbricate scales which are a little larger on the ventral than on the dorsal surface.

Limbs short, the adpressed hind limb barely reaching the axilla. Fingers and toes short, free ; 8 lamellae under 4th, 4 under inner toe. Distal joint of inner digit extremely short.

Tail short, somewhat depressed as a whole, triangular in vertical section.

Colouration.—Dorsal surface dull olivaceous, with a dark longitudinal line extending on either side from the tip of the snout above the eye to the base of the tail ; this line followed below by a pale one and then by a second dark one ; two very narrow dark lines separated by a pale interspace below the second broader one on the sides of the belly ; tail dark olivaceous ; ventral surface yellowish green, suffused with dark olivaceous on the tail.

Total length		70 mm.
Length of head and body		33 „
Length of tail		37 „
Length of head 10 „
Length of fore limb 10 „
Length of hind limb 8 „

Habitat.—Bilimora, Bombay Presidency (*T. Bainbrigge Fletcher*), 13-ii-11.

Type.—No. 17020, Ind. Mus. (♀).

We are indebted to Mr. T. Bainbrigge Fletcher for the unique specimen of this curious little lizard, which appears to be very distinct from any hitherto described. It belongs to the section of the genus typified by *H. frenatus*, D. & B., and is perhaps nearer the S. Indian *T. reticulatus*, Beddome, than any other species. It may be distinguished from that form by its flattened head, broad rostral and longitudinal dark stripes.

***Japalura bengalensis*, sp. nov.**

(Plate v, fig. 4.)

Japalura yunnanensis, Annandale (*nec* Anderson), *J.A.S.B.*, 1906, p. 288.

In the paper cited I referred to the specimen here described as the male of *Japalura yunnanensis*, Anderson, but having since had an opportunity of examining an example of that species and having partially dissected the one from Buxa, I find that the latter is a female and differs from the Chinese form. It may be described as follows :

Habit stout ; body moderately compressed ; size large.

Head triangular ; its dorsal surface sloping downwards and forwards from behind the eyes, slightly concave between the orbits ; scales of dorsal surface irregular, keeled, largest on snout ; 9 upper and 9 lower labials ; snout longer than diameter of eye, bluntly pointed. A small gular pouch, no transverse fold across the throat.

Scales of back and sides small, almost granular, mixed with much larger keeled scales which tend to be grouped in small patches ; no dorso-lateral rows of enlarged scales ; ventrals larger than largest dorsals, strongly keeled, imbricate, pointed behind ; scales on upper surface of limbs irregular in size, leaf-shaped, strongly keeled.

Crest.—A well-developed nuchal crest (in the female) consisting of upright lanceolate scales shorter than half the diameter of the eye rising from a fold of skin covered with almost granular scales ; no dorsal crest on the anterior half of back, a very slight one on the posterior half.

Limbs moderate. Adpressed hind limb reaches anterior border of orbit ; tibia about as long as skull ; third and fourth fingers equal.

Colouration.—Head dull olivaceous with two narrow brown cross-bars on the dorsal surface, one just in front of, the other just behind the eye ; dark lines radiating from the eye ; lips with dark vertical bars ; sides of head irregularly marked with dark brown ; back and sides (in spirit) livid bluish-grey with a fine reticulation of dark brown ; tail obscurely barred ; limbs marked irregularly ; ventral surface yellowish ; a black patch on the gular pouch.

Total length	.	..	310 mm.
Length of head and body			110 "
Length of tail			200 "
Length of head			33 "
Length of fore limb			55 "
Length of hind limb	.	.	85 "

Locality.—Buxa, Jalpaiguri district, Bengal frontier of Bhutan.

Type.—No. 12564, Ind. Mus. (♀)

It is probable that this species inhabits the foot-hills of Bhutan, perhaps replacing *J. variegata* east of the R. Tista and being replaced by *J. andersoniana* at some unknown point situated still further east.

J. bengalensis is one of the largest species in the genus and easily surpasses either the Sikhim, the Dafla or the Khasi form in this respect. It is distinguished from the first (*J. variegata*) not only by its larger size but also by the nature of its crest, its colouration and scale-characters. From *J. planidorsata* the shape of its body will readily distinguish it, while its hind limbs are much shorter than in *J. andersoniana*.

***Tropidophorus assamensis*, sp. nov.**

This species is closely related to *Tropidophorus cochinensis*, D. & B., from which it differs in its more slender form, shorter limbs, keeled ventrals, etc.

Habit slender ; cylindrical.

Head.—Snout sharply pointed, narrow, no longer than orbit. Dorsal scales strongly ridged ; a single fronto-nasal, which is almost as broad posteriorly as it is long ; praefrontals short, forming a median suture ; frontal shorter than fronto-parietals and interparietal together ; 4 supraoculars, the 1st and 4th longest ; 5 supraciliaries, 1st longest ; fronto-parietals longer than interparietal, forming a suture behind it ; 5 upper labials, 4th longest ; 3rd, 4th and 5th beneath eye ; an azygous mental ; tympanum smaller than eye-opening.

Scales, 30 round body, all strongly keeled and spinously produced ; ventrals largest ; throat scales not strongly keeled but ending in a sharp point posteriorly ; a pair of large praenals ; dorsal and lateral tail-scales keeled ; ventral tail-scales smooth.

Limbs short but well-formed. Hind limb reaches wrist. Infra-digital lamellae smooth.

Tail cylindrical, tapering, longer than head and body.


Colouration.—Dorsal surface dark brown obscurely marbled with yellow, three cross-bars of latter colour ; one across hips, a second across shoulders and a third (less distinct than others) across back of neck ; tail rather darker, obscurely banded ; head brown ; sides and lips black with small white spots ; ventral surface yellowish ; small black spots on chin, throat and chest ; ventral surface of tail dark with an interrupted mid-ventral pale stripe, which becomes obscure distally, and numerous small yellowish spots.

Total length	90 mm.
Length of tail	51 ,,
Length of head	9 ,,
Breadth of head	5 ,,
Length of fore limb	. 12 ,,
Length of hind limb	. 14 ,,

Type.—No. 17029, Ind. Mus.

Locality.—Haraigaj range, 550 ft., Sylhet hills, Assam.

A single specimen was obtained by Mr. G. Mackrell, who has kindly presented it to the Indian Museum. It differs widely from *T. beramorii*, of which we possess the type. *T. yunnanensis*, Boulenger, is in my opinion synonymous with the latter species, of which I have examined many Burmese specimens. The scales seem to vary greatly in the degree to which they are keeled and in some cases are quite smooth, while the number of longitudinal rows of scales is not by any means constant.



A. C. Chowdhary, del.

1. *Typhlops diversiceps*, sp. nov. 2. *Aproaspidops antecursorum*, gen. et sp. nov.

3. *Ablabes pavo*, sp. nov. 4. *Japalura bengalensis*, sp. nov.

5. *Lygosoma courcyanum*, sp. nov.

III. LEPIDOPTERA.

By CAPTAIN W. H. EVANS, R.E.

The collection of Lepidoptera obtained on the Abor Expedition is not a large one, a fact which is doubtless explained by the season during which the military operations were undertaken. In addition to the specimens found by Mr. Kemp, a collection made by Captain H. W. Price of the Supply and Transport Corps and presented by him to the Indian Museum, has been examined while the records of other specimens collected by Major E. H. Sweet of the 2nd Gurkha Rifles are included. Permission to incorporate identifications of the species comprised in the latter collection was obtained through the courtesy of Major E. H. Sweet and the Secretary of the Bombay Natural History Society.

In all seventy-four species have been examined. The order and nomenclature followed in the list is that adopted in my "List of Indian Butterflies" published in the *Journal of the Bombay Natural History Society*, March 31st, 1912.

1. *Danaïa aglea melanoides*, M.
1 ♂ Sadiya : Nov. (S.W.K.); 1 ♀ Kobo : Dec. (S.W.K.).
1 ♂ " Abor Expedition " (E.H.S.).
2. *Danaïa melaneus*, Cr.
1 ♀ Between Kobo and Janakmukh (H.W.P.).
3. *Danaïa plexippus*, L.
1 ♂ " Abor Expedition " (E.H.S.).
4. *Euploea diocletianus*, Fab.
1 ♂ Between Kobo and Janakmukh (H.W.P.).
5. *Ypthima baldus*, Fab.
1 ♂ " Abor Expedition " (E.H.S.).
6. *Ypthima affectata*, El. and Ed.
1 ♂ Rotung : March, dry season form (S.W.K.).

Above the specimen is uniform dark brown on the forewing with no male brand : the hindwing has all the ocelli showing and some white powdering on the margin, especially towards the anal angle. Below the ground colour is very dark brown, loosely powdered pale ochreous : the ocelli on the hindwing are minute : there are no distinct bands, but the powdering is darker on the disc of the hindwing, showing up as a dull ochreous irregular patch. The underside resembles in general appearance the underside of *Y philomela indecora*, M., from the Western Himalayas.

7. *Lethe chandica*, M.
1 ♂ Rotung : March (S.W.K.).

8. *Mycalesis perseoides khasia*, Evans.
1 ♂ "Abor Expedition," dry season form (*E.H.S.*).
9. *Mycalesis visala*, M.
1 ♂ "Abor Expedition," dry season form (*E.H.S.*).
10. *Mycalesis meda*, Fab.
1 ♂, 1 ♀ "Abor Expedition," dry season forms (*E.H.S.*).
11. *Melanitis leda ismene*, Cr.
3 ♂, 2 ♀ "Abor Expedition," dry season forms (*E.H.S.*).
12. *Melanitis phedima bela*, M.
1 ♂ "Abor Expedition," dry season form (*E.H.S.*).
13. *Hestina nama*, Db.
1 ♂ "Abor Expedition" (*E.H.S.*).
14. *Stibochiona nicea*, Gray.
1 ♂ Rotung : March (*S.W.K.*) ; 1 ♀ between Kobo and Janakmukh (*H.W.P.*).
15. *Liminitis procris*, Cr.
1 ♂, 1 ♀ Kobo : Dec. (*S.W.K.*).
16. *Pantoporia zeroca*, M.
1 ♂ Sadiya : Nov. (*S.W.K.*).
17. *Neptis hylas varmona*, M.
1 ♂ Dibrugarh : Nov. (*S.W.K.*).
18. *Neptis soma*, M.
2 ♂ Sadiya and Kobo : Nov. (*S.K.W.*).
19. *Neptis hordonia*, Stoll.
1 ♂ Kobo : Dec. (*S.W.K.*).
20. *Cyrestis thyodamas*, Bdl.
2 ♂ Sadiya : Nov. (*S.W.K.*).
21. *Junonia iphita*, Cr.
1 ♂ Sadiya : Nov. (*S.W.K.*).
22. *Vanessa indica*, Herbst.
1 ♂ Kobo : Dec. (*S.W.K.*).
23. *Symbrenthia hippoclus khasiana*, M.
1 ♂ Sadiya : Nov. (*S.W.K.*).
24. *Rhinopalpa polynice birmana*, Fruh.
4 ♂ Kobo : Nov. and Dec. (*S.W.K.*) ; and between Kobo and Janakmukh (*H.W.P.*).

This insect is rare in Assam generally but seems common in the Abor country.

25. *Hypolimnas elina*, L.
1 ♂ Between Kobo and Janakmukh (*H.W.P.*).
26. *Kallima inachus*, Bdl.
1 ♂ Kobo ; Dec. (*S.W.K.*) ; 2 ♂ between Kobo and Janakmukh (*H.W.P.*).

27. *Cynthia erota*, Fab.
1 ♂ Between Kobo and Janakmukh (*H.W.P.*).
28. *Cirrhochroa aoris*, Db.
1 ♂ " Abor Expedition " (*E.H.S.*); 1 ♂ Sadiya : Nov. (*S.W.K.*); 1 ♀ Rotung : Dec. (*S.W.K.*).
29. *Argynnis hyperbius*, Joh.
1 ♂ Kobo : Dec. (*S.W.K.*); 1 ♀ Sadiya : Nov. (*S.W.K.*).
30. *Pseudergolis wedah*, Koll.
3 ♂ Kobo : Nov. and Dec. and Sadiya : Nov. (*S.W.K.*).
31. *Cethosia cyane*, Dr.
2 ♂ , 1 ♀ Kobo : Dec. (*S.W.K.*).
32. *Cethosia biblis*, Dr.
1 ♀ Rotung : Dec. (*S.W.K.*).
33. *Zemeros flegyas*, Cr.
2 ♂ " Abor Expedition " (*E.H.S.*); 1 ♂ Sadiya : Nov. and 2 ♂ Rotung : Dec. and March (*S.W.K.*).
- The specimens have the white apical spots well developed, showing a close resemblance to the race *confucius*, Fruh., from Southern Burma.
34. *Papilio philothenus polycuctes*, Db.
1 ♂ Rotung to Renging (*S.W.K.*).
35. *Papilio helenus*, L.
1 ♂ " Abor Expedition " (*E.H.S.*); 2 ♀ between Kobo and Janakmukh (*H.W.P.*).
36. *Papilio polytes romulus*, Cr.
1 ♂ " Abor Expedition " (*E.H.S.*).
37. *Papilio memnon agenor*, L.
1 ♀ form *butlerianus*, Roth., between Kobo and Janakmukh (*H.W.P.*).
38. *Papilio protenor euprotenor*, Fruh.
1 ♀ Sadiya : Nov. (*S.W.K.*).
39. *Papilio paris*, L.
1 ♀ Between Kobo and Janakmukh (*H.W.P.*).
40. *Leptocircus curius*, Fab.
1 ♀ Sadiya : Nov. (*S.W.K.*).
41. *Delias aglaia*, L.
2 ♂ Kobo : Dec. (*S.W.K.*).
42. *Pieris canidia*, Spaw.
1 ♂ Kobo : Dec. (*S.W.K.*).
43. *A. pias lalage*, Db.
1 ♂ " Abor Expedition " (*E.H.S.*); 1 ♂ Sadiya : Nov., 3 ♂ Rotung : Dec. and 1 ♂ Kobo : Dec. (*S.W.K.*).
44. *Dercas verhueli doubledayi*, M.
1 ♂ Rotung : March (*S.W.K.*).

45. *Catopsilia florella*, Fab.
1 ♂ Sadiya : Nov. (S.W.K.).
46. *Colias fieldii*, Mén.
2 ♀ near Kalek, 2,500 ft., March (S.W.K.).
- The specimens from the locality should be true *fieldii*, but are small and dull resembling the western *edusina*, But., more than the Chinese *fieldii*, Mén.
47. *Terias hecabe*, L.
3 ♂ Kobo : Dec. and 1 ♀ Sadiya : Nov., dry season forms (S.W.K.).
48. *Terias silhetana*, Wall.
1 ♀ Kobo : Dec., dry season form (S.W.K.).
49. *Hebomoia glaucippe*, L.
1 ♀ Between Kobo and Janakmukh (H.W.P.).
50. *Pareronia avatar*, M.
1 ♂ " Abor Expedition " (E.H.S.).
51. *Neopithecops zalmora*, But.
2 ♂ Rotung : Dec. and March (S.W.K.).
52. *Cyaniris transpecta*, M.
1 ♂ Sadiya : Nov. and 2 ♂, 1 ♀ Kobo : Dec. (S.W.K.).
53. *Cyaniris dilecta*, M.
1 ♂ Rotung : Dec. (S.W.K.).
54. *Cyaniris limbata placida*, de N.
2 ♂ Kobo : Dec., dry season forms (S.W.K.).
55. *Catochrysops strabo*, Fab.
1 ♂ Kobo : Dec. (S.W.K.).
56. *Castalius rosimon*, Fab.
2 ♂ Sadiya : Nov. (S.W.K.); 1 ♀ " Abor Expedition " (E.H.S.).
57. *Castalius elna*, Hew.
1 ♂ Sadiya : Nov. and 2 ♂, 2 ♀ Kobo : Dec. (S.W.K.).
58. *Lycaenesthes emolus*, God.
1 ♂ Sadiya : Nov. (S.W.K.).
59. *Nacaduba bhutea*, de N.
1 ♂ Kobo : Dec. (S.W.K.).
60. *Nacaduba nora*, Fd.
1 ♂ Rotung : Dec. (S.W.K.).
61. *Nacaduba noreia*, Fd.
1 ♂ Kobo : Dec. (S.W.K.).
62. *Lampides bochus*, Cr.
1 ♀ Sadiya : Nov. (S.W.K.).
63. *Lampides elpis*, God.
1 ♂ Kobo : Dec., typical and 1 ♂ Rotung : March (S.W.K.).

The specimen from Rotung has the border above reduced to a fine line, as in *pura*, M., and below all the markings very dull.

64. *Polyommatus boeticus*, L.
2 ♂ Rotung : March (S.W.K.).
65. *Poritia hewitsoni tavoyana*, Doh.
2 ♂, 2 ♀ Between Kobo and Janakmukh (H.W.P.).

The females have the lower part of the forewing very pale blue, while the orange spot, prominent in true *hewitsoni*, is only faintly indicated in one specimen and absent in the other.

66. *Ilerda epicles*, God.
3 ♂ Sadiya : Nov., 2 ♂ Kobo : Dec. and 1 ♂ Rotung : Dec. (S.W.K.).
67. *Arhopala silhetensis*, Hew.
1 ♀ Dibrugarh : Nov. (S.W.K.).
68. *Rapala schistacea*, M.
1 ♂ Kobo : Dec. (S.W.K.).
69. *Hypolycaena erylus*, God.
1 ♂ Kobo : Dec. (S.W.K.).
70. *Cheritra freja*, Fab.
1 ♀ Kobo : Dec., dry season form (S.W.K.).
71. *Zeltus etolus*, Fab.
3 ♂ Kobo : Dec. and 3 ♂ Sadiya : Dec. (S.W.K.).
72. *Loxura atymnus*, Cr.
1 ♂ Dibrugarh : Nov. and 1 ♂ Sadiya : Nov. (S.W.K.).
73. *Ampittia maro*, Fab.
1 ♂ " Abor Expedition " (E.H.S.).
74. *Rhopalocampta benjaminii*, Guér.
1 ♂ Rotung : March (S.W.K.).

IV. PORIFERA.

By N. ANNANDALE, *D.Sc., F.A.S.B., Superintendent of the Indian Museum.*

In the absence of ponds, lakes and slow-running streams it was not to be expected that sponges would be found in the Abor country, but two species were obtained by Mr. Kemp and Mr. J. Coggin Brown of the Geological Survey in N.-E. Assam. The only species hitherto found in Assam was *Spongilla (Euspongilla) proliferens*, Annandale, which was taken near Mangaldai to the north of the Brahmaputra by Mr. Kemp on a former occasion and is widely distributed in the Oriental Region. Specimens were found by Messrs. Kemp and Brown on their way to the Abor country at Dibrugarh on the south bank of the Brahmaputra. The same collectors obtained specimens of another sponge (*Spongilla (Eunapius) crassissima* var. *crassior*, Annandale) in a small muddy pond at Mariani in the Sibsagar district. This form had only been found hitherto at Rajshai (Rampur Bhulia) on the north bank of the Ganges in Bengal, while the typical form of the species is only known from the Gangetic delta and Orissa.

All the specimens from N.-E. Assam were preserved in November. Although apparently killed during a period of full vegetative vigour, they are full of gemmules.

I can detect no buds on the examples of *S. proliferens*, but these structures, although characteristic of the species, are not always present. The gemmules possess well-formed foraminal tubules.

The specimens of *S. crassissima* var. *crassior* agree well with the types except in being of a bright green colour. In external appearance (except as regards colour) they closely resemble *S. decipiens calcuttana*, which belongs to the same subgenus, but they can be easily distinguished therefrom by their much stouter megascleres and more compact skeleton.

S. proliferens is fully described on pp. 72-76 of my volume in the "Fauna" on the Freshwater Sponges, etc. and *S. crassissima* and its variety on pp. 98 and 99 of the same work.

V. SCOLOPENDRIDAE.

By F. H. GRAVELY, *M.Sc.*, Assistant Superintendent, Indian Museum.

Out of the eight species collected two are new, one is only known from the E. Himalayas and Assam ; but all the rest occur south and in most cases west of this area as well.

Subfamily *CRYPTOPINAE*.

Genus *Cryptops*.

C. doriae, Pocock.

Renging, 2,150 ft., 19-xii-11 (under bark).

This species has hitherto been recorded from the area between and including New Guinea and Burma. It must extend westwards right across India, however, for I recently obtained specimens from Taloshi (Satara District, Bombay) at an altitude of about 2,000 ft. in the Western Ghats.

Subfamily *OTOSTIGMINAE*.

Genus *Otostigmus*.

O. burn-murdochi, n. sp.

Upper Rotung, ca. 2,000 ft. Jan., 1912.

Collected by the 32nd Sikh Pioneers when roadmaking and named after Mr. I. Burn-Murdoch of that regiment.

Description.—Length (apart from appendages) up to 63 mm. Antennae 19-jointed (rarely 17-jointed on one side and 19-jointed on the other, the terminal joint being present in both), the first three joints large, polished and hairless, the rest smaller and pilose. Dorsal surface of head and body dark olive green, finely and sparsely but evenly punctured throughout, the punctures on the head and first one or two segments often a trifle larger and less clearly defined than the rest ; segments 3-20 with a pair of very fine and sometimes incomplete longitudinal grooves ; these grooves usually more strongly impressed close to the posterior margin especially in the anterior or middle segments of the body, where they are often joined by a transverse groove between which and the posterior margin the surface may be rugose. Marginal grooves present behind segments 8-11. Anal segment laterally with strong marginal ridges ; posterior margin composed of two straight halves meet-

ing in an angle of about 120° Ventral surface paler than dorsal, similarly punctured, but without grooves; sternocoxal plate armed with five strong teeth (sometimes four or six on one side only). Sternum of anal segment narrower behind than in front, posterior margin more or less concave. Pseudopleural processes at least as long as sternum of anal segment, sometimes very slender behind, armed with 1-2 distal, 1-2 external, and 1-4 dorsal spinules. Tooth on inner side of femur of poison-jaws strongly developed. Legs distally of a more bluish tinge than the body. First pair of legs armed with 1 distal spine on the patella, 1 on the tibia, 2 on the proximal tarsus, and 2 at the base of the claw; second with 2 spines on the proximal tarsus and claw, with or without 1 on the tibia; third to twentieth with 2 at the base of the claw, 1, or usually in many of the anterior segments 2, at the end of the proximal tarsus; anal legs with 2 at the base of the claw only; femur of anal legs with a strong dorsal tooth on the inner side at the end ("eckdorn") with 1-3 smaller teeth behind it, beneath on the outer side with 3, on the inner with 2, 2-3, 3 teeth often irregularly arranged.

This species is allied to those grouped under the number 5 in Kraepelin's table [Revision der Scolopendriden, p. 99, in Mitt. Naturhist. Mus. Hamburg, XX, 1902 (1903)]. It differs from all of them, however, in combining the presence of external spinules on the pseudopleural processes with the absence of longitudinal grooves on the sterna.

O. insularis, Haase.

Kobo, 400 ft., 8-xii-11.

Rotung, 1,300-1,400 ft., 31-xii-11 and 1-i-12.

This species is recorded from the Seychelles, Ceylon, the Phillipines and the E. Himalayas.

Several specimens have recently been added to the Museum collection. In addition to the Abor records above noted, the following localities may be given:—

Ceylon: Kandy and Peradeniya, 1,500-2,000 ft.

E. Himalayas: Darjiling District—Ghumti, ca. 1,800 ft.

Assam: Mangaldai District—Assam-Bhutan Frontier. Dikrang Valley.

Malay Peninsula: Penang.

Individuals differ greatly one from another in many respects. Kraepelin has pointed out that the Himalayan form differs from the Ceylon one in having only two instead of two and half segments of the antenna smooth and hairless. This is confirmed by the specimens I have seen, the Penang specimen agreeing with the Himalayan ones in this respect, but as two of the Himalayan ones have two and half segments bare on one antenna I am inclined to think he was right in not giving the two forms separate varietal names. Great differences are also found in the size, proportions, and femur-armature of the anal legs. Our Ceylon specimens agree more or less perfectly with Kraepelin's description,

but in the Himalayan ones the number of spines on the femur is commonly less and the femur as a whole is apt to be shorter ; neither of these characters seems, however, to be constant nor are the shortest femora necessarily those with the fewest spines. This frequent reduction of the number of spines on the femora brings the species still nearer to *O. scaber*, Pocock, and although I have never yet seen a specimen with two tarsal spurs on the legs of less than about eighteen segments I cannot help thinking that the two will in course of time prove to be identical. The pseudopleural processes are also subject to variation, and the distinctness of the longitudinal keels, as well as the presence or absence of a pair of longitudinal grooves and the whole texture of the dorsal surface, is extraordinarily variable. The dorsal surface may be smooth and polished between the keels throughout, or may be granular or finely spinulose laterally, or finely spinulose throughout.

***O. rugulosus*, Por.**

Sadiya, N.-E. Assam, 25-xi-11.

Kobo, 400 ft., 30-xi—3-xii-11 (rotten wood).

Rotung, 1,300 ft., 30-xii-11 (under stones).

Renging, 2,150 ft., 19-xii-11 (rotten wood).

Previously recorded only from the Seychelles, Mauritius, Andamans, Burma, and Siam. There are however also specimens in the Indian Museum collection which were recently collected by Mr. Kemp in Assam as far west as Mangaldai and the Assam-Bhutan frontier north of that district.

The extent to which the longitudinal grooves of the sterna are developed varies greatly ; and in one or two of the Kobo specimens they are entirely absent.

Genus ***Rhysida***, Wood.

***R. nuda* (Newp.).**

Dibrugarh, N.-E. Assam, 17—21-xi-11.

Sadiya, N.-E. Assam, 25-xi-11.

This species occurs throughout the Oriental Region and also in Australia and Paraguay. The specimens from the Abor country are nearer *R. immarginata* (Por.) than *R. nuda* as these species are defined by Kraepelin (loc. cit., pp. 141 and 143-4), but as specimens combining the characters of the two seem to be far commoner in India than the latter and quite as common as the former, I feel unable to separate them, and Newport's name has priority over Porat's. Complete or incomplete marginal grooves are often present on several segments in front of the twenty-first, which alone has strong marginal ridges.

R. cuprea, Krpln.

Kobo, 400 ft., 30-xi—3-xii-11 (one in rotten wood and one in earth).
Balek, 600 ft., 24-iii-12.

Up to the present the only published record for this species is Bhutan. The Indian Museum has, however, recently received specimens from the following localities :—

E. Himalayas : Darjiling District—Kurseong, *ca.* 5,000 ft. ; Ghumti Tea Estate, *ca.* 4,000 ft.

Assam : Sylhet—Shamshernagar, *ca.* 100 ft.

This species is much nearer *R. stuhlmanni*, Krpln., and *R. petersi* (Por.) from Africa than Kraepelin supposed ; for the fourth joint of the antenna is often broader than long, the first stigmata may approach an S-shape, the pseudopleura may be terminated by 3 instead of 2 spines, and the nineteenth and twentieth legs may have 2 and 1 instead of 1 and 0 tarsal spurs respectively. It can be distinguished however by its 20-21-jointed antenna. In one specimen one of the antennae is only 18-jointed and has the terminal segment present, but presumably this is due to regeneration after mutilation.

Genus **Ethmostigmus**, Poc.**E. pygomegas** (Kohlr.).

Kobo, 400 ft. (rotten wood).

Upper Rotung, *ca.* 2,000 ft. (collected by M. de Courcy).

This species is found throughout the E. Himalayas, Assam, Burma and the Nicobars.

Subfamily SCOLOPENDRINAE.

Genus **Scolopendra** (L.) Newp.**S. mazbii**, n. sp.

Upper Rotung, *ca.* 2,000 ft., Jan., 1912.

One small specimen was found under the leaf-stem of a plantain ; the rest were obtained by Capt. the Hon. M. de Courcy and the 32nd Sikh Pioneers when road-making ; the name chosen for the species is taken from that of the caste to which the men of this regiment belong.

Description.—Length (apart from appendages) up to 95 mm. Antennae 17-jointed, the first five joints hairless, the rest uniformly pubescent throughout. Dorsal surface of head dark bluish in front, pale greenish behind, marked with a fine longitudinal groove in the middle line. First segment also pale greenish above, with or without the fine median groove ; second and third segments dark bluish with a pair of distinct longitudinal grooves ; remaining segments with the exception of the last olivaceous

in front with a dark bluish band along the posterior margin¹ and a pair of distinct longitudinal grooves throughout, about half of these segments with raised lateral margins ; last segment olivaceous, little darker behind than in front, entirely without paired or median grooves, lateral margins very strongly raised, posterior margin curved. Ventral surface uniformly olivaceous or with head and first segment browner or paler than the rest ; sternocoxal plate armed with 5 minute and widely separated teeth, sterna of segments 2-19 or 20 with a pair of distinct longitudinal grooves ; sternum of the last segment rather slender, narrower behind than in front, posterior margin straight or very nearly so. Pseudopleural processes conical, rather slender, the apex simply pointed. Tooth on inner side of femur of poison-jaws strongly developed ; first pair of legs paler in colour than the rest, legs 1-19 armed with 1 tarsal and 2 claw spurs, the twentieth with 2 claw spurs and 0(-1) tarsal spurs, the anal legs usually without spurs, sometimes, however, with 1 claw-spur. Femur of anal legs armed normally with 2, 2 ventral outer, 2 ventral inner, 2 dorsal inner and 1 long and always simply pointed posterior dorsal spine ; the arrangement, however, is sometimes less regular than this when one or two additional spines are developed ; and the proximal ventral spine is often so much reduced that it is not improbable that it may occasionally be absent. Distal tarsus of anal leg moderately slender in young specimens ; much stouter proportionally in adults in which it is sometimes actually shorter than the very large claw.

This species comes next to *S. morsitans* when compared with Kraepelin's table (loc. cit., pp. 226-232), but can be distinguished therefrom at once by the simply pointed pseudopleural processes and the similarly simple posterior dorsal spine ("eckdorn") of the anal legs as well as by other minor characters.

¹ Mr. Kemp informs me that the colours were much more brilliant in life, the ground colour which is now olivaceous being then a straw-yellow.

VI. HYMENOPTERA ANTHOPHILA.

By C. A. PAIVA, *Assistant, Indian Museum.*

Among the Hymenoptera collected during the Abor Expedition there are 40 specimens of Bees, most of which appear to be fairly widely distributed. I have been able to accurately identify most of them, but there are a few the identity of which I am not quite sure about. I therefore propose to leave them undetermined for the present.

1. *Sphcodes*, sp.

One male collected at Sadiya, N.-E. Assam, on the 27-xi-11 (*Kemp*).
A genus very widely distributed in both hemispheres.

2. *Halictus*, spp. (♂ ♂).

There seem to be at least two species among the 3 specimens of this genus which were collected at Sadiya, 28-xi-11, and Dibrugarh, 11—19-xi-11, N.-E. Assam and at Kobo, 400 ft., 3-xii-11 (*Kemp*). Those from Sadiya and Kobo come very near *H. funebris*, Cam., and the one from Dibrugarh agrees with the description of *Halictus pulchriventris*, Cam., differing only in having the 2nd joint of the flagellum of the antenna black beneath instead of testaceous yellow.

There is also a very badly preserved female belonging to this genus collected at Sadiya, N.-E. Assam, 28-xi-11 (*Kemp*).

3. *Halictus rufo-zonatus*, Vachal.

Ann. Mus. Civ. Genoa (2a), 1894, p. 442, ♀; Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 439, ♀.

Six females were found under bark at Renging, ca. 2,150 ft., 19-xii-11 (*Kemp*).

Bingham records the species from Bhamo, Upper Burma.

4. *Andrena mollis*, Smith.

New Sp. Hym. B. M., p. 50, ♀; Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 446.

Three specimens were collected at Sadiya, N.-E. Assam, 28-xi-11, and two at Dibrugarh, N.-E. Assam, 17—19-xi-11 (*Kemp*).

Recorded by Bingham from Bombay, Assam, Burma and Tenasserim.

Represented in the Indian Museum collection from United Provinces :—Jhansi, 850 ft., 2-viii-05 (*Brunetti*).

U. Burma :—Rangoon, 12-vi-97 (*Bingham*).

E. Himalayas :—Sukna, 500 ft., 2-viii-08 (*Annandale*).

A very variable species.

5. *Nomia aurifrons*, Smith.

Trans. Ent. Soc. London, 1875, p. 43, ♀; Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 455; Paiva, Rec. Ind. Mus., i, pt. i, 1907, p. 16.

One specimen from Dibrugarh, N.-E. Assam, 11—19-xi-11 (*Kemp*).

Recorded by Bingham from Bengal, Sylhet, Burma and Tenasserim.

Represented in the Indian Museum collection from Katmandu, Nepal Valley, 4,500 ft., Oct., 1906, and from Nagarkote, Nepal, ca. 6,000 ft., Oct., 1906 (*Hodgart*).

Represented in the Pusa collection from Bihar :—Pusa, Chupra.

6. *Nomia terminata*, Smith.

Trans. Ent. Soc. London, 1875, p. 56; Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 459; Paiva, Rec. Ind. Mus., i, pt. i, 1907, p. 16.

Three specimens taken at Kobo, 400 ft., 6—8-xii-11 (*Kemp*).

Recorded by Bingham from Sikhim, Burma and Tenasserim.

Represented in the Indian Museum collection from Assam :—Margherita.

E. Himalayas :—Sureil, Darjiling district, 5,000 ft., April, 1905 (*Alcock*).

7. *Lithurgus dentipes*, Smith.

Cat. i, p. 146, ♂; Horne, Trans. Zool. Soc. Lond., vii, 1872, p. 175; Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 470.

One specimen from Sadiya, N.-E. Assam, 27-xi-11 (*Kemp*).

Recorded by Bingham from Sikhim; Manipuri, N.-W P. (*Horne*); Tenasserim and New Holland (*Smith*).

Not previously represented in the Indian Museum collection.

Represented in the Pusa collection from Bihar :—Pusa, Chupra.

8. *Ceratina hieroglyphica*, Smith.

Cat. ii, 226, ♀ ♂; ? *Ceratina flavopicta*, Morawitz (*nec* Smith), Hor. Soc. Ent. Ross., xxiv, 1890, p. 356. *Ceratina hieroglyphica*, Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 503, pl. iv, fig. 3; Rec. Ind. Mus., ii, 1908, p. 365.

One specimen from Sadiya, N.-E. Assam, 28-xi-11 (*Kemp*).

Bingham records it from Mussoorie, Dehra, Barackpore, Sikhim, Bangalore, Assam, Burma and Tenasserim extending into China and the Malayan region.

Represented in the Indian Museum collection from Kashmir 8,000—9,000 ft.

W. Himalayas :—Mussoorie, ca. 7,000 ft., 20—26-v-05 (*Brunetti*).

Bihar :—Barh (*Burkill*).

Bengal :—Calcutta, Khulna, 10-iii-07 (*Caunter*).

Assam :—Sadiya and Margherita.

Borneo :—10 miles south of Kuching, Sarawak, 24-vi-10 (*C. W. Beebe*).

Represented in the Pusa collection from Bihar :—Pusa, Chupra, Daulatpur, Darbhanga.

W. Himalayas :—Mussoorie, 7,000 ft.

Bombay :—Kanara, Belgaum, 2,500 ft.

9. *Crocisa emarginata*, Lepel.

Lepelletier, Hym., ii, p. 449, ♀ ♂; Smith, Cat. Hym. Ins. B.M. ii, p. 277; Journ. Linn. Soc., xi, 1876, p. 390. *Crocisa decora*, Smith, Trans. Ent. Soc. London, n. s., ii, 1852, p. 41, ♀. *Crocisa elegans*, Smith (*nec* Moscary), New species Hym. B. M., p. 107, ♀. *Crocisa emarginata*, Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 517; Proc. Zool. Soc. London, 1896, p. 454; Paiva, Rec. Ind. Mus., i, 1907, p. 16.

One specimen from Dibrugarh, N.-E. Assam, 19-xi-11 (*Kemp*).

Recorded by Bingham from Sikkim, Calcutta, Bombay, Malabar, Ceylon, Pundaloya (*Green's coll.*), Burma and Tenasserim; on the west extending to South Africa, on the east to China and the Malay Archipelago.

Represented in the Indian Museum collection from Bombay :—Poona Ghats.

United Provinces :—Lucknow.

Bengal :—On the road from Magra to Trebeni, Hooghly district, 31-vii-09 (*Mus. collr.*); Basanti Forest Station, Sunderbuns, L. Bengal, 16-xi-09 (*Jenkins*); Calcutta, 7-xi-06 (*Brunetti*), 9-vi-09 (*Mus. collr.*).

Nepal :—Soondrijal, Nepal Valley, 1906 (*Hodgart*).

Assam :—Sibsagar; Margherita; Naga Hills.

S. India :—Bangalore; Oorgaum, ca. 2,500 ft., 20-x-10 (*Mus. collr.*).

U. Burma :—Between Tengueh and Tali Fu, Yunnan, W. China, 1909-10 (*J. Coggin Brown*).

L. Burma :—Moulmein; Mergui; Tenasserim.

Ceylon : Kandy, vi-10.

Malay Archipelago :—Sinkep I.

Malay Peninsula :—Perak.

Represented in the Pusa collection from S. India :—Naduvatom, 7,000 ft., Nilgiris.

Bengal :—Madhyapara, Dacca district.

Bombay :—Belgaum, 2,500 ft.

10. *Habropoda fulvipes*, Cam.

Cameron, Ann. Mag. Nat. Hist., xiii (7th series), 1904, p. 211.

One specimen in very good condition taken in a jungle path, Sadiya, N.-E. Assam, 23-xi-11 (*Kemp*).

Originally described from Khasi Hills (*Rothney coll.*).

Not previously represented in the Indian Museum collection.

11. *Anthophora zonata*, Linn.

Apis zonata, Linn. Syst. Nat. (ed. 10), i, p. 576. *Anthophora subcaerulea*, Lepel., Hym., ii, p. 30, ♀♂. *Anthophora zonata*, Bingham, Proc. Zool. Soc. Lond., 1896, p. 455; Faun. Brit. Ind. Hym., i, 1897, p. 527; Paiva, Rec. Ind. Mus. i, 1907, p. 16.

One specimen from Sadiya, N.-E. Assam, 23-xi-11 (*Kemp*).

Recorded by Bingham from all over India, Burma, Tenasserim, and Ceylon, extending through the Malay regions to Australia.

Represented in the Indian Museum from W. Himalayas :—Mussoorie, ca. 7,000 ft., 20—24-vi-05 (*Brunetti*); Almora, Kumaon, ca. 5,500 ft., 15-vii-11 (*Paiva*); Kalka, base of Simla Hills, alt. 2,400 ft., 18-vii-11 (*Mus. collr.*).

United Provinces :—Dehra Dun base of W. Himalayas; Kaladhunji, Naini Tal district (plains) 20-iv-09 (*Mus. collr.*); Meerut, 25-iv-05 (*Brunetti*).

E. Himalayas :—Sikhim; Siliguri, base of E. Himalayas.

Bihar :—Rajmahal, 5-vii-09 (*Annandale*); Barh (*Burkill*).

Chota Nagpur :—Paresnath 4,300—4,400 ft., 15-iv-09 (*Annandale*).

Bengal :—Murshidabad; Calcutta, 23-iv-07, 27-v-09, 9-vi-09; Netrakona, Mymensingh district, 4-x-09 (*Bagchi*).

Assam :—Margherita; Sadiya; Naga Hills; Sibsagar; Mazbat, Mangaldai district, 11—15-ix-10 (*Kemp*).

Burma :—Tenasserim Valley.

Madras :—Gopkuda I., Lake Chilka, Ganjam, N.-E. Madras, 7—15-viii-07; S. end of Lake Chilka, N.-E. Madras, 4-iii-10 (*Annandale*).

S. India :—Bangalore; Shasthancottah, 12 miles N.-N.-E. of Quilon; Trivandrum, 6-xi-08 (*Annandale*); Ayasamtengi, S. end of Lake Kayankulum, Travancore, 6-xi-08; Maddathoray, W. base of W. Ghats, Travancore, 17-xi-08 (*Annandale*); Trivandrum, August, 1903, November, 1890.

Ceylon :—Peradeniya, 10-vii-10, 24-v-10, June, 1910.

Malay Peninsula :—Perak.

Represented in the Pusa collection from Bihar :—Pusa, Chupra.

Orissa :—Cuttack.

Central Provinces :—Bilaspur.

Bombay :—Matheran, 2,500 ft.

12. *Xylocopa latipes* (Drury).

Apis latipes, Drury, Ill. Exot. Insects, ii, pl. 48, fig. 2. *Xylocopa latipes*, Fab., Syst. Piez., p. 337 ; Smith, Cat., ii, p. 353 ; Trans. Ent. Soc. Lond., 1874, p. 267 ; Bingham, Proc. Zool. Soc. Lond., 1896, p. 456 ; Faun. Brit. Ind. Hym., i, 1897, p. 536, pl. iv, fig. 7 ; Cameron, Faun. and Geo. Mald. and Lacc. Arch., vol. i, Hym., 1900-1903, p. 62.

One specimen from Sadiya, N.-E. Assam, 27-xii-11 (*Kemp*).

Recorded by Bingham from Sikhim, Barrackpore, Kumaon, Terai, Central India, Malabar, Coonor, Assam, Burma, Tenasserim extending to China and the Malayan region.

Cameron records it from Mamaduwari, Mahlos Atoll, Maldives.

Represented in the Indian Museum from E. Himalayas :—Sikkim ; Tindharia, 2,820 ft. (*Richardson*).

United Provinces :—Saharanpur.

Chota Nagpur :—Ranchi.

Bengal :—Murshidabad ; Calcutta.

Assam :—Shillong ; Margherita ; Naga Hills ; Sibsagar ; Mazbat, Mangaldai district, 11-15-x-09 (*Kemp*).

Burma :—Tenasserim ; Tavoy ; Mergui.

S. India :—Bangalore ; Trivandrum ; Travancore ; S. Malabar.

Ceylon :—Colombo, 25-vi-4.

Malay Archipelago :—Sinkep I.

Malay Peninsula :—Perak ; Johore.

Borneo :—Kapit, Sarawak, 21-vii-10 (*Beebe*) ; Sandakan.

Represented in the Pusa collection from Assam :—Naharkatia ; Nangpoh, 3,000 ft. to 5,000 ft., Khasi hills.

13. *Xylocopa collaris*, Lep.

Lepelletier, Hym., ii, p. 189, ♀ ; Smith, Cat., ii, p. 353 ; Journ.

Linn. Soc., xi, p. 393 ; Trans. Ent. Soc. Lond., 1874, p. 270.

Xylocopa dejeanii, Lepel., Hym., ii, p. 209, ♂ ; Smith, Cat.

Hym. Insects B.M., ii, p. 357 ; Journ. Linn. Soc., xi, p. 394.

Xylocopa collaris, Bingham, Proc. Zool. Soc. Lond., 1896, p. 456 ;

Faun. Brit. Ind. Hym., i, p. 543.

Seven females and two males from Yembung, 1,100 ft., taken on the 14-i-12, boring into rotten wood, and another taken at Rotung, 1,400 ft. in rotten wood on 23-xii-11 (*Kemp*).

Recorded by Bingham from Sikhim, Allahabad, Kumaon, Burma, Tenasserim, Ceylon (Pundaloya), extending into the Malay region and Siam.

Represented in the Indian Museum collection from Tibet (*Moller*).

W. Himalayas :—Simla.

E. Himalayas :—Sikhim.

Assam :—Naga Hills ; Sibsagar ; Sadiya ; Margherita.

Burma :—Mergui ; Upper Tenasserim.

S. India :—Trivandrum, Feb., 1893.

Malay Peninsula :—Johore.

Represented in the Pusa collection from Bengal :—Buxa Duars.

14. *Bombus orientalis*, Smith.

Cat. ii, p. 402, ♀. *Bombus buccinatoris*, Sm., New Sp. Hym. Ins. B.M., p. 132, ♀. ? *Bombus simulus*, Grib., Bull. Ent. Soc. Ital., xxiii, 1891, p. 114, ♀. *Bombus orientalis*, Bingham, Faun. Brit. Ind. Hym., i, 1897, p. 555, pl. iv, fig. 9 ; Paiva, Rec. Ind. Mus., i, 1907, p. 17.

Two specimens from Rotung, 1,400 ft., 26—27-xii-11, one from near Renging, 1,500 ft., 19-xii-11, and one from Sirpo, ca. 1,300 ft., March, 1912 (*Kemp*).

The specimens vary very much in size, the smallest is 12 mm., the next 15 mm. and the largest 26 mm. in length.

Recorded by Bingham from Sikhim, Kunawar. "One of the commonest species about Darjiling" (*Bingham*).

Represented in the Indian Museum collection from Nepal :—Katmandu, Nepal Valley, 4,500 ft., Oct., 1906 ; Soondrijal, Nepal Valley ; Chitlong, little Nepal Valley ; Chandragiri, ca. 8,000 ft., Oct., 1906 (*R. Hodgart*).

E. Himalayas :—Sikhim ; Kurseong, 4,700—5,000 ft., 22, 24, 25-vi-10, 5, 7, 9-ix-09 (*Annandale*) ; 6,000 ft., 13-x-09 (*D'Abreau*) ; Gumti, Darjiling district, ca. 4,000 ft., vii-11 (*Gravelly*).

Assam :—Cheera Punji, Khasi Hills (*Warren*).

W. Himalayas :—Kanaul, British Garhwal, 18-x-07 (*Mus. collr.*).

Represented in the Pusa collection from E. Himalayas —Phoobsering, Lebong, Darjiling district.

It might be useful to note that each of the species *Bombus orientalis* and *B. haemorrhoidalis*, which to the casual observer would appear alike, has its range of distribution quite distinct. *B. orientalis*, though recorded so far west as Kanaul in British Garhwal, is not met with in any of the western ranges of the Himalayas. It is found as far west as the western borders of Nepal and probably in some of the Kumaon Hills where the two species may meet. *B. haemorrhoidalis*, though described from Chusan in the hills of North China, has not yet been found in any Indian locality to the east of the Kumaon Hills. I found it in fairly large numbers in Almora, visiting roses.

15. *Apis dorsata*, Fab.

Fabricius, Ent. Syst., ii, p. 328 ; Syst. Piez., 370 ; Smith, Cat. ii, p. 415 ; Journ. Linn. Soc., xi 1876, p. 396 ; Horne, Trans. Zool. Soc. Lond., vii, p. 181, pl. 22, fig. 3 ; Lep., Hym., i, p. 405, ♀. *Apis testacea*, Sm., Journ. Linn. Soc., ii, 1858, p. 49. *Apis zonata*, Sm. (*nec Guér.*), Journ. Linn. Soc., iii, 1859, p. 8. *Apis dorsata*, Bingham, Proc. Zool. Soc. Lond., 1896, p. 457 ; Faun.

Brit. Ind. Hym., i, 1897, p. 557, pl. iv, fig. 11 ; Paiva, Rec. Ind. Mus., i, p. 17, 1907.

One specimen from Dibrugarh, N.-E. Assam, 17—19-xi-11 (*Kemp*).
Recorded by Bingham from throughout India, Burma, Tenasserim, Ceylon, extending into China and the Malayan region to Java.

Represented in the Indian Museum collection from W. Himalayas :—Mussoorie, ca. 7,000 ft., 12-viii-05 (*Brunetti*) ; Kumaon ; Simla, 7,000 ft., 7-v-10 (*Annandale*).

United Provinces :—Dehra Dun, base of W. Himalayas ; Dhikala, Naini Tal district, 22-iv-08 ; Boxar, Naini Tal district, 19-iv-08, Gularbojh, Naini Tal district, 17—20-iv-08 ; Bareilly 15—22-iii-07 ; Lucknow, 5-xi-07 ; Meerut, 8—14-iii-07 ; Kaira, base of W. Himalayas, Naini Tal district, 24-iii-10 ; Bhogpur, Bijnor district, 2-iii-10 ; Amangarh, Bijnor district, 24-ii-10 (*Mus. collr.*).

Nepal frontier :—Thamaspur, 18—20-ii-08 (*Mus. collr.*).

E. Himalayas :—Sikhim.

Chota Nagpur :—Paresnath, 4,400 ft., 11-iv-09 (*Annandale*).

Bengal :—Murshidabad ; Calcutta, 20-iii-08 (*Annandale*).

Assam :—Naga Hills ; Sibsagar ; Shillong ; Margherita ; Mazlat, Mangaldai district, 11—15-x-10 (*Kemp*) ; Lushai Hills, 3,600 ft., 7-vi-04 (*E. C. MacLeod*).

Burma :—Base of Dawna Hills, 2-iii 08 (*Annandale*).

Bombay :—Belgaum.

Malay Archipelago :—Sinkep I.

Represented in the Pusa collection from Punjab :—Lahore ; Wazirabad ; Jamoo.

Central Provinces :—Bula ; Hoshangabad ; Itarsi.

Bihar :—Chupra.

Bengal :—Rangpore.

REFERENCES.

- Bingham, C. T., 1896.—“ A contribution to the knowledge of the Hymenopterous Fauna of Ceylon.”—Proceedings of the Zoological Society of London.
- „ 1897.—Fauna of British India including Ceylon and Burma, Hymenoptera, vol. i (Wasps and Bees).
- „ 1908.—“ Aculeate Hymenoptera in the Indian Museum, I.”—Records of the Indian Museum, vol. ii, Calcutta.
- Cameron, P., 1900-1903.—Fauna and Geography of the Maldive and Laccadive Archipelago, vol. i, Hymenoptera.
- „ 1904.—“ Descriptions of new species of Aculeate and Parasitic Hymenoptera from Northern India.”—Annals and Magazine of Natural History, vol. xiii, 7th series, London.

- Drury, D., 1770-82.—Illustrations of Natural History, etc. Exotic Insects, London.
- Fabricius, J. C., 1804.—Systema Piezatorum, Brunswick.
- Gribodo, G., 1891.—“Sopra alcune specie nuove o poco conosciute di Imenotteri Antofili.”—Bulletino della Societa entomologica Italiana, Florence.
- Horne, C., 1872.—“Notes on the habits of some Hymenopterous Insects from the North-West Provinces of India.”—Transactions of the Zoological Society of London, vol. vii.
- Lepelletier, A. L., 1841-46—Histoire naturelle des Insectes Hyménopteres, 4 vols., Paris.
- Linné, C., 1758.—Systema Naturae, Ed. x, Stockholm.
- Morawitz, F. 1890.—“Hymenoptera Aculeata from China and Mongolia.”—Horae Societatis Entomologicae Rossicae, St. Petersburg.
- Paiva, C. A., 1907.—“Records of Hemiptera and Hymenoptera from the Himalayas.”—Records of the Indian Museum, vol. i.
- Smith, F., 1852.—“Descriptions of some new and apparently undescribed species of Hymenopterous Insects from North China, collected by Robert Fortune.”—Transactions of the Entomological Society of London, vol. ii, new series.
- „ 1853-54.—Catalogue of Hymenopterous Insects in the collection of the British Museum, London.
- „ 1858.—“Catalogue of the Hymenopterous Insects collected at Sarawak, Borneo, Mount Ophir, Malacca and at Singapore, by A. R. Wallace.”—Journal of the Proceedings of the Linnean Society, vol. ii, London.
- „ 1859.—“Catalogue of Hymenopterous Insects collected at Celebes, by Mr. A. R. Wallace.”—Journal of the Proceedings of the Linnean Society, vol. iii, London.
- „ 1874.—“Monograph of the Genus *Xylocopa*, Latr.”—Transactions of the Entomological Society of London.
- „ 1875.—“Descriptions of new species of Indian Aculeate Hymenoptera, collected by Mr. G. R. James Rothney.”—Transactions of the Entomological Society, London.
- „ 1876.—“A catalogue of the Aculeate Hymenoptera and Ichneumonidae of India and the Eastern Archipelago.”—Journal of the Proceedings of the Linnean Society, vol. xi, London.

- 1912.] C. A. PAIVA : *Hymenoptera Anthophila*. 83
- Smith, F., 1879.—Descriptions of new species of Hymenoptera in the collection of the British Museum, London, parts i and ii.
- Vachal J 1894.—“ Nouvelles espèces d'Hyménoptères des genres *Halictus*, *Prosopis*, *Allodape* et *Nomioides* rapportées par M. Fea de la Birmanie.”—Annuario del Museo Zoologico della R. Università di Napoli, Naples.

PRINTED BY
GOVERNMENT OF INDIA PRESS
CALCUTTA

Arrangements are being made to figure, in a final paper dealing with the Zoological Results of the Abor Expedition as a whole, a number of the new species of insects described without figures in this Part.