

RECORDS

of the

INDIAN MUSEUM

(A Journal of Indian Zoology)

Vol. 51, Part 4

December, 1953

	<i>Page</i>
On a new evolutionary phenomenon : the sharp increase of intraspecific variation in minimum populations, as evidenced by the Desert Locust. <i>M. L. Roonwal</i>	481
Siluroid Fishes of India, Burma and Ceylon XIV. Fishes of the genus <i>Mystus scopoli</i>. <i>K. C. Jayaram</i>	527
On some variations in the nematode genera <i>Porrocaecum</i> and <i>Echinocephalus</i> <i>T. D. Soota</i>	559

Edited by the Director, Zoological Survey of India.

ON A NEW EVOLUTIONARY PHENOMENON: THE SHARP INCREASE OF INTRASPECIFIC VARIATION IN MINIMUM POPULATIONS, AS EVIDENCED BY THE DESERT LOCUST.

By M. L. ROONWAL, M.Sc., Ph.D. (CANTAB.), F.N.I.

Forest Entomologist, Forest Research Institute, Dehra Dun.

(PLATES XV—XVIII.)

(With 4 Plates, 15 Text-figures and 5 Tables.)

CONTENTS.		PAGE.
I.—Introduction	. . .	481
(a) General	. . .	481
(b) Acknowledgements	. . .	482
II.—Some Data on the Desert Locust	483
(a) Geographical distribution, etc., of Desert Locust	. . .	483
(b) Population-density and amplitude and periodicity of population-flux in Desert Locust		486
III.—Discussion on Known Types of Evolutionary Phenomena or “ Effects ”		490
(a) Some general considerations	490
(b) Known types of evolutionary “ Effects ”	. . .	492
IV.—A New Evolutionary Phenomenon or “ Effect ”	. . .	498
(a) General remarks		498
(b) Increased variability in <i>solitaria</i> phase of Desert Locust		499
(c) Discussion	508
V.—Evolution of Various Types in Desert Locust		516
VI.—Summary	516
VII.—References	517
VIII.—Explanation of Plates		525

I.—INTRODUCTION.

(a) GENERAL.

The importance of the abundance or otherwise of the population of a species playing a role in the production of variations has been realized since the days of Darwin (1876) who stated that in plants the more abundant species show greater variation, that is to say, a larger number of varieties, than the rarer species. Adlerz (1902), however, was the first author to bring forth clearly an example of the effect of population fluctuation on the production of variations in a species. He observed that in the Swedish butterfly, *Polyommatus virgaureae* L., which shows periodic eruptions in numbers, morphological variation

is greatly increased during years of mass-abundance than in normal years. Since then, several other such examples have come to light. The theoretical study of the statistical effects of population-size alone, other factors being constant, has been made in recent years by Hagedoorn & Hagedoorn (1921), Dubinin (1931), Wright (1931-1946) and others. From these analyses it is clear that population-size itself is capable of influencing variation-intensity to a considerable extent.

As a result of these observations and analyses, there have been revealed a variety of evolutionary phenomena (which, in the literature, have come to be termed as "Effects") concerning the effect of population-size on the intensity of the production of variations. Our knowledge of these phenomena is based in some cases on field observations alone, and in others on theoretical analyses alone. The following 4 types of phenomena are hitherto known to exist :—(i) The automatic maintenance (equilibrium) of genetic variability in spite of crossing (the Hardy Effect); (ii) the rapid random (automatic) decay of variation-intensity in the total absence of selection pressure (the Hagedoorns Effect); (iii) the increase of variation-intensity with high populations (the Adlerz Effect); and (iv) the increase of variation-intensity neither in very high nor in very low populations but in populations of intermediate sizes (the Wright-Dubinin Effect).

In the present paper I have described a New Phenomenon or Effect which is characterised by the fact of the *highest* variation-intensity in *minimum* populations and the *lowest* variation-intensity in *maximum* populations. The New Effect has been found from a study of variations in the Desert Locust, *Schistocerca gregaria* (Forsk.). This species is subject to periodic mass-outbreaks in comparatively short cycles of about 11 years. The outbreak years are characterised by intensive swarming in enormous numbers and extensive overflow-migrations. In the intervening years the species is confined to restricted permanent breeding areas; and the seasonal migrations are restricted to this area, there being no overflow-migrations. The swarming and the non-swarming individuals show marked differences in colour, morphology, physiology and behaviour (Uvarov, 1923) and were called the two "phases", *viz.*, phase *gregaria* and phase *solitaria*. The existence of increased variability in the *solitaria* phase was shown by Roonwal (1936 *et seq.*). In a preliminary account (Roonwal, 1947*d*) I had regarded this increased variation-intensity as an example of the Wright-Dubinin Effect, but later work has shown that it differs markedly from that Effect and is an altogether new phenomenon not hitherto described. I have also critically reviewed the available information on the various phenomena mentioned above, so that the new phenomenon presented here may be viewed against a suitable background.

(b) ACKNOWLEDGEMENTS.

It is a pleasure to express my indebtedness to the following gentlemen for the assistance which they kindly rendered in various ways: To Shri Y. Ramachandra Rao, lately Deputy Director, Bureau of Plant Protection and Quarantines, New Delhi, and formerly Locust Research

Entomologist to the Indian Council of Agricultural Research, for supplying valuable information regarding eye-stripe polymorphism in the Desert Locust from his field records and collections ; to Dr. K. R. Nair, Statistician at the Forest Research Institute, Dehra Dun, for assistance in the calculations and for discussing some problems of correlations ; to Brigadier Sir Oliver Wheeler, lately the Surveyor-General of India, New Delhi, for advice regarding suitable map-projections to use and regarding the method of computing land-areas ; and, finally, to Dr. B. P. Uvarov, Director, Anti-Locust Research Centre, London, for providing some information on the distribution of the Desert Locust.

II.—SOME DATA ON THE DESERT LOCUST.

(a) GEOGRAPHICAL DISTRIBUTION, ETC., OF DESERT LOCUST.

(Pl. XV.)

(i) *Geographical distribution.*

The geographical distribution of the Desert Locust differs materially in the swarming or *gregaria* phase and the non-swarming or *solitaria* phase. The following are the more important authorities who have contributed records on which our present knowledge of the distribution is based :—Drouet (1861), Walker (1870), Darwin (1876), Uvarov (1923, 1928, 1932), Chopard (1932), Faure (1932), Vayssi re (1933), Predetchensky (1935), Rao (1937, 1945), Waloff (1946*a*, *b*), Roonwal (1950), ‘F.A.O.’ (1952) and Fortesque-Foulkes (1953).

Phase gregaria.—In the swarming or *gregaria* phase, this locust extends over 3 continents, *viz.*, Europe, Africa and Asia. Each of these continents is invaded to the extent indicated below : ASIA : In Asia it covers the whole of the Western, Middle and S. E. Asia, east (in India) up to Assam and south up to about the level of Madras (Rao, 1945 ; and Roonwal, 1950). Walker (1870) has recorded some specimens from “Ceylon” as being present in the collection of the British Museum (Natural History), London. But Dr. Uvarov (*in litt.*, 10th November, 1947) has informed me that these specimens, which are still present, were almost certainly mislabelled “Ceylon” and were most probably from the mainland of India. There are several records from Nepal and Kashmir. Swarms are recorded north from Afghanistan and Transcaspia (southern U. S. S. R.) to almost the level of the northern tip of the Caspian Sea. The whole of the Western and Middle Asian region (Iran, Iraq, Arabia, Syria, Israel and Turkey) is subject to its invasions. EUROPE : Stray specimens in Southern Europe, evidently migrants from North Africa, have been recorded from Corfu, Noxos, Portugal, and southern England, *vide* Lucas (1920), Uvarov (1923, 1928) and Waloff (1946*b*). AFRICA : The whole of Africa, with the probable exception of the Central Sahara and the tropical rain forests of the Belgian Congo, is subject to its invasions. Swarms are common in Lybia, Algeria, Morocco, French West Africa, Egypt, the Sudan, Eritrea, Ethiopia, Somaliland, Kenya, Uganda, Tanganyika and French Equatorial Africa. Records from South Africa are scanty but specimens in the

gregaria phase have been recorded as far south as the Cape Province. There are interesting records from the Atlantic islands off the western coast of Africa, *viz.*, from Madeira, Canary, Azores, Cape Verde and Ascension, *vide* Drouet (1861), Walker (1870), Darwin (1876), Chopard (1932), Vayssiére (1933) and Waloff (1946*b*). Scudder (1883) has recorded a swarm that came on board a ship in latitude 25° 28' N. and longitude 41° 31' W., *i.e.*, about midway between Africa and America, with the nearest point of land about 1,200 miles off.

Phase solitaria.—The distribution of the *solitaria* or non-swarmling phase is restricted to Northern and South (?) Africa and Western Asia up to and including the Indian desert region of Baluchistan, Sind, the Punjab, Rajasthan and Kutch. Thence, the region extends to Iran, Iraq, Arabia, etc., right up to the western coast of Africa, including the Red Sea coast, Eritrea, Somaliland, Egypt, the Sudan, Morocco, French West Africa, Senegal, etc. *Solitaria* phase specimens have also been recorded from South Africa. For further details *vide* Uvarov (1923, 1928, 1932), Faure (1932), Maxwell-Darling (1934-1937), Vayssiére (1936), Rao (1937, 1945) and Roonwal (1946*b*, 1950).

As already pointed out by me (Roonwal, 1949*d*, pp. 167-168), Bodenheimer's (1937) claim of the "Sudano-Deccanian" region (extending from the Sudan in North Africa to the Deccan in peninsular India) as the permanent home of the Desert Locust, is unwarranted in so far as the Deccanian portion is concerned. We know definitely that the permanent home in India is confined to the Western desert regions of Baluchistan, Sind, the Punjab, Rajasthan and Kutch. The locusts reach the Deccan only in very exceptional swarming years and never breed there. The other equally unwarranted claims of Bodenheimer are discussed below.

(ii) *Surface-area of distribution.*

It will be seen from the foregoing account that the area occupied by the Desert Locust in its swarming phase is much more extensive than that occupied in the non-swarmling phase, and a quantitative comparison of the two is of considerable interest.

The following method was followed for determining the total area of distribution:—An outline map of the world (Pl. XV) was drawn according to the Re-centred Sinusoidal Projection (Bartholomew) developed from the Sanson-Flamsteed's Equal Area Projection. This projection is area-true for all lands, while distortion is reduced to a minimum. The natural scale on this map was 1 : 100 000,000, which, during reproduction in Pl. XV, was considerably reduced. A scale (in cm.) which was *natural size in the original map* is also given for calculating the degree of reduction from the original during reproduction.

The choice of the map projection to be used was based on the advice of Brigadier Sir Oliver Wheeler, lately the Surveyor-General of India, New Delhi, who wrote (*in litt.*, 12th April, 1945) as follows :—

- “ 1. The simplest way is to use a map that is drawn on one of the equal-area projections and, as you suggest, count the squares on a piece of squared paper laid over it. This is far quicker than planimeter for an approximation. Planimeter areas have to be cross-checked against blunders and are troublesome. Some equal-area projections commonly used in Atlases are: Bonne’s (Modified conical equal-area); Sanson’s (Sinusoidal equal-area); Lambert’s Third (Cylindrical equal-area); Lambert’s Sixth (Zenithal equal area).”
- “ 2. Mercator’s projection should on no account be used *except* by sub-dividing your area into East/West belts, and ascertaining the average scale of each belt. This is easily possible on some world maps by direct reading of a varying (with latitude) scale; in the absence of such a scale it is necessary to compute the average scale of each belt.”
- “ 3. In the absence of a map on a suitable projection or knowledge of what the projection is; a method that can be applied to *any* projection is to draw (or superimpose) “quadrilaterals” bounded by meridians and parallels and add up the square miles enclosed within each quadrilateral.”

“ This method is probably more meticulous and troublesome than is necessary for your purposes but if you have to resort to it, we have published tables of the area of $\frac{1}{4}^{\circ}$ quadrilaterals that are good between 44° N. and 44° S. They do not go to higher latitudes than 44° ”

Kellaway (1949, p. 118) has also stated that “ for a map of Africa *Sanson-Flamsteed’s Projection* is satisfactory for most general purposes ” The bulk of my areas lie in Africa.

On this map the distribution of the Desert Locust in the *gregaria* and the *solitaria* phases was carefully plotted and outlined. A transparent squared paper, with $\frac{1}{2}$ centimetre squares drawn in black ink, was then superimposed on the distribution map, and the number of squares in each of the two areas, *viz.*, the *gregaria* and the *solitaria* distribution-areas, were counted off. Due allowance was made for fractions of squares which were suitably added to make up approximations of full squares. In counting the squares only the land-area was, as a rule, taken into account, and not the entire area encircled in the outlines. Thus, the oceanic areas of the Atlantic and the Indian Oceans were omitted. But smaller waters such as the Persian Gulf, the Red Sea, the Mediterranean Sea and the Caspian Sea (which the Desert Locust normally has no great difficulty in crossing, and, as a rule, does cross almost regularly in its periodic migrations) were included in counting. On the scale on which the original map was drawn, 13 linear half-centimetres equalled 4,000 miles. In other words, 13×13 or 169 half-centi-

metre squares covered an area equal to $4,000 \times 4,000$ or 16,000,000 square miles. Therefore, one half-centimetre square represents about 94674.6 square miles. On this basis the following approximate land-areas were obtained :—

(a) *Phase gregaria*.

	Square miles.
Europe	[142,012
Africa	11,360,952
Asia	4,970,417
	<hr/>
TOTAL or say	16,473,381 16,500,000
	<hr/>

(b) *Phase solitaria*.

	Square miles.
North Africa and Asia	8,662,726
South Africa	805,734
	<hr/>
TOTAL or say	9,468,460 9,500,000
	<hr/>

It will be seen that the *gregaria* area is about 1.7 times or nearly $1\frac{2}{3}$ rds the *solitaria* area.

(b) POPULATION-DENSITY AND AMPLITUDE AND PERIODICITY OF POPULATION-FLUX IN DESERT LOCUST.

(i) *Relative numbers*.

Having discussed the area of distribution, it is necessary now to obtain some idea on the density of population—a factor which is partly, but not wholly, interlinked with the area of distribution. At the same time, population-density is a factor of great importance in the mechanisms of speciation. Abundant evidence has now been accumulated to show that populations in no species are constant; they are, on the other hand, in a continuous state of flux. While this broad fact may be taken as well established, it must be said that a more or less precise knowledge on the nature of the flux is limited to but a few species, and here too many important aspects of the flux need fuller investigation. This knowledge mainly concerns 4 aspects, *viz.*, (i) the absolute numbers; (ii) the periodicity and amplitude of flux; (iii) the coincidental changes in the morphology, physiology and behaviour of the species undergoing population-flux; and, finally, (iv) the effect of all these factors on evolutionary processes.

Fortunately, the Desert Locust has proved to be an almost ideal material for the problems stated above. What is more, our knowledge of its population-densities, of the periodicity and amplitude of population-flux, and of the coincidental changes in its morphology, physiology and behaviour, is perhaps more complete than for any other species. There are also other advantages, namely, that (i) the amplitude of flux is very great; and (ii) the periodicity is not immoderately long, one full cycle usually lasting only about 11.1 years on an average (*vide* Roonwal 1950; and Table 1 below).

TABLE 1.—Swarming periods of the Desert Locust in India from 1860 onwards. (*Modified from Roonwal, 1950.*)

Swarming periods.	Non-swarming intervals.	Period for each full cycle.
I. 1860—66 (7 years) ..	—	—
— ..	2 yrs.	9 yrs.
II. 1869—73 (5 years)	—	—
— ..	2 yrs.	7 yrs.
III. 1876—81 (6 years)	—	—
—	7 yrs.	13 yrs.
IV. 1889—98 (10 years)	—	—
—	1 yr.	11 yrs.
V. 1900—07 (8 years)	—	—
—	4 yrs.	12 yrs.
VI. 1912—19 (8 years)	—	—
—	6 yrs.	14 yrs.
VII. 1926—31 (6 years)	—	—
—	8 yrs.	14 yrs.
VIII. 1940—46 (7 years)	—	—
— ..	2 yrs.	9 yrs.
IX. 1949—	—	—
AVERAGE : 7.1 years	4 years	11.1 years

Precise knowledge of population-densities and of the amplitude and periodicity of population-flux under natural conditions in the field has been obtained in the last 15 years, largely as a result of long and painstaking investigations by Indian entomologists, particularly Y. Ramachandra Rao and his co-workers, working in the desert regions of Baluchistan, Sind, the Punjab and Rajasthan.

Regarding periodicity, it may be stated that since 1860. there have been nine swarming cycles in India (Table 1), with the swarming periods varying from 5-10 years (average 7.1 years), the non-swarming intervals varying from 1-8 years (average 4 years), and a complete cycle varying from 7-14 years (average 11.1 years).

As for population-densities, it has now been established that in the *solitaria* or non-swarming years densities of adults in the permanent breeding grounds, such as Rajasthan, Sind and Mekran (Baluchistan), never exceed about 1,000-2,000 per square mile, and are usually much less. Until recently, we had no knowledge of the critical densities in the year which inaugurates a new swarming cycle, but the recent work of Misra, Nair & Roonwal (1952) has shown that this figure is not less than about 18,000 adult individuals per square mile; the increased density inaugurated the recent or 9th cycle which began in 1949 and is now (1953) continuing with vigour. It is of interest to mention here that in the middle of the 1932-1939 non-swarming period of low population, there occurred in July 1935 (as reported by Rao, 1936, and further discussed by Roonwal, 1949*d*) a sudden increase of numbers and the population increased from the *solitaria* figure of about 460 per sq. mile to 13,000 per sq. mile in Mekran and 40,000 per sq. mile in the Sind-Rajputana desert. This 'incursion', however, fizzled out and no swarming cycle set in; by December of the same year the new brood showed a population of only about 700 per sq. mile. No precise figures are available of the population-density in swarming years but we may tentatively take it as not less than about 30,000 per sq. mile; usually, it is certainly much higher. The position as regards population-density may be summed up as follows in a simplified way:—

- | | |
|---|----------------------------------|
| 1. In <i>solitaria</i> or non-swarming period | 1,000 per sq. mile. |
| 2. In the inaugurating year of a new swarming cycle | 20,000 per sq. mile. |
| 3. In the swarming period | .. 30,000 and over per sq. mile. |

For the hopper stages, I have observed in the swarming years 1940-1946 in the breeding fields in the Rajasthan desert, hopper masses giving maximum densities of about 25,000 per square metre in the freshly hatched first stage hoppers and about 1,050 per square metre in the fifth stage. Singh & Bhatia (1952) have recently recorded somewhat similar maximum densities, *viz.*, 2,840 hoppers per square foot in the first stage and 94 in the fifth stage.

(ii) *Absolute numbers.*


(Text-fig. 1 & Table 2.)

We may now discuss the significance of these population-densities on the absolute numbers of adults present at any one time. *Theoretically,*

if we assume that the adults are uniformly distributed in the densities stated above, the total absolute numbers in the area of its entire range of distribution (*vide supra*) would be as follows (Table 2) :—

In the *gregaria* phase period : 495,000,000,000 adults.

In the *solitaria* phase period : 9,500,000,000 adults.



TEXT-FIG. 1.—*Schistocerca gregaria*. Diagrammatic representation of the nature of distribution of population in the *gregaria* (swarming) and the *solitaria* (non-swarming) phase periods. The total surface land-areas occupied during the two periods are shown in their proportionate dimensions; the *solitaria* area is about 9,500,000 sq. miles and the *gregaria* area about 16,500,000 sq. miles, or 1.7 times larger (*see text* for fuller details). Distribution is more uniform in the *solitaria* period, and comparatively more discontinuous in the *gregaria* period. However, populations in the *gregaria* period are much more concentrated and dense (*ca.* 30,000 or more per sq. mile) than in the *solitaria* period (*ca.* 1,000 or less per sq. mile).

(a) Phase *gregaria* period. (b) Phase *solitaria* period.

TABLE 2.—Theoretical (calculated) absolute numbers of Desert Locust adults present at any given moment in the entire land-area of its distribution, on the assumption of uniform distribution in the given densities.

Phase.	Total land-area of known distribution (in square miles).	Known approximate densities (per square mile).	Calculated absolute numbers.
(A) Ph. <i>gregaria</i>	16,500,000	30,000	495,000,000,000
(B) Ph. <i>solitaria</i>	9,500,000	1,000	9,500,000,000
Times increase of A over B	1.7	30	52

Actually, however, these numbers are misleading, since neither in the swarming nor in the non-swarming years is the distribution of population uniform. The distribution is always in 'pockets' or 'islands' of varying shapes and dimensions, leaving considerable tracts entirely void. This feature of discontinuity is especially pronounced in the *gregaria* phase, the distribution in the *solitaria* phase being

comparatively more uniform (Text-fig. 1). The dimensions of these islands of distribution vary from one to several square miles. They are generally larger in the *solitaria* phase and smaller (more or less confined to individual swarms and the areas where they oviposit) in the *gregaria* phase. This situation results in a certain contradiction, namely, that in the *gregaria* phase the individual 'islands of distribution' are generally smaller than in the *solitaria* phase, although the total land-area of distribution is much larger (Pl. XV).

The numbers with which, then, we are concerned as that which is dynamically important for interactions in the processes of speciation or production of variations are not the absolute theoretical numbers but the actual densities in small islands of distribution. As already stated, these densities are the lowest (*ca.* 1,000 per sq. mile) in the *solitaria* phase and the highest (*ca.* 30,000 at the least, or nearly 30 times or usually more) in the *gregaria* phase. In discussing their respective effects on the production of variations we shall proceed on the basis of these rough proportions.

III.—DISCUSSION OF KNOWN TYPES OF EVOLUTIONARY PHENOMENA OR "EFFECTS"

(a) SOME GENERAL CONSIDERATIONS.

No two individuals of a species are alike. It is this *existence* of variability, howsoever caused, which is the basis of speciation. Since variation first arises in individuals and then spreads to the group, we may say with justification that the progress of speciation ultimately depends upon the behaviour of variability in natural populations. This behaviour, in its turn, depends upon a variety of factors such as genetic differences (mutation rate), geographical and ecological distance (isolation; migration or cross-breeding rate between different groups, whatever the cause), elimination of variant individuals by 'natural' and other forms of selection (selection pressure), and, finally, population structure. Briefly, the following are the chief factors at work in speciation:—

1. Mutation pressure.
2. Selection pressure.
3. Migration pressure.
4. Population pressure :
 - (a) Population size.
 - (b) Random drift or genetical-automatic processes.

Speciation theoretically implies that a genetically homogeneous population-group, that is, a group in which the balanced gene-frequency in respect of a particular gene *AA* has reached 100 per cent. at level C, first achieves heterogeneity by the introduction of a new allelomorph *aa*, either by crossing or by mutation, and ultimately achieves a new genetic homogeneity at level D. In actual practice, however, in the first place not one but perhaps scores of genes are involved in the origin

of a new species. Secondly, complete homogeneity is seldom achieved, except in pure line (clones) and perhaps in extremely small populations of bisexually reproducing forms; there occurs, instead, 'a balanced heterogeneity' which jumps from one peak of stability to another.

The factors leading to genic homogeneity (less variation) or heterogeneity (more variation) are briefly given in Table 3. The state of variability or gene frequencies in a species or population-group is in a continual state of vibration about the equilibrium point in a distribution curve the form of which depends upon the relations between population number and the various pressures indicated above.

TABLE 3.—Factors leading to genic homogeneity or heterogeneity.

FACTORS FOR GENIC HOMOGENEITY	FACTORS FOR GENIC HETEROGENEITY
1. Gene duplication.	1. Gene mutation (u, v).
2. Gene aggregation.	2. Random division of aggregate.
3. Mitosis.	3. Chromosome aberration.
4. Conjugation.	4. Reduction (meiosis).
5. Linkage.	5. Hybridization (m).
6. Population size. (Varying effects reported.)	6. Population size. (Varying effects reported).
7. Environmental or selection pressure (s).	7. Pressure of local environments of subgroups (s_1).
8. Cross-breeding (migration pressure).	8. Individual adaptability.
9. Individual adaptability.	

The relation subsisting between population number (*i.e.*, abundance) on the one hand, and variability on the other, had been realised since a long time on observational grounds (see Darwin, 1876; Adlerz, 1902). Hagedoorn & Hagedoorn (1921) have pointed out the importance of fluctuation in numbers in causing specific uniformity; they, however, believed that the total number of individuals in a species is constant. Elton (1924) has well emphasised the existence of periodic fluctuations in various groups of animals, especially mammals, and has expressed the view that they must be caused by periodic climatic changes acting over wide areas. He further believed that population fluctuation causes uniformity because the whole population re-arises from a few surviving individuals after a period of low numbers; the genotypic constitution tends to become pure. He wrote (p. 159): "In fact there is in many animals a mechanism by which variations in the population are alternately acquired and weeded out or kept as indifferent characters which in turn may become useful at some crisis in the history of the animal at a later date" According to Elton, fluctuation explains many things in evolution, such as: (i) how, by a temporary cessation of

the struggle for existence, new and indifferent (non-adaptive) mutations can get a footing and spread ; (ii) how some characters are selected periodically ; (iii) how species remain on the whole uniform ; and (iv) how adaptive fineness may arise without reference to selection and crossing. " There is ", he wrote (p. 160), " so to speak, an entrance examination by natural selection, which weeds out the worst candidates, but the final selection is by lot "

Attempts at theoretical interpretations on mathematical grounds are of recent origin and, for certain types of phenomena, are largely due to the analyses of Chetverikov (1926), Fisher (1930), Haldane (1932) ; Wright (1931-1946), Dubinin (1931), Dubinin & Romaschoff (1932) and Kolmagorov (1935). Although these analyses are theoretical and refer to one or two types of phenomena only, they have been valuable in providing clarifications which could not be achieved by other means. It now remains to apply similar analytical techniques to the other types of phenomena discovered in the field by actual biological observations on wild populations. (For a somewhat fuller discussion of the problem, see Roonwal, 1949b.)

(b) KNOWN TYPES OF EVOLUTIONARY " EFFECTS "

A survey of the published literature shows that four different types of evolutionary phenomena have been recorded in regard to the intensity of the production of variation in relation to population-flux ; and a fifth one is presented below in this paper. One of these phenomenon has been conveniently termed in recent literature as the " Wright-Dubinin Effect " and, following this practice, I have termed the other ones also by a similar nomenclature, associating them with the names of their discoverers. These phenomena are as follows :—

1. The Hardy Effect.—Automatic maintenance of genetic variability in spite of crossing.
2. The Hagedoorns Effect.—Rapid random (automatic) decay of variation-intensity in the total absence of selection pressure.
3. The Adlerz Effect.—Increase of variation-intensity with increased abundance (high populations).
4. The Wright-Dubinin Effect.—Increase of variation-intensity in intermediate populations, *i.e.*, populations which are neither too large nor too small.
5. The New Effect (presented here).—Increase of variation-intensity in *minimum* populations.

These effects may now be discussed in some detail (also see Roonwal, 1949b).

1. *The Hardy Effect*

Hardy (1908) has demonstrated that genetic variability, once gained by a population, is automatically maintained and is not reduced by

crossing. If AA and aa are the new allelomorphs, and their populations are mixed in the proportions q and $1-q$, the second and subsequent generations will show the following structure of gene-frequencies (Hardy's Equilibrium Formula) :—

$$q^2 AA : 2q(1-q) Aa : (1-q)^2 aa,$$

assuming that the numbers are fairly large so as to ensure random mating, that the sexes are evenly distributed among the three varieties (dominant AA , heterozygous Aa and recessive aa), and that all are equally fertile. Proportions will change up to the second generation, but thereafter they will strike an equilibrium and will have no tendency to either increase or decrease. Hardy concluded thus : "In a word there is not the slightest foundation for the idea that a dominant character should show a tendency to spread over a whole population, or that a recessive should tend to die out "

2. *The Hagedoorns Effect*

Hagedoorn & Hagedoorn (1921) visualised the theoretical possibility of a rapid, random (automatic) decay of variability in a stable population in the total absence of mutation and selection pressures. Actual instances are not known. Fisher (1922) has criticised the Hagedoorns Effect and has remarked that such random decay of variance is so extremely slow as to be quite unimportant. He has further pointed out that if k is the rate of decay and n the number of individuals breeding at random, the rate of extinction is $k = \frac{1}{2n}$ and it would require $2n^*$ generations to reduce its variance in the ratio 1 to e^\dagger or $1.4n$ generations to halve it. When the number of interbreeding individuals in a species is large (say not less than 10,000), it will be seen that the rate of decay of variance is so slow as to be quite negligible. In a later paper Fisher (1930*b*, p. 204) has stated that " random survival, while of great importance in conditioning the fate of an individual mutant gene, is a totally unimportant factor in the balance of forces by which the actual variability of species is determined " He has further stated that this is because, in the total absence of mutation-pressure and selection-pressure, the quantity of variation of an interbreeding group would decrease by random survival at a rate such that the time of relaxation is $2n$ generations (n = number of breeding individuals in each generation). This number is usually very large. The time needed to halve the variance by random extinction of genes in the total absence of mutations is 1.4 times the number of potential parents in each generation (p. 219) ; and the loss of variance due to this cause is very trifling. Thus, the steady random decay of variation as visualized by Hagedoorn & Hagedoorn (1921) will not occur and, in the absence of selection, variability will remain constant but will now evidently depend on the size of the population. If the number of breeding individuals n is small, then the Hagedoorns Effect may become significant.

* Fisher (1930*b*) later on corrected his original (1922) figure of the " time of relaxation " from $4n$ to $2n$; thus, 2.8 also becomes 1.4.

† Evidently e = gene-extinction.

Hagedoorn & Hagedoorn (1921) have also pointed out the importance of population fluctuation for reducing variations and producing homogeneity. They, however, believed in the constancy of the number of individuals of a species from year to year. By population fluctuation they merely implied migration to new habitats or to the effect of unusual climatic and biological changes, and so on. They also suggested that in spite of the constancy of the total number in a species (which we now know is not the case), only a certain number breed each year out of the surviving population (here again, no evidence is brought forth). This, if true, will have the same effect as fluctuation of population and may, in special cases, be significant. It may be stated that Hagedoorn & Hagedoorn were evidently not aware of the observations of Adlerz (1902) on the effect of population-flux on variation.

3. *The Adlerz Effect*

Darwin (1876, Chapter 2, pp. 42-43) believed that the more abundant species show greater variation (a large number of varieties) than the rarer ones, but he did not publish any detailed evidence for this belief except saying that he had examined, more or less statistically, a fairly large number of cases among plants. He wrote :

“ Alphonse de Candolle and others have shown that plants which have very wide ranges generally present varieties ; and this might have been expected, as they are exposed to diverse physical conditions, and as they come into competition (which, as we shall hereafter see, is an equally or more important circumstance) with different sets of organic beings. But my tables further show that, in any limited country, the species which are the most common, that is abound most in individuals, and the species which are most widely diffused within their own country (and this is a different consideration from wide range, and to a certain extent from commonness), oftenest give rise to varieties sufficiently well-marked to have been recorded in botanical works. Hence it is the most flourishing, or, as they may be called, the dominant species,—those which range widely, are the most diffused in their own country, and are the most numerous in individuals,—which oftenest produce well-marked varieties, or, as I consider them, incipient species. And this, perhaps, might have been anticipated ; for, as varieties, in order to become in any degree permanent, necessarily have to struggle with the other inhabitants of the country, the species which are already dominant will be the most likely to yield off-spring, which, though in some slight degree modified, still inherit those advantages that enable their parents to become dominant over their compatriots. In these remarks on predominance, it should be understood that reference is made only to the forms which come into competition with each other, and more especially to the members of the same genus or class having nearly similar habits of life. With respect to the number of individuals or commonness of species, the comparison of course relates only to the members of the same group. One of the higher plants may be said to be dominant if it be numerous in individuals and more widely diffused than the other plants of the same country, which live under nearly the same conditions. A plant of this kind is not the less dominant because some conferva in-

habiting the water or some parasitic fungus is infinitely more numerous in individuals, and more widely diffused. But if the conferva or parasitic fungus exceeds its allies in the above respects, it will then be dominant within its own class."

Adlerz (1902) was the first authority to demonstrate clearly the existence of increased variation with the increase of abundance in a species whose population showed periodic fluctuations. His observations, however, appear to have been overlooked. He observed this phenomenon in the Swedish butterfly, *Polyommatus virgaureae* L.* In July 1896 this species greatly increased in numbers in Sweden, and at the same time showed a variety in females which had nowhere been known before. This variety was characterised by the presence of pale blue spots inside the golden-red band on the upper side of the hindwings. The number and clearness of the spots varied greatly. The greater number was 5, and from this to the chief form of the species (with no spots) there were all grades of intermediates. During the following years the numbers declined, and the absolute as well as relative proportion of the said variety became rarer than during the period of mass-increase. In the summer of 1901 again there was mass-increase in the numbers of this species, and again the said variety became more common—more than half the number of captured specimens showed the characters of the new variety.

A similar phenomenon has been subsequently reported by several workers in other species of animals. Thus, Fisher & Ford (1926-1928) found that in several species of Lepidoptera the more abundant species show a greater degree of variation (in depth of pigmentation) than the rarer ones. They also showed that this situation is to be expected even on theoretical (mathematical) grounds, assuming that Mendelian inheritance alone occurs. A similar theoretical conclusion was arrived at earlier by Fisher (1922), *vide infra*.

Fisher (1937a) reiterated the same position after statistically studying the egg-size of some British birds, although he was at the same time cautious. He wrote (p. 3): "The theoretical reasons for anticipating a higher variability associated with a numerically larger population are recondite and by no means fully understood. They depend on the balance of selective and mutational agencies, by which the genetic variability of natural populations must be determined. It is because we cannot easily predict this *a priori* that direct empirical knowledge is of importance. In respect of variations conditioned environmentally, there is no reason to suppose that these would be greater or less in a dense than in a sparse population. The same is true of that fraction of the variability which is caused by deleterious mutations. To these all species appear to be exposed." He further wrote (pp. 3-4): "If species of different populations were supplied at the same absolute rate with different neutral mutations, each occurring with extreme rarity, then calculation shows that the number of factors maintained in the heritable variance of the species would be nearly proportional to the logarithm of the population." And again (p. 23): "The excess variability appears to be strongly concentrated among the few most numerous

*Evidently the same as *Chrysophanus virgaureae* Linn. (family *Lycaenidae*).

species of each group, suggesting that evolutionary progress is much more rapid among these than among other species. This important indication may, however, be due to our comparative ignorance of the real relative numbers of the population of the less numerous species.”

Ford & Ford (1930) studied the variation in a small colony (of 6-7 miles radius) of the butterfly *Melitaea aurina* Rott. in Great Britain for a period of 49 years (1881-1930). During this period the species fluctuated in numbers as follows: Abundant 1881-1897; less common 1898-1906; uncommon 1907-1912; very rare 1912-1920; increase 1920-1924; abundant 1924-1930. The degree of parasitism was largely responsible for the fluctuation of population. Variation was small in the first and second periods of abundance but very great while the numbers were increasing rapidly. This variation concerned size, shape and colour; also, a high proportion of individuals were deformed in various ways. When the rapid increase of population ceased, the undesirable elements (deformities) practically disappeared, and the species settled down to a comparatively uniform type which, however, was distinct from that prevailing during the first period of abundance. Ford & Ford stated that the lowering of selection-intensity resulted in increased variation and also in increased numbers. In this example two points are worthy of note. First, that the greatest variation occurs while the population is rapidly increasing. Secondly, that the high variation-intensity ceased and was replaced by comparative uniformity when the rapid increase of population stopped and the population presumably settled down at a high but more or less steady level. The cause of the latter decline of variation is not clear.


On theoretical grounds, Fisher (1922) stated that increased abundance is accompanied by increased variation. He wrote (p. 324): “In all cases it is worth noting that the rate of mutation required varies as the variance of the species, but diminishes as the number of individuals is increased. Thus a numerous species, with the same frequency of mutation, will maintain a higher variability than will a less numerous species: in connection with this fact we cannot fail to remember the dictum of Charles Darwin, that ‘wide ranging, much diffused and common species vary most’ (1, chap. ii).” Fisher also gave graphs of distribution of variance in a species under various theoretical conditions.

4. *The Wright-Dubin in Effect*

The important theoretical demonstration that variation (gene-frequencies) may fluctuate merely as a statistical consequence of the ‘effective’ population-size, all other pressures being constant, we owe largely to the analyses of Chetverikov (1926), Dubinin (1931), Dubinin & Romaschoff (1932), Wright (1931-1946) and Kolmogorov (1935), but more particularly of Dubinin and of Wright. Dubinin termed this process the ‘genetico-automatic process’, while Wright’s term for it was ‘random drift’ or merely ‘drift’. They concluded that speciation occurs most rapidly neither in very large nor in very small populations but in populations of certain intermediate sizes. Great reduction of population leads to fixation (of non-adapted genes) and loss of genes;

great increase only to slow change. The most rapid change occurs in a large population which is subdivided into a number of partially isolated groups between which occasional crossing takes place.

In discussing evolution in populations in approximate equilibrium, Wright (1935, p. 257) has considered the "evolutionary processes in populations in which the selective values of different grades of a character depend on the (squares) deviations of the latter, from an optimum. It is assumed that the effects of different genes on the character combine additively (no epistasis)". Having reached an optimum (in selective variation), there would be no further change and new mutations would now be harmful to start with (*vide* also Fisher, 1930). But if genes have multiple effects, and harmonious adjustment of characters is arrived at, the possibility of numerous optimum "peaks" would arise (Text-fig. 2). Thus, all possible combinations may occur by a trial and error mechanism, and the locus of a population (regarding gene-variations) may be carried across a saddle from one peak to another and perhaps to a higher one. This view contrasts with that of steady progress under natural selection, as developed in its extreme form by Fisher (1930). Haldane (1932) occupies an intermediate position.



TEXT-FIG. 2.—Diagrammatic representation of the field of gene-combinations in two dimensions instead of many thousands. Dotted lines represent contours with respect to adaptiveness. (*After* Wright, 1932.)

Wright (1935) has shown that the effectiveness of this trial and error mechanism (random drift) of gene-frequencies is normally slow, but is enormously increased if the population is subdivided into many local groups which breed largely within themselves. The distribution of gene-frequencies under random sampling is here determined by the relation between the effective size of the local group and the cross breeding index. "At any time combinations might be reached by chance in particular local groups, with effects of the first order of importance, leading to expansion of such groups (intergroup selection)....." (p. 265).


These analyses, though important, are theoretical, and actual examples of the Wright-Dubin Effect are hitherto wanting. As pointed out by me earlier (Roonwal, 1949*b*, pp. 129-130), perhaps the extraordinary degree of speciation in birds and mammals among island forms,

particularly in such areas as the island-groups of the East Indies and Australasia, where a species has become divided into a large number of subspecies, one for almost every tiny island in its range of distribution, may be regarded as an example of this Effect. In my earlier accounts (Roonwal, 1947*d*; 1949*b*, p. 130) I had regarded the case of the increased variation in the *solitaria* phase of the Desert Locust, *Schistocerca gregaria*, also as an example of the Wright-Dubinin Effect, but it is now clear that it represents an altogether new and different kind of Effect, described below, which is characterised by increased variation-intensity in *minimum* populations.

IV.—A NEW EVOLUTIONARY PHENOMENON OR “EFFECT”.

(a) GENERAL REMARKS.

The data for the New Effect presented here are based on the writer's findings (Roonwal, 1936-1954) in the Desert Locust, *Schistocerca gregaria*



TEXT-FIG. 3.—*Schistocerca gregaria*. Diagrammatic representation of the periodic population fluctuation and its relationship to the eye-stripe number. (Modified from Roonwal, 1949*a*.) The number of peaks and valleys in the figure has no quantitative significance. The papillae in the region of the *congregans* population (left) represent the “outbreak centres”, while similar papillae in the *dissocians* population (right) represent isolated pockets of high population before the dispersal leading to low population (characteristic of the *solitaria* periods) occurs.

e, “eye-stripe figure” (see Text, p. 31, foot-note, for explanation).

(Forsk.) of greatly increased variation-intensity in the *solitaria* or non-swarming phase (where the population is the lowest, being of the order of about 1,000 - 2,000 adults, usually much less, per square mile at the maximum), as compared to the *gregaria* or swarming phase (where the population is the highest, being of the order of about 30,000 adults per square mile at the minimum).

The existence of two extreme phases, the *gregaria* and the *solitaria*, had already been established by Uvarov (1923, 1928) and later on confirmed by Faure (1932) and several other workers.

The *solitaria* and the *gregaria* phase periods alternate (*vide supra*, pp. 7-8; and Text-fig. 3). There is also a difference in geographical distribution in the two periods, as shown above. In the *solitaria* period the distribution is more restricted (being about 9,500,000 sq. miles of land-area) than in the *gregaria* period where it is about 16,500,000 sq. miles or nearly 1.7 times as great (Pl. XV).

The two phases also differ from each other in several other respects. Apart from differences of behaviour (the *gregaria* phase being the swarming phase, the *solitaria* the non-swarming one), there exist some colour differences (especially in the hoppers, see Pl. XVII) and differences in the shape and proportion of the body-parts and the related ratios, particularly the E/F ratios (concerning the elytron-length, E, and the hind-femur-length, F; see Text-figs. 4 and 5) and ratios connected with the head-width, C (*vide* Roonwal & Misra, 1952; Nair, 1952a). Other phase differences (Roonwal, 1946a) have also been established. Roonwal (1947a) has also shown a difference in the distribution of pigment in the cells of the ommatidia of the compound eyes (Pl. XVIII). Other phase differences are illustrated in Pls. XV-XVII and Text-figs. 4-9.

(b) INCREASED VARIABILITY IN *solitaria* PHASE OF DESERT LOCUST.

(i) *Some variable characters.*


Here, however, we are mainly concerned with the non-phasic type of variations which have been shown to exist in the *solitaria* phase by Roonwal (1936-1952). He showed (Roonwal, 1936) that 2 types of individuals occur in the *solitaria* phase, *viz.*, with 6 and 7 vertical eye-stripes. Later on, several other differences were shown to exist, such as variations in the number of antennal segments, the occurrence of 5- and 8-eye-striped individuals (Text-fig. 6), differences in sex-ratios correlated with the eye-stripe number, differences in the number of moults and the mode of development of the eye-stripes in the 5-, 6-, 7- and 8-striped forms (Roonwal, 1947a), and so on. The more important differences as regards increased variation between the *gregaria* and the *solitaria* phases on the one hand, and within the *solitaria* phase as on the other, are briefly mentioned below. For brevity, the words "*gregaria* phase" are abbreviated to "G." and "*solitaria* phase" to "S." :—

(A)—*Morphological Differences* (Pls. XVI-XVIII; and Text-figs. 4-9.)

1. On the whole, G. less variable than S.
2. G. has 6 eye-stripes only. S. has 5, 6, 7 and 8 eye-stripes, but mostly 6 and 7 (Roonwal, 1936, 1947a) (see Text fig. 6).
3. Eye-stripes in G. develop by one type only. In S. there are varying modes of development (Roonwal, 1947a).
4. The degree of sexual dimorphism (Text-fig. 7) as expressed by the Sexual Dimorphism Percentage or S. D. P. $[(\frac{\text{♀}}{\text{♂}} - 1) \times 100]$ in respect of the length of the body-parts is more marked in G. (S. D. P. for

elytron-length or E, 9.76 ; for hind-femur length or F, 8.72) than in S. Within the latter phase, S., the S. D. P. is higher in the 7-eye-striped than in 6-eye-striped forms, thus : 7-striped, for E 20.08, for F 18.33 ; 6-striped, for E 18.04, for F 15.63 (*vide* Roonwal, 1949c, 1952b ; Roonwal & Nag, 1950, 1951 ; and Roonwal & Misra, 1952).

5. The number of hind-tibial spines in adults show a somewhat greater range of variation in S. than in G. The outer spines vary from 8-12 in S. and 8-10 in G. The inner spines vary from 8-12 in S. and 10-12 in G.



TEXT-FIG. 4.—*Schistocerca gregaria*. Pronotum showing differences in the *solitaria* and the *gregaria* phases. Figures (a) and (c) in dorsal view ; (b) and (d) in lateral view ; (a), (b) Phase *solitaria* ; (c), (d) Phase *gregaria*.

l.m., lower margin of lateral lobe ; *l.s.*, lower pale horizontal stripe of lateral lobe ; *m.k.*, mid-dorsal keel ; *m.s.*, pale mid-dorsal stripe ; *u.s.*, upper pale horizontal stripe of lateral lobe.

6. The size of the body-parts, *e.g.*, length of elytron, length of hind-femur, width of head, length and width of pronotum, vary far more in the *solitaria* than in the *gregaria* phase, and show, in the former, various groupings which are correlated with the number of eye-stripes (Text-fig. 8)—*vide* Roonwal (1949c, 1952b), Roonwal & Nag (1950, 1951) and Roonwal & Misra (1952).


7. Similar increased variations in the *solitaria* phase, as compared to the *gregaria*, occur in respect of the biometrical ratios such as E/F or elytron-length/hind-femur-length (Text-fig. 5)—*vide* Roonwal (1949c) Roonwal & Nag (1950, 1951) and Roonwal & Misra (1952).

8. There are 26 antennal segments in the 6-eye-striped G. In S. the number varies from 25-30, as follows (Text-fig. 9): 5-eye-striped 25, 6-striped 26-27, 7-striped 28-29 and 8-striped 30 (*vide* Roonwal, 1952a).

(B)—Other Differences

1. Usually only 5 moults (excluding the “intermediate moult” in G. and 5-7 in S.)—*vide* Roonwal (1946a, 1947a).

2. Only one type of hopper (the yellow-black) and one type of adult (pink when immature and yellow when mature) in G. In S. two kinds of hoppers (the “green” type and the “fawn” type), and two kinds of adults (the “blue-grey” type and the “fawn” type)—*vide* Roonwal (1946b).



TEXT-FIG. 5.—*Schistocerca gregaria*. Graphs to show the variation in E/F ratios (elytron-length/hind-femur-length) in various types of individuals, *viz.*, phase *gregaria* 6-eye-striped, and phase *solitaria* 6-, 7- and 8-eye-striped. The mean E/F ratio is highest in the phase *gregaria* (6-striped) and lowest in the 8-striped phase *solitaria*. (*Adapted from* Roonwal, 1949a, with the addition of data for 8-eye-striped individuals.)


(ii) Periodic variation in eye-stripes during 25 years (1926-1950).

It will be seen from the foregoing account that whereas in the *gregaria* phase individuals there is a narrowing down of variation, in the *solitaria* phase individuals there is a remarkable flowering of variation. In other words, the *gregaria* phase is variation-stable, the *solitaria* phase variation-unstable. This difference is most strikingly demonstrated in the number of eye-stripes. Data for three cycles in India covering a period of 25 years (1926-1950) are now available* and we are thus in

* Data for the earlier years are also available from swarms and has already been made use of (*vide* Roonwal, 1949c); these data are, however, comparatively scanty. For a portion of the data for the period 1926-1950 I am indebted to Shri Y. Ramachandra Rao (*in litt.*, 25th April 1947).

a position to observe the periodic appearance and disappearance of this polymorphism simultaneously with the decrease and increase respectively of populations. The following is a brief description of this periodicity of polymorphism :—

1. *Swarming (gregaria) period 1926-31.*—Except during the last year (1931), all the individuals collected during the 1926-31 swarming cycle were 6-eye-striped. Of the several specimens collected at Pasni (Mekran, Baluchistan coast) in April and May 1931, a good many proved




TEXT-FIG. 6.—*Schistocerca gregaria*. Compound eyes of adults in various forms of individuals. (a) In *gregaria* or swarming phase, with 6 eye-stripes. (b)-(e) In *solitaria* or non-swarming phase, with 6, 7, 8 and 5 eye-stripes. In all figures the anterior side is on the right, the posterior one on the left.

d., dorsal spot of eye; *is.*, interstripe; *s1-s8*, first to eighth eye-stripes, counting from the posterior side.

to be of the *solitaria* phase (as judged by E/F ratios and other characters) and one specimen was found to have 7-eye-stripes. Collections made in October 1931 near Pasni also included one or two 7-eye-striped forms.

2. *Non-swarming (solitaria) period 1932-39.*—As mentioned above, the first indication of the appearance of 7-eye-striped individuals was in the last year of the swarming cycle of 1926-31. During the

solitaria years 1932-39, the relative proportions of the 6- and 7-eye-striped forms varied greatly (*vide* Rao, 1942, and *in litt.*, 1947 ; and Roonwal, 1949*d*)—from about 29-92 per cent in the case of 6-striped forms and, correspondingly, 8-71 per cent in the case of 7-striped ones. The greatest proportion of 6-striped forms was found during the July 1935 incursion of phase *gregaria*-like forms. Again, in 1937 there was a sharp rise in the proportion of 6-striped forms almost to the point of eclipsing the 7-striped ones, but towards the end of the year there



TEXT-FIG. 7.—*Schistocerca gregaria*. Bar diagrams showing the gradient in Sexual Dimorphism Percentage (S. D. P.) in biometric characters, *viz.*, E (length of elytron), F (length of hind-femur), C (width of head at the genal level) and P (length of pronotum at the dorsal keel), ranging from the 7-eye-striped *solitaria* to the 6-eye-striped *gregaria* populations under various conditions of population density and swarming. The polymorphism in the *solitaria* phase is evident, and gradually declines in the extreme *gregaria* phase. (After Roonwal & Misra, 1952.)


Ajmer sw. (6) 1950 (2nd yr.), population from a small swarm at Ajmer in June 1950, *i.e.*, in the 2nd year of the new swarming cycle in India ; *greg. (6)*, typical extreme phase *gregaria* population (from data of Roonwal, 1949*a* ; and Roonwal & Nag, 1950) ; *inc. sw. (6) 1949 (1st yr.)*, population from a locust concentration at Kakko (Bikaner, Rajasthan) in July 1949, *i.e.*, in the 1st year of a new swarming cycle in India (from data of Misra, 1950) ; *sol. (6)*, *sol. (7)*, typical 6-eye-striped and 7-eye-striped *solitaria* populations respectively (from data of Roonwal, 1949*a* ; and Roonwal & Nag, 1951).

was a sharp rise in the number of the latter form. In the peak *solitaria* year, 1936, the proportion of the 7-striped forms was comfortably higher than that of the 6-striped ones. Furthermore, 8-striped forms were first noted in the field in 1936 and most of them were collected during that year, only a few being seen in 1937 and 1938. The proportion of the 8-striped forms, as compared to that of the 6- and 7-striped ones, was always low, not more than about 1 out of every 20 to 50 locusts (Rao, *in litt.*) and the population in these surveys ranged from about 100-2,000 individuals per square mile. Very few 8-striped forms were

found in 1937, and only one in 1938. In 1939, no 8-striped form was found and the proportion of the 6- and 7-striped forms varied as 66-91 : 9-34.

As the 8-eye-striped form represents the extreme development of the *solitaria* period, the following details about the wild forms, kindly supplied by Shri Y. R. Rao (*in litt.*), are of interest :—

- (1) 18.ii.1936 : Jasurana (Jaisalmer Dist., Rajasthan) ; ♂ ; E/F 1.99.
- (2) 16.vi.1936 : Pholia (Jaisalmer Dist., Rajasthan) ; ♂ ; in *solitaria* phase.
- (3) 2.ix.1936. Mithra (Bahawalpur State, Punjab) ; ♀ ; E/F 1.97.
- (4) 13.ix.1936. Udaramsar (Bikaner Dist., Rajasthan) ; ♂ ; E/F 1.94.
- (5) 14.ix.1936. Sardarshahar (Bikaner Dist., Rajasthan) ; ♂ ; E/F 2.08.
- (6) 30.ix.1936. Naka Kharrari (Lasbela State, Baluchistan) ; ♀ ; E/F 1.91.
- (7) 28.x.1936. Bhagori (Lasbela State, Baluchistan) ; ♀ ; E/F 2.11.
- (8) —.ii.1937. Ambagh (Lasbela State, Baluchistan).
- (9) 3.x.1937. Girasar (Jaisalmer Dist., Rajasthan) ; ♀ ; E/F 2.00.
- (10) 25.vii.1938. Gurrani (near Pasni, Baluchistan coast) ; ♀ ; E/F 1.99.



TEXT-FIG. 8.—*Schistocerca gregaria*. Polymorphic variability in the size of certain body-parts, viz., elytron-length (E), hind-femur-length (F), head-width in genal level (C), and prenatal width at the keel (P), under various conditions of population density from extreme phase *solitaria* (low populations) to extreme phase *gregaria* (high populations). Each bar represents the mean size, in mm., of the body-part in females (higher level) and males (lower level of same bar) under defined population conditions (*see inset for explanation*). The difference between the values in the two sexes is represented by the blank rectangle at the top of each bar. Note the graduality in the fall of values, the male E being exceptional. The graduality in the decline of sexual differences with increasing population density is noticeable. For explanation of abbreviations, *see Text-fig. 7.* (After Roonwal & Misra, 1952.)

TABLE 4.—Showing E/F ratios (elytron length/hind-femur length) in the various types of individuals of *Schistocerca gregaria* (Forsk.). (Data for the 6- and 7-eye-striped forms are *after* Roonwal, 1949c ; and Roonwal & Nag, 1951.)

Abbreviations.—*greg.*, phase *gregaria* ; *sol.*, phase *solitaria*.

Phase, sex and number of eye-stripes	E/F ratio	
	Range	Mean, with Standard Error
<i>greg.</i> ♂♂ : (6-striped)	2.04—2.30	2.17±0.02
<i>sol.</i> ♂♂ : (6-striped)	1.88—2.23	2.05±0.01
<i>sol.</i> ♂♂ : (7-striped)	1.85—2.13	2.00±0.01
<i>sol.</i> ♂♂ : (8-striped)	1.94—2.08	2.003±0.41
<i>greg.</i> ♀♀ : (6-striped)	2.08—2.43	2.25±0.02
<i>sol.</i> ♀♀ : (6-striped)	1.96—2.25	2.09±0.01
<i>sol.</i> ♀♀ : (7-striped)	1.93—2.17	2.03±0.007
<i>sol.</i> ♀♀ : (8-striped)	1.91—2.11	1.996±0.032


The E/F ratios in the 8-striped forms are the lowest (Table 4) as compared to the other forms determined earlier (Roonwal, 1949c ; and Roonwal & Nag, 1951). The combined (male and female) E/F ratios for the *four* types of individuals are as follows (Text-fig. 5):— (a) Phase *gregaria* : 6-striped, 2.22. (b) Phase *solitaria* : 6-striped, 2.07 ; 7-striped, 2.02 ; and 8-striped, 1.999.

The sex-ratios also vary in relation to eye-stripe number, as has been shown for the 6- and 7-striped forms (*vide* Roonwal, 1936 ; 1941 ; 1945 ; Nair, 1952*b* and ; Misra, 1953). As in the 7-striped *solitaria* individuals, the proportion of females is higher than males in the 8-striped *solitaria* forms, although the number of available samples of the latter is admittedly small. Of the nine 8-striped individuals collected in the field during 1936-38, 4 were males and 5 females ; and all the five 8-striped individuals bred in the laboratory were females. Thus, we get a ♂ : ♀ ratio of 4 : 10 or 29 per cent ♂ and 71 per cent ♀. The proportion of the sexes in the various types of individuals is roughly as follows :—

- (a) Phase *gregaria* (6-eye-striped) : ♂ 50 : ♀ 50
 (b) { „ *solitaria* (6- „ „) : ♂ 60 : ♀ 40
 „ „ (7- „ „) : ♂ 35 : ♀ 65
 „ „ (8- „ „) : ♂ 29 : ♀ 71

3. *Swarming (gregaria) period 1940-46.*—All the individuals taken from actual swarms had 6-eye-stripes only. However, it was noticed again and again that, both in Baluchistan and Rajasthan, 7-eye-striped forms appeared in the next generation obtained from the *residual* populations left after the departure of the swarms. No 8-striped forms were obtained. The occurrence of 7-striped forms emphasises the fact

that even during swarming years whenever and wherever populations fall, the 7-striped forms begin to appear. Thus, in the winter-spring period, when populations are generally lower than in the summer and autumn periods, there is an appreciable proportion of 7-striped forms, but these



TEXT-FIG. 9.—*Schistocerca gregaria*. Antennae of adults showing variation in the number of segments. (a) Right antenna, with 26 segments, of a 6-eye-striped phase *gregaria* ♀ from a swarm in Peshawar (N.W.F.P.). (b) Left antenna, with 26 segments, of a 6-eye-striped phase *solitaria* ♀ from Lasbela State, Baluchistan. (c) Right antenna, with 27 segments, of a 6-eye-striped phase *solitaria* ♀ from Lasbela State, Baluchistan. (d) Left antenna, with 28 segments, of a 7-eye-striped phase *solitaria* ♀ from Baluchistan. (After Roonwal, 1952a.)

1, first (basal) segment or scape; 26, 27, 28, twenty-sixth, twenty-seventh and twenty-eighth segments respectively.

forms tend to disappear almost completely in the summer and autumn periods. It must be stated that, on the whole, the 1940-46 cycle was considerably milder than the 1926-31 cycle. It is this mildness of swarma-

ing concentrations which, in my opinion, largely accounts for the occurrence of the 7-striped forms throughout the 1940-46 cycle, whereas in the previous cycle no 7-striped forms were found, as seen by an examination of old collections.

The eye-stripe proportions in the various years during the period 1940-46 are given in Table 5.

4. *Non-swarming (solitaria) period 1947-48.*—During the brief *solitaria* interval between the cessation of the last (1940-46) cycle and the beginning of the present one (1949 to date, *i.e.*, 1953), 7-striped forms were again common, although their proportions varied from generation to generation. No 8-striped forms were reported, although it is not unlikely that they may have occurred in small numbers and their existence overlooked.


5. *Swarming (gregaria) period (1949 to date, i.e., 1953).*—The year 1949 saw the occurrence of locust concentrations in the Indian desert regions, the population rising at Kakko (in the Bikaner District, Rajasthan) from below 1,000 per sq. mile in June to over 18,000 per sq. mile in July (Misra, 1952). By 1950, extensive swarmings had started and it was clear that a new cycle was on. The 5-, 6- and 7-eye-stripe proportions in the Kakko concentration were 0.4 : 94 : 5.6, and for the first time in nature a 5-striped individual was found. A sample from a 1950 swarm gave the 6- and 7-eye-stripe proportions as 93 : 7 (Roonwal & Misra, 1952).

TABLE 5.—Summary of data on the relative proportions of 6- and 7-eye-striped forms in Desert Locust populations in India during the swarming cycle of 1940-1946. (*From data of Pruthi, 1951.*)

Year	Periods and eye-stripe proportions					
	Winter—Spring		Summer		Autumn	
	6	7	6	7	6	7
1940 .	?	?	67—100	0—33	85—100	0—15
1941	80—100 (mostly 94—100)	0—20 (mostly 0—6)	90—100	0—10	91—100	0—9
1942 .	64—100	0—36	90—100	0—10	86—100	0—14
1943	77—100	0—23	97—100	0—3	92—100	0—8
1944 . .	92—100	0—8	60—100 (mostly 90—100)	0—40 (mostly 0—10)	47—95	5—53
1945 .	93—100	0—7	100	0	96—100	0—4
1946 .	?	?	100 (one sample only)	0	?	?

(c) DISCUSSION.

It will now be abundantly clear that eye-stripe polymorphism (and, along with it, polymorphism in several other characters as well) periodically appears and disappears. Similarly, the range of distribution of the species periodically swells and shrinks. In short, the following phenomena are noticeable :—(i) The population fluctuates periodically, the 'plague period' of high numbers corresponding with a considerable increase of range of the species. (ii) As regards variation-intensity, the species is comparatively stable during the plague or *gregaria* period but highly polymorphic during the *solitaria* period of *minimum* populations. (iii) The phenomenon is periodic and evidently occurs over a wide range (India to Africa) more or less simultaneously. (iv) It would appear that the 5-, 7- and 8-striped individuals, which make their appearance during *solitaria* periods, when selection-pressure is low probably owing to reduced numbers, are eliminated, presumably by selection, during the *gregaria* periods of high population and increased selection-pressure, leaving behind only 6-striped individuals.




TEXT-FIG. 10.—*Schistocerca gregaria*. Diagrams to show the periodic relationship between (a) population-size, (b) variation-intensity and (c) global area of distribution

Crossing results (unpublished) show that 5-striped, 6-striped and 7-striped individuals produce both 6- and 7- striped forms in F_1 and F_2 generations. Other results need not be discussed here for the present.

The relationship of the three Phenomena, *viz.*, population-size, variation-intensity and global distribution-range are diagrammatically represented in Text-fig. 10.

This dynamic relationship can also be expressed in another way, thus : Let p represent the small population-size and P the large population-


size ; r the reduced global area of the species during the non-swarming (*solitaria*) years and R the increased global area during the swarming (*gregaria*) years ; and, finally, v the reduced variation-intensity (*i.e.*, a variation-stable situation) and V the increased variation-intensity (*i.e.*, a variation-unstable situation) ; then the periodic relationship may be diagrammatically expressed as in Text-fig. 11.



TEXT-FIG. 11.—*Schistocerca gregaria*. Chart to show the correlation and periodicity of flux of population-size, global area occupied and variation-intensity. For explanations, see text.

We may represent this relationship in yet another way, as in Text-fig. 12. The X-axis represents increasing *solitaria*-ness, and the point S on it represents the extreme state (100 per cent) of *solitaria*-ness. The Y-axis represents increasing *gregaria*-ness, and the point G on it represents the extreme state (100 per cent) of *gregaria*-ness. The lengths of the lines PS and PG are assumed to be equal. The points on the line SG represent the position of *solitaria*-ness or *gregaria*-ness on the *solitaria-gregaria* continuum. The shaded portion represents the variation-unstable population (*i.e.*, one with high variation-intensity), while the unshaded portion represents the variation-stable population (*i.e.*, one with zero variation-intensity). It is assumed for the sake of convenience that : (i) in the line PQ dividing the variation-stable area from the variation-unstable area, the point Q lies *midway* between the points S and G on the S-G continuum ; and (ii) this line is a straight one, *i.e.*, the rate of fall of variation-intensity remains constant. The extent of distribution of variability in a population at any one point is read off horizontally, *i.e.*, parallel to the X-axis, and thus the amount of variation-intensity present in a population can be read off easily on this diagram. To take a few examples : Point A on the S-G continuum represents 80 per cent *solitaria*-ness (and, correspondingly, 20 per cent *gregaria*-ness) ; in such a population, out of a total of 8 units about 6/8ths or 3/4ths of it is variation-unstable and the remainder variation-stable (with zero variation-intensity). Similarly, point B represents 70 per cent *solitaria*-ness (and 30 per cent *gregaria*-ness) ; in such a population, out of a total of 7 units about 4/7ths of it are variation-unstable, and the remainder variation-stable. Again, point C represents 30 per cent *solitaria*-ness (and 70 per cent *gregaria*-ness) ; in such a population, with a total of 3 units, all the individuals are variation-stable. These results will, of course, vary with the angle which the line PQ sustains on the X-axis. At point S (of 100 per cent *solitaria*-ness) the entire population is variation-unstable. But, with the point Q being midway on the S-G continuum, the entire population will be variation-stable beyond point Q towards G. In other words, at point Q we get a sudden extinction of variation and the entire population is variation-stable and remains so until point G (100 per cent *gregaria*-ness) is reached.

This idealised condition (of point Q being midway between S and G) need not, and indeed does not, apply to all the variabilities, but varies from character to character. Two examples, which are taken from the data of Roonwal (1945*a*, 1949*c*, *d*), will suffice to illustrate this point. In regard to the character E/F ratio (elytron-length/hind-femur-length) it is known (Roonwal, 1949*c*, p. 161, Text-fig. 1 ; also *vide* Text-fig. 5 above) that it vibrates on the *solitaria-gregaria* continuum in 2 groups or peaks of temporary stability (in the sense of Wright, 1932, *vide* Text-fig. 2 above) as follows :—(i) The *solitaria* group with *three* peaks (*i.e.*, with high variability) for the 6-, 7- and 8-eye-striped forms (with possibly a fourth peak for the 5-eye-striped forms). All these peaks are characterised by low ratios. (ii) The *gregaria*-group with *one* peak (*i.e.*, with comparatively low variability or, ideally speaking, none) of high E/F ratios. The change from the *solitaria* group of peaks to the



TEXT-FIG. 12.—*Schistocerca gregaria*. Idealised diagram to represent the relationship between variation-intensity on the one hand, and the *gregaria* and *solitaria* phases on the other. The shaded area represents the variation-unstable population (*i.e.*, one with great variation-intensity) and the unshaded portion the variation-stable population (*i.e.*, one with no variation), on the assumptions: (i) that the dividing line, PQ, between the two areas lies at a point midway on the line S-G which represents the *solitaria-gregaria* continuum; and (ii) that the line PQ is a straight one.

gregaria group, *i.e.*, from the variation-unstable position to the variation-stable position, takes place *below* the point Q = 50, as is evident from the data of Roonwal (1949*a*, p. 174 ; also *vide* Text-figs. 13 and 14 here). In other words, the sudden change or 'extinction of variation' referred to above occurs very early or long before the movement of S towards G has gone half-way. Roonwal (1949*a*, p. 174) wrote thus in this connection

in regard to E/F ratios : " It may be tentatively assumed that the dividing line between *solitaria* and *gregaria* populations is at $g^*=50$. Actually perhaps, this figure may prove to be a little too high, but it may be adopted until more data are available. "

The position regarding the variability in the eye-stripe character is, however, different. The complete or near-complete extinction of variability in this character occurs long after $Q = 50$. As shown in my " First Hypothesis " (Roonwal, 1945a ; *vide* also Nair, 1952b for discussion), in a mixed 6- and 7-eye-striped (*i.e.*, variation-unstable) population travelling from S to G, the elimination of the 7-striped forms is very rapid after the proportion of about 70 per cent 6-striped forms is reached. In other words, $Q = ca. 70$. In any case, whatever the position of Q on the S-G continuum, the triangle PQS will be shaded to represent the variation-unstable population showing high variation-intensity, and the triangle PQG will then represent the variation-stable population of zero variation-intensity, the proportions of the two at any one point in a population being read off horizontally, *i.e.*, parallel to the X-axis.

Since in the Desert Locust the determining factor in the degree of variation-intensity is the population density rather than absolute numbers, let us see what are the theoretical possibilities. Let n represent the total number of individuals in an area, s the area occupied, and d the density per unit area. If we further assume that

$$d = \frac{n}{s},$$

then there are 3 theoretical possibilities, A, B, C, as regards the change of d . These, together with the conditions under which they will be realised, are listed below :—

(A)— d remains constant : *This condition will be realised when :*

1. s increases ; and n increases proportionately to s .
2. s decreases ; and n decreases proportionately to s .

(B)— d increases : *This condition will be realised when :*

3. s is constant ; and n increases.
4. s decreases ; and n increases.
5. s decreases ; and n is constant.
6. s increases ; and n increases more rapidly than s .
7. s decreases ; and n decreases less rapidly than s .


(C)— d decreases : *This condition will be realised when :*

8. s increases ; and n is constant.

* The symbol g means the " E/F shift figure " as conceived by Roonwal (1949d, p. 171). It means the " shift ", as determined by the E/F ratios, towards *gregaria*-ness in a *solitaria-gregaria* population. Similarly, the symbol e means the " eye-stripe figure " by which is meant the percentage of 6-eye-striped forms in a mixed population of 6- and 7-striped forms (*see* Text-figs. 13 and 14).

9. s is constant ; and n decreases.
10. s increases ; and n decreases.
11. s decreases ; and n decreases more rapidly than s .
12. s increases ; and n increases less rapidly than s .

Subject to the correction that the population is not uniformly distributed throughout the range of distribution but is confined to niches or islands, the following situations would seem to occur in the Desert Locust :—In the *solitaria* phase the population density (in the niches in the permanent breeding grounds) is only about 1,000 per sq. mile, whereas in the *gregaria* phase this figure is about 30,000 per sq. mile, or nearly 30



SERIAL NUMBER OF POPULATION SAMPLE (TABLE 2).

TEXT-FIG. 13.—*Schistocerca gregaria*. Graph showing the correlation between the “E/F shift figure g ” and the “eye-stripe figure e ” in several population samples. (After Roonwal, 1949a.) For further explanations, see text, p. 31, foot-note.

times as great (Table 2). The increase in the area of distribution in the *gregaria* phase is only about 1.7 times that of the *solitaria* phase (*vide supra*, p. 6 & Pl. XV). Thus, the increase of density during the *gregaria* phase takes place by means of situation B(6) mentioned above. And, *vice versa*, during the passage from the *gregaria* to the *solitaria* condition, the situation C(11) evidently comes into play.


The part which the migration or crossing factor might play in controlling variation-intensity in the Desert Locust is not well understood.

But in so far as the larger migratory movements are concerned, *viz.*, the seasonal migrations, they are similar in the *gregaria* and the *solitaria* phases. There is a free exchange of population between the large geographical areas concerned, as the N. African populations migrate seasonally to Arabia, and populations from the latter region migrate seasonally to Iran and India. The reverse movements occur with equal regularity. Thus, we may assume with justification that a thorough mixing up of these populations over vast areas occurs regularly in both phases, with this difference that in the *gregaria* phase the usual bounds of the permanent breeding grounds are erupted and migrations occur over greater distances and areas.

The exact mode of the periodic population-increase itself is not understood, although a number of explanations have been put forward (*vide* Roonwal, 1949a, for discussion). How misleading some of these explanations can be when they are not rigidly checked against facts will be evident from the comparatively recent attempt of Bodenheimer (1937). This author has attempted to explain periodicity on the basis of a difference in the number of annual generations between the migration and the non-migration periods, potential increase, environmental resistance, and so on. His account, however, contains so many errors and unfounded assumptions that it is not surprising that he has completely confused the issues. It will suffice to point out a few of his errors. First, his purely imaginary scheme for the building up of Desert Locust outbreaks runs in the following sequence : Population ebb ; preparatory years (2-6 generations) ; prodromal years (1-3 generations) ; eruption (2-10 generations) ; crisis (1-2 generations) ; and population ebb (a few to many generations). Bodenheimer has left unexplained the precise meaning of the terms 'preparatory', 'prodromal', 'eruption' and 'crisis'. But even if we were to give an approximate meaning to these terms, there is in fact not the slightest factual evidence for the sequence of events mentioned above. All that can be distinguished are the plague or mass-outbreak years (usually showing some sort of a peak) with very high population (of about 30,000 or more adults per sq. mile in swarms), and the non-plague years with a very low but fluctuating population (of not over about 1,000—2,000 non-swarmling adults per sq. mile). However, in the year preceding the start of full-scale swarming, it is now evident that there are certain initial concentrations of population up to about 20,000 per sq. mile for a generation, as was recently found in Kakko in Bikaner District, Rajasthan, in 1949 by Misra (1952). The number of generations assigned by Bodenheimer to each one of his imaginary periods has, of course, no basis in fact.

Secondly, Bodenheimer has claimed that all the species of locusts, such as those of the genera *Schistocerca*, *Locusta*, *Calliptamus* and *Doclostaurus*, which migrate extensively are able to develop more than one annual generation in their *gregaria* phase, while the *solitaria* phase has only one generation. This claim certainly does not hold true for the Desert Locust, *Schistocerca gregaria*, which in the *solitaria* phase has 2-3 annual generations (theoretically perhaps 4), and in the *gregaria* phase 3-4 (theoretically perhaps 5), as shown below. The existence

of 2-3 annual generations (depending on the area) in the *solitaria* phase in nature in western India has been demonstrated by Rao (1942) who concluded that theoretically "there is the possibility of four successive generations following one another in the course of the year, though ordinarily not more than three broods can be expected" (p. 242). It should be stated, however, that in places like Pasni (Mekran Coast, Baluchistan), which is an area of scanty winter rains and no summer rains at all, there is normally only one annual generation, *viz.*, in winter. In the *gregaria* phase, in South-Western Asia, Fortescue-Foulkes (1953, pp. 12-14) has concluded that there may be as many as 2, or sometimes even 3, monsoon generations in India. He has distinguished 3 major



TEXT-FIG. 14.—*Schistocerca gregaria*. Graph obtained when the "eye-stripe figure *e*" is plotted as a function of the "E/F shift-figure *g*". (After Roonwal, 1949a.) For further explanations, see text, p. 31, foot-note.

seasonal breeding areas in S. W Asia as follows :—(i) Only one breeding season a year—"monsoon" breeding (Sind, central and southern Rajasthan, Cutch, Saurashtra, N. Bombay). (ii) Two breeding seasons a year—"winter-early spring" and "main-spring" breeding seasons (S. Iran). (iii) Three breeding seasons a year—"monsoon", "winter-early spring" and "main spring" breeding seasons (N. Pakistan; and the Punjab and Uttar Pradesh in North India). Fortescue-Foulkes concluded (p. 17) that "it is thus theoretically possible for five successive generations to be produced during the year in South-Western Asia, an

estimate which corresponds with the maximum number of generations which might be produced by swarms migrating between North-East Africa and Middle East (Davies 1952). The production of all five generations, however, depends not only on suitable climatic conditions but also on the continuity of swarm displacements between the seasonal breeding zones, and can only be established by a more detailed study of particular breeding seasons and years." It may be mentioned in this connection that under favourable conditions of temperature and soil moisture under laboratory conditions, Husain & Ahmad (1936) have obtained as many as 8 successive generations per year at 40° C., about 7 at 36° C. and 5 at 30° C. in the incubator. I have observed during the years 1935-38 in the *solitaria* phase in Pasni (Mekran Coast, Baluchistan) that whereas in nature there is only one annual breeding in December-February coinciding with the winter rains, in cages under semi-natural conditions with moist soil and an abundance of fresh food, breeding was continuous and about 2½ annual generations were obtained; it is hardly necessary to mention that in summer one generation was over in about 2-3 months, whereas in winter, owing to low temperatures, the period of development was considerably prolonged.

Thirdly, the accuracy of Bodenheimer's claim of the "Sudano-Deccanian" region as being the permanent home of the Desert Locust, has already been refuted above (see p. 4) in so far as the Deccan and the adjoining regions in India are concerned. Fourthly, his statement that the mortality of adults in the so-called imaginal diapause is certainly very great cannot be substantiated. There is no diapause in any stage in the Desert Locust; only, during the winter months (November to February) as compared to the hotter months, the rate of development is very much slowed down and the life-duration, especially of the adults, considerably prolonged as a result of low temperatures. Fifthly, there is no evidence to support Bodenheimer's assertion that mortality is greater in the winter than in the summer months.

Finally, a word may be said about the mutual relationships of the various types of evolutionary Effects or phenomena discussed above, ranging from the Hardy Effect to the New Effect described here. It is difficult to believe that all these Effects, which are in appearance distinct from one another, are completely independent and are caused by totally different mechanisms. Rather, it would seem that the mechanism which brings them into play is fundamentally the same, and that the various Effects are merely different phases or expressions of that mechanism.

A further question which arises is whether the types of polymorphism which periodically arise and are swamped away as in the Desert Locust, can be of any significance in the actual process of speciation. In other words, will these variations ever become stable and thus produce separable specific populations. It is difficult to answer this question, but we have no reason to believe that the variations discussed here are utterly transitory and of no speciation-value.

V.—EVOLUTION OF VARIOUS TYPES IN DESERT LOCUST.


(Text-figure 15.)

Having considered the various forms that exist in the Desert Locust, it would be appropriate to discuss their evolution. The following types have so far been recognized:—

Group (A).—Phase *gregaria* type (6-eye-striped).

Group (B).—Phase *solitaria* types (5-, 6-, 7- and 8-eye-striped types). Two further types have also been recognised, viz., the “blue-grey type” and the “fawn type” (Roonwal, 1945*b*; 1946*b*); the former is more common (91 %) than the latter (9%), and in both types 6- and 7-eye-striped individuals occur.

Leaving aside these 2 colour-types in the phase *solitaria* group, the evolution of the various forms of *Schistocerca gregaria* would appear to be as follows (Text-fig. 15):—



TEXT-FIG. 15.—*Schistocerca gregaria*. Diagram to show the evolution of the various polymorphic forms.

The 6-eye-striped *solitaria* must be regarded as the *basic type*, since it is the most common and the most stable type. Under certain conditions, it produces or evolves into 2 different groups, viz., (i) the *gregaria* phase group in which only 6-eye-striped forms occur; and (ii) the *solitaria* phase group in which, in addition to the 6-striped forms, forms with 5, 7, and 8 eye-stripes occur. It would seem that all these different forms arise directly from the ‘basic type’ (6-eye-striped *solitaria*). This conclusion is supported by preliminary results of the crossing experiments briefly mentioned above (p. 28). The position of the 5-striped forms is not quite clear. Misra (1952) has reported the first and only example of a 5-striped form taken wild in 1949 which was the first year of a new swarming cycle; previous records are all from breeding cages. Tentatively, I am inclined to place the 5-striped forms with the *solitaria* group and as having evolved from the ‘basic type’ (6-striped *solitaria*) rather than from the 6-striped *gregaria*.

VI.—SUMMARY.

1. A number of different evolutionary phenomena or “Effects” are known to exist in regard to the influence of population-structure on the processes of speciation or production of variations. Here I have brought forth evidence, based on the Desert Locust, for the existence of a new type of phenomenon.

2. New data on the land-area occupied by the Desert Locust, *Schistocerca gregaria* (Forsk.), are given with the aid of a world map drawn on an "equal-area" projection. This area is about 16,500,000 sq. miles in the swarming (phase *gregaria*) periods and about 9,500,000 sq. miles in the non-swarming (phase *solitaria*) periods, or in the ratio 1.7 : 1.

3. The density of population and the amplitude and periodicity of population-flux in the Desert Locust are discussed. The population density is about 1,000—2,000 per sq. mile or less in the *solitaria* phase and about 30,000 per sq. mile or more in the *gregaria* phase. The periodicity of plague cycles varies thus : 5-10 years (average 7.1) in the swarming years ; 1-8 years (average 4) in the non-swarming years ; and 7-14 years (average 11.1) for the complete cycles.

4. A critical review is given of the known types of phenomena or "Effects" concerning the production of variations in relation to population density and other characteristics. The various phenomena are : (i) the Hardy Effect ; (ii) the Hagedoorns Effects ; (iii) the Adlerz Effect ; and (iv) the Wright-Dubin Effect.

5. (a) A new phenomenon or "Effect", as evidenced by the case of the Desert Locust, is described. This Effect consists in the fact that intraspecific variability (polymorphism) is sharply increased during the non-swarming or phase *solitaria* periods of *minimum* populations, and almost completely obliterated during the swarming or phase *gregaria* periods of *maximum* populations. The *gregaria* phase is variation-stable and the *solitaria* phase variation-unstable.

(b) The periodic appearance and extinction of these polymorphic variations are traced over a period of 25 years (1926-50) during which the phenomenon was observed in 3 swarming periods and 2 non-swarming periods.

(c) Some new data on the characteristics of the 8-eye-striped individuals are provided. Their E/F ratio is the lowest among the phase *solitaria* group of individuals, being lower than in the 6- and 7-striped *solitaria* forms. They also have the highest proportion of females, the ♂ : ♀ ratio being 29 : 71.

(d) The various factors which lead to the increase or decrease of population density are discussed theoretically.

6. The evolution of the various types of polymorphic forms in the Desert Locust is traced. The 6-eye-striped phase *solitaria* form is considered as the *basic type* from which the other types are believed to have been derived directly in two directions : (i) the 'gregaria phase group' with a single type (6-eye-striped) ; and (ii) the 'solitaria phase group' with 5-, 6-, 7- and 8-eye-striped types. The exact position of the 5-eye-striped type is rather problematical owing to want of sufficient data.

VII. REFERENCES.

* Those not seen in the original are marked with an asterisk.

ADLERZ, G. (1902). Periodische Massvermehrung als Evolutionsfaktor.—*Biol. Cbl.*, Leipzig, 22, pp. 108-119.

- BODENHEIMER, F. S. (1937). Factors controlling locust populations.—*Proc. 4th int. Locust Conf. (Cairo, 1936)*, Cairo, Appendix 9, pp. 1-11.
- CARTER, G. S. (1951). *Animal Evolution. A Study of Recent Views of Its Causes*. xv+368 pp.—London (Sidgwick and Jackson).
- *CHETVERIKOV (TSCHETWERIKOFF), S. S. (1926). *J. Biol. exp.*, Moscow, 2, pp. 3-54.
- CHOPARD, L. (1932). Voyage de MM. L. Chopard et A. Mequignon aux Açores (aout-septembre 1930). I. Orthopteres.—*Ann. Soc. ent. Fr.*, Paris, 101, pp. 55-68, 2 pls.
- CUENOT, L. (1936). *L'Espèce*. 310 pp.—Paris (Doin & Co.).
- DARWIN, C. (1876). *The Origin of Species by Means of Natural Selection or the Preservation of Favoured Races in the Struggle for Live*. (6th ed.).—London.
- DAVIES, D. E. (1952). Seasonal breeding and migrations of the Desert Locust (*Schistocerca gregaria* Forskal) in north-east Africa and the Middle East.—*Anti-Locust Mem.*, London, 4, 57 pp., 13 maps, 2 figs.
- DOBZHANSKY, T. (1941). *Genetics and the Origin of Species*. (2nd ed.) xviii+446 pp.—New York.
- DROUET, H. (1861). *Elemente di la fauna Acoreene*.—Paris (J. R. Bailliere).
- DUBININ, N. P. (1931). Genetico-automatic processes and their significance for the mechanism of organic evolution. (In Russian).—*J. Biöl. exp.*, Moscow, 7, pp. 463-479.
- DUBININ, N. P. AND ROMASCHOFF, D. D. (1932). Die genetische Struktur der Art und ihre Evolution. (In Russian, with German summary.)—*Biol. Zh.*, Moscow, 5, pp. 939-976.
- ELTON, C. (1924). Periodic fluctuations in numbers of animals: Their causes and effects.—*Brit. J. exp. Biol.*, Cambridge, 2, pp. 119-163.
- ELTON, C. (1930). *Animal Ecology and Evolution*.—Oxford.
- ELTON, C. (1951). *Animal Ecology* (5th impr.). xx+209 pp., 8 pls.—London (Sidgwick and Jackson).
- FAURE, J. C. (1932). The phases of locusts in South Africa.—*Bull. ent. Res.*, London, 23, (3) pp. 293-405, 25, pls., 1 figd. map.
- FISHER, R. A. (1922). On the dominance ratio.—*Proc. R. Soc. Edinb.*, Edinburgh, 42, p. 323.
- FISHER, R. A. (1930a). *The Genetical Theory of Natural Selection*. 272 pp.—Oxford (Clarendon Pr.).
- *FISHER R. A. (1930b). The distribution of gene ratios for rare mutations.—*Proc R. Soc. Edinb.*, Edinburgh, 50, pp. 204-224.
- FISHER, R. A. (1937a). The relation between variability and abundance shown by the measurements of the eggs of British nesting birds.—*Proc. R. Soc. Lond. (B)*, London, 122, pp. 1-26.

- FISHER, R. A. (1937*b*). The wave of advance of advantageous genes.—*Ann. Eugen.*, Cambridge, **7**, p. 355.
- FISHER, R. A. and FORD, E. B. (1926). Variability of species.—*Nature*, London, **118**, pp. 515-516.
- FISHER, R. A. and FORD, E. B. (1928). The variability of species in the Lepidoptera with reference to abundance and sex.—*Trans. ent. Soc. Lond.*, London, **76**, pp. 367-384, 1 pl.
- FOOD AND AGRICULTURAL ORGANISATION (1952). The Desert Locust (*Schistocerca gregaria* Forsk.). International control measures against plague swarms.—*F. A. O. Fact Sheet*, Rome, No. **X2**, cyclostyled, pp. 1-8.
- FORD, E. B. (1937). Problems of heredity in the Lepidoptera.—*Biol. Rev.*, Cambridge, **12**, pp. 461-503.
- FORD, E. B. (1940). Polymorphism and taxonomy. pp. 493-513.—In J. Huxley's *The New Systematics*. Oxford.
- FORD, H. D. and FORD, E. B. (1930). Fluctuation in numbers and its influence on variation in *Melitaea aurinia* Rott. (Lepidoptera).—*Trans. ent. Soc. Lond.*, London, **78**, pp. 345-351, 1 pl.
- FORTESCUE-FOULKES, J. (1953). Seasonal breeding of the Desert Locust (*Schistocerca gregaria* Forskal) in South-Western Asia.—*Anti-Locust Mem.*, London, **5**, 36 pp., 14 maps (figd.), 1 fig.
- GAUSE, G. F. (1927). Zur Kenntnis der Variabilität der Wanderheuschrecke.—*Z. angew. Ent.*, Munich, **13**, pp. 247-266.
- HAGEDOORN, A. L. and HAGEDOORN, A. C. (1921). *The Relative Value of Processes Causing Evolution*.—The Hague (Martinus Nijhoff).
- HALDANE, J. B. S. (1932). *The Causes of Evolution*. 235 pp.—London (Longmans, Green & Co.); and New York (Harper Bros.).
- HARDY, G. H. (1908). Mendelian proportions in mixed populations.—*Science* (N. S.), New York, **28**, pp. 49-50.
- HUSAIN, M. A. and AHMAD, T. (1936). Studies on *Schistocerca gregaria* Forsk. II. The biology of the Desert Locust, with special reference to temperature.—*Indian J. agric. Sci.*, Delhi, **6**, pp. 188-262.
- HUXLEY, J. (Ed. by). (1940). *The New Systematics*. 583 pp.—Oxford.
- HUXLEY, J. (1944). *Evolution: The Modern Synthesis*. (3rd impr.). 645 pp.—London.
- KELLAWAY, G. P. (1949). *Map Projections* (2nd ed.). viii+127 pp.—London (Methuen & Co.).
- *KOLMOGOROV, A. (1935). [Mathematical distribution of gene frequencies in populations.].—*C. R. Acad. Sci. U. R. S. S.*, Moscow, **3** (7), pp. 129-132.
- LONG, D. B. (1952). Some problems of polymorphism in insects.—*Proc. R. ent. Soc. Lond. (A)*, London, **27** (10-12), pp. 99-110.

- LUCAS, W. J. (1920). *A Monograph of the British Orthoptera*. xii+264 pp., 25 pls.—London.
- MAXWELL-DARLING, R. C. (1934). The solitary phase of *Schistocerca gregaria* in North-Eastern Kordofan (Anglo-Egyptian Sudan).—*Bull. ent. Res.*, London, **25** (1), pp. 63-84, 2 pls.
- MAXWELL-DARLING, R. C. (1936). The outbreak centres of the *Schistocerca gregaria* Forsk. on the Red Sea Coast of the Sudan.—*Bull. ent. Res.*, London, **27** (1), pp. 37-70, 2 pls.
- MAXWELL-DARLING, R. C. (1937). The outbreak centres of the Desert Locust (*Schistocerca gregaria* Forsk.) in Arabia.—*Bull. ent. Res.*, London, **28** (4), pp. 605-618, 1 pl.
- MISRA, S. D. (1952). Studies of eye-stripe characteristics and the biometrical ratios of the body-parts in a concentration of Desert Locust individuals met with in July 1949 in the Rajputana Desert, that brought about incipient swarming and thus started a new locust cycle in India. 100-112 pp. (*In* Misra, Nair & Roonwal: Studies in intraspecific variation. Part VI, *vide infra*).—*Indian J. Ent.*, New Delhi, **14** (2), pp. 95-152.
- MISRA, S. D. (1953). Confirmation of Roonwal's Hypotheses for prediction of swarming of Desert Locust from data collected in 1949.—*Saugar Univ. J.*, Sagar, **1** (2), pp. 233-238.
- MISRA, S. D., NAIR, K. R. and ROONWAL, M. L. (1951). Dynamics of eye-stripe composition, biometrical ratios and certain other characters during phase transformation in the Desert Locust.—*Current Sci.*, Bangalore, **20** (6), pp. 164-165.
- MISRA, S. D., NAIR, K. R. and ROONWAL, M. L. (1952). Studies in intraspecific variation. Part VI. Dynamics of variability in respect of eye-stripe characters, sex-ratios and body-size of Desert Locust populations during the initial years (1949-1950) of a new swarming cycle in India, together with a statistical note on Roonwal's Hypotheses on prediction of swarming.—*Indian J. Ent.*, New Delhi, **14** (2), pp. 95-152.
- NAIR, K. R. (1952a). Use of measurements of more than one biometric character regarding size of body-parts for discriminating between phases in the case of six-eye-striped specimens of the Desert Locust. 126-136 pp. (*In* Misra, Nair & Roonwal: Studies in intraspecific variation. Part VI, *vide supra*).—*Indian J. Ent.*, New Delhi, **14** (2), pp. 95-152.
- NAIR, K. R. (1952b). A statistical note on Dr. Roonwal's Hypotheses for prediction of the swarming of the Desert Locust. 136-146 pp. (*In* Misra, Nair & Roonwal: Studies in intraspecific variation. Part VI, *vide supra*).—*Indian J. Ent.*, New Delhi, **14** (2), pp. 95-152.
- PREDETCHENSKY, S. A. (1935). The annual cycle of the Desert Locust (*Schistocerca gregaria* Forsk.), its migrations and periodicity in Persia and adjacent countries of tropical and subtropical Asia. (*In* Russian.)—*Bull. Plant Prot.* (1, *Ent.*) Leningrad, **12**, pp. 1-135.

- PRUTHI, H. S. (1951). The Desert Locust cycle of 1940-46 in India : Its progress and control.—*Indian J. Ent.*, New Delhi, **12** (1) (of June 1950), pp. 67-98 ; **12** (2) (of December 1950), pp. 191-232 ; and **13** (1) (of June 1951), pp. 29-63.
- RAO, Y. RAMACHANDRA. (1936). The locust incursion of 1935 in North East India. Its significance in the study of locust problem.—*Indian J. agric. Sci.*, Delhi, **6**, pp. 1031-1053.
- RAO, Y. RAMACHANDRA. (1937). Some outbreak centres of *Schistocerca gregaria* in Mekran.—*Proc. 4th int. Locust Conf. (Cairo, 1936)*, Cairo, Appendix 30, pp. 1-9.
- RAO, Y. RAMACHANDRA. (1942). Some results of studies on the Desert Locust (*Schistocerca gregaria*, Forsk.) in India.—*Bull. ent. Res.*, London, **33**, pp. 241-265.
- RAO, Y. RAMACHANDRA. (1945). The areas of origin of locust flights in different parts of India with reference to the question of their control.—*Current Sci.*, Bangalore, **14** (2), pp. 31-33.
- ROBSON, G. C. (1928). *The Species Problem*.—London.
- ROBSON, G. C. and RICHARDS, O. W. (1936). *The Variation of Animals in Nature*. xvi+425 pp., 2 col. pls.—London.
- ROONWAL, M. L. (1936). On the existence of two different types of striped eyes among solitary type specimens of the Desert Locust, *Schistocerca gregaria* Forsk.—*Current Sci.*, Bangalore, **5** (1), p. 24.
- ROONWAL, M. L. (1937). [Development of eye-stripes in the Desert Locust, *Schistocerca gregaria* (Forsk.)].—In Y. R. Rao's *Rept. Work Res. Staff under Locust Res. Entom. to Imp. Counc. agric. Res., Karachi, during 1936*. Simla (Govt. India Pr.), pp. 25-26 ; and 150-151.
- ROONWAL, M. L. (1941). Sex-ratios and eye-stripes in a swarm of the Desert Locust.—*Indian J. Ent.*, New Delhi, **3** (2), pp. 340-341.
- ROONWAL, M. L. (1944). Do the army-worm moths in India have phases ?—*Current Sci.*, Bangalore, **13**, pp. 134-135.
- ROONWAL, M. L. (1945a). New hypotheses for the prediction of the swarming of the Desert Locust.—*Bull. ent. Res.*, London, **35** (4), pp. 391-393.
- ROONWAL, M. L. (1945b). Two colour-types in *solitaria*-phase adults and hoppers of the Desert Locust.—*Nature*, London, **155**, p. 792.
- ROONWAL, M. L. (1945c). Remarks on variation in eye-stripes, etc., in the Desert Locust and the species-problem.—*Proc. 31st Indian Sci. Congr. (Delhi, 1944)*, Pt. 4, *Discussions*, Calcutta, p. 33.
- ROONWAL, M. L. (1946a). On a new phase character, the metasternal interspace, in the Desert Locust, *Schistocerca gregaria* (Forsk.) [Orthoptera, Acrididae].—*Proc. R. ent. Soc. Lond. (A)*, London, **21** (1-3), pp. 13-15. *Erratum* : Page 13.—In explanation of Fig. 1, line 2 : For " fig. 1a " read " fig. 2a."

- ROONWAL, M. L. (1946b). Studies in intraspecific variation. I. On the existence of two colour-types in the adults and hoppers of the *solitaria* phase in the Desert Locust, *Schistocerca gregaria* (Forskål) [Orthoptera, Acrididae].—*Rec. Indian Mus.*, Benares [Calcutta], **44** (4), pp. 369-374.
- ROONWAL, M. L. (1946c). On variation in the number of ovarioles and its probable origin in the Desert Locust, *Schistocerca gregaria* (Forskål). [Orthoptera, Acrididae].—*Rec. Indian Mus.*, Benares [Calcutta], **44** (4), pp. 375-384.
- ROONWAL, M. L. (1946d). Studies in intraspecific variation. II. New rules governing the correlation between normal- and extra-moulting and directional reversal of the elytron-wing complex in the Desert Locust and other Acrididae (Orthoptera).—*Indian J. Ent.*, New Delhi, **7** (1-2) [Dec. 1945], pp. 77-84.
- ROONWAL, M. L. (1947a). Variation and structure of the eyes in the Desert Locust, *Schistocerca gregaria* (Forskål).—*Proc. R. Soc. London (B)*, London, **134** (no. 875), pp. 245-272, 3 pls. *Errata*: See same journal, **134** (no. 877), p. 545.
- ROONWAL, M. L. (1947b). On the simulation of background colours by the Desert Locust, *Schistocerca gregaria* (Forskål) [Orthoptera, Acrididae]. Experiments with painted boxes.—*Proc. nation. Inst. Sci. India*, Calcutta, **13** (1), pp. 25-28.
- ROONWAL, M. L. (1947c). On variation in the number of hind-tibial spines in the Desert Locust, *Schistocerca gregaria* (Forskål) [Orthoptera, Acrididae].—*Indian J. Ent.*, New Delhi, **8** (1) [June 1946], pp. 71-77.
- ROONWAL, M. L. (1947d). Evolutionary significance of periodicity of variation-intensity and population-flux in the Desert Locust.—*Nature*, London, **159**, pp. 872-873.
- ROONWAL, M. L. (1947e). Evolutionary and practical aspects of variation in locusts. [Abstract of Special Lecture delivered before the Zoological Society of Bengal on 3rd February 1947 at Calcutta].—*Sci. & Culture*, Calcutta, **12** (10), pp. 484-485.
- ROONWAL, M. L. (1949a). The geographical and geological aspects of the migration of animals. [Lecture delivered before the National Geographical Society of India on 4th December, 1948, at Banaras.]—*Bull. nation. geogr. Soc. India*, Banaras, No. 11, pp. 1-33, 2 pls.
- ROONWAL, M. L. (1949b). Modern trends in systematics. (Presidential Address to the Section of Zoology and Entomology, Indian Science Congress, 1949.)—*Proc. 36th Indian Sci. Congr. (Allahabad, 1949)*, *Presidential Addresses*, Calcutta, pp. 111-138.
- ROONWAL, M. L. (1949c). Studies in intraspecific variation. III. Body-size and biometrical ratios in various types of individuals of the Desert Locust, *Schistocerca gregaria* (Forskål). [Orthoptera, Acrididae].—*Rec. Indian Mus.*, Benares [Calcutta], **45** (2-3) [1947], pp. 149-165.

- ROONWAL, M. L. (1949*d*). Studies in intraspecific variation. IV The role of some variations, *e.g.*, eye-stripes, etc., as population indicators in the Desert Locust, and their practical importance.—*Rec. Indian Mus.*, Benares [Calcutta], **45** (2-3) [1947], pp. 167-180.
- ROONWAL, M. L. (1950).—Locust and grasshopper damage to Indian crops.—*Chemical Age*, Bombay, No. 2, pp. 175-182, 1 col. pl.
- ROONWAL, M. L. (1952*a*). Variation and post-embryonic growth in the number of antennal segments in the *phadká* grasshopper (*Hieroglyphus nigrorepletus* Bolivar), with remarks on the Desert Locust and other Acrididae (Insecta : Orthoptera).—*Proc. nation. Inst. Sci. India*, Calcutta, **18** (3), pp. 217-232, 1 figd. table.
- ROONWAL, M. L. (1952*b*). General introduction. Pp. 96-100. (In Misra, Nair and Roonwal : Studies in intraspecific variation. Part VI. *vide supra*).—*Indian J Ent.*, New Delhi, **14** (2), pp. 95-152.
- ROONWAL, M. L. (1954). On a new evolutionary phenomenon : The sharp increase of intraspecific variation in minimum populations, as evidenced by the Desert Locust. (A preliminary account).—*Bull. nation. Inst. Sci. India*, Calcutta, No. 3. (*In press*).
- ROONWAL, M. L. and MISRA, S. D. (1952). Variability in a sample of Desert Locust population taken from a swarm in the Rajputana Desert, India, in 1950, the second year of a swarming cycle, together with a discussion on the dynamics of phase-transformation. Pp. 112-126. (In Misra, Nair and Roonwal : Studies in intraspecific variation. Part VI, *vide supra*).—*Indian J Ent.*, New Delhi, **14** (2), pp. 95-152.
- ROONWAL, M. L. and NAG, M. K. (1950). Statistical analysis of variability in body-size, etc., of various types of individuals of Desert Locust (Insecta : Orthoptera : Acrididae).—*Proc. 37th Indian Sci. Congr. (Poona, 1950)*, Pt. 3, *Abstracts*, Calcutta, p. 249.
- ROONWAL, M. L. and NAG, M. K. (1951). Studies in intraspecific variation. Part V Statistical supplement to the analysis of biometrical data on body-size, etc., of various types of individuals of the Desert Locust, presented in Part III.—*Rec. Indian Mus.*, Calcutta, **47** (1) [1949], pp. 13-23.
- SCUDDER, S. H. (1883). [Swarms of *Schistocerca gregaria* in the Atlantic].—*Psyche*, Boston, **2**, p. 124.
- SINGH, G. and BHATIA, K. R. (1952). Population density of hopper bands of the Desert Locust (*Schistocerca gregaria* Forsk.) in different instars.—*Indian J. Ent.*, New Delhi, **14** (2), pp. 161-164.
- TIMOFFEFF-RESSOVSKY, N. W. (1940). Mutation and geographical variation. Pp. 72-136.—In J. Huxley's *The New Systematics*.—Oxford.
- UVAROV, B. P. (1921). A revision of the genus *Locusta*, L. (= *Pachytylus*, Fieb.), with a new theory as to the periodicity of migrations of locusts.—*Bull. ent. Res.*, London, **12** (2), pp. 135-163.

- UVAROV, B. P. (1923). Notes on locusts of economic importance, with some new data on the periodicity of locust invasions.—*Bull. ent. Res.*, London, **14** (1), pp. 31-39.
- UVAROV, B. P. (1928). *Locusts and Grasshoppers. A Handbook for Their Study and Control.* 352 pp.—London (Imp. Inst. Entom.).
- UVAROV, B. P. (1932). The locust outbreak in Africa and Western Asia, 1925-1931.—*Econ. Advis. Counc. Committ. on Locust Control*, London, 87 pp., 8 maps.
- VAISSIÈRE, P. (1933). Le criquet pèlerin (*Schistocerca gregaria* Forsk.) aux Açores. [Orth. Acrididae].—*Bull. ent. Soc. Fr.*, Paris, **38**, pp. 85-86.
- VAYSSIÈRE, P. (1936). Etat de recherches sur les sauterelle migra trices.—*Rev. Ass. Fr. Sci.*, Paris, **64** (3), pp. 111-118.
- VOLTERRA, V (1931). Variations and fluctuations of the number of individuals in animal species living together. Pp. 409-448.—*In* R. N. Chapman: *Animal Ecology, with Especial Reference to Insects*, x+464 pp. New York & London (McGraw-Hill Book Co.).
- WALKER, F. (1870). *Catalogue of the Specimens of Dermaptera and Saltatoria in the Collection of the British Museum.* Part III. *Locustidae (concluded) and Acrididae (part).* Pp. 4+425-604.—London (Brit. Mus. Nat. Hist.).
- WALOFF, Z. (1946a). Seasonal breeding and migrations of the Desert Locust (*Schistocerca gregaria* Forskål) in Eastern Africa.—*Anti-Locust Mem.* London, **1**, pp. 1-74.
- WALOFF, Z. (1946b). A long range migration of the Desert Locust from Southern Morocco to Portugal, with an analysis of concurrent weather conditions.—*Proc. R. ent. Soc. Lond. (A)*, London, **21** (10-12), pp. 81-84.
- WILLIS, A. G. (1922). *Age and Area. A Study in Geographical Distribution and Origin of Species.*—Cambridge.
- WRIGHT, S. (1931a). Evolution in Mendelian populations.—*Genetics*, Brooklyn, N. Y., **16**, pp. 97-159.
- *WRIGHT, S. (1931b). Statistical theory of evolution.—*Proc. Amer. Stat. Assoc.*, **26** (Suppl.), pp. 201-208.
- WRIGHT, S. (1932). The roles of mutation, inbreeding, crossbreeding and selection in evolution.—*Proc. 6th int. Congr. Genet. (Ithaca)*, **1**, pp. 356-366.
- WRIGHT, S. (1935). Evolution in populations in approximate equilibrium.—*J. Genet.*, Cambridge, **30**, pp. 257-266.
- WRIGHT, S. (1940a). The statistical consequences of Mendelian heredity in relation to speciation. Pp. 161-183.—*In* J. Huxley's *The New Systematics.*—Oxford.
- WRIGHT, S. (1940b). Breeding structure of populations in relation to speciation.—*Amer. Nat.*, **74**, pp. 232-248.
- WRIGHT, S. (1946). The differential equation of the distribution of gene frequencies.—*Proc. nation. Acad. Sci.*, Washington, **31**, pp. 382-389.

VIII.—EXPLANATION OF PLATES XV—XVIII

LETTERING


<i>b.</i> , basement membrane.	<i>nu. s.</i> , nuclei of secondary pigment cells.
<i>c. c.</i> , crystalline cone.	<i>om.</i> , ommatidia.
<i>c. g.</i> , cerebral ganglia.	<i>op.</i> , opticon.
<i>cr.</i> , cornea.	<i>op. n.</i> , optic nerve.
<i>cr. c.</i> , corneagen cells.	<i>p.</i> , primary pigment cells.
<i>cr. i.</i> , inner layer of cornea.	<i>p. cp.</i> , periopticon.
<i>cr. o.</i> , outer layer of cornea.	<i>pr. n. f.</i> , post-retinular nerve fibres.
<i>e. op.</i> , epipticon.	<i>pr. p. i.</i> , post-retinular pigment layer.
<i>fn.</i> , fenestra or hole in basement membrane.	<i>r.</i> , retinula.
<i>n. c.</i> , nerve cells.	<i>r. c.</i> , retinular or visual cells.
<i>n. c. s.</i> , sheath of nerve cells.	<i>r. n. f.</i> , retinular nerve fibres
<i>n. f.</i> , nerve fibres.	<i>rh.</i> , rhabdom.
<i>nu. c. c.</i> , nuclei of crystalline cone (Semper's nuclei).	<i>rp.</i> , reddish brown pigment granules.
<i>nu. p.</i> , nuclei of primary pigment cells.	<i>s.</i> , secondary pigment cells.
<i>nu. r. d.</i> , distal nuclei of retinular or visual cells.	<i>t. d.</i> , distal or small tracheae.
<i>nu. r. p.</i> , proximal nuclei of retinular or visual cells.	<i>t. p.</i> , proximal or large tracheae.
	<i>y. p.</i> , dirty-yellow pigment granules.

PLATE XV

Map of the World, showing the distribution of the Desert Locust, *Schistocerca gregaria* (Forsk.) in the *solitaria* or non-swarming phase (diagonals) and the *gregaria* or swarming phase (area enclosed by thick line). The former is *ca.* 9,500,000 sq. miles, and the latter *ca.* 16,500,000, or 1.7 times as great.

The map has been drawn on a Re-centred Sinusoidal Projection (Bartholomew) developed from the Sanson-Flamsteed's Projection which is area-true for all lands and in which distortion is reduced to a minimum. It is particularly accurate for the African region where lies the bulk of the distribution range of the Desert Locust.

The natural scale on which the original map was drawn was 1 : 100,000,000. A scale (in cm.) which was *natural-size* in the *original map* is also given for calculating the degree of reduction from the original during reproduction.



Roonwal.

Distribution of the Desert Locust


PLATE XVI

Photographs of the Desert Locust, *Schistocerca gregaria* (Forsk.) in the *gregaria* (swarming) and the *solitaria* (non-swarming) phases. FIGS. 1 and 2 approximately natural size ; FIG. 3 enlarged.


FIG. 1.—A specimen in the *gregaria* phase, taken from a dense swarm in Lucknow (Uttar Pradesh) in 1930.

FIG. 2.—A specimen in the *solitaria* phase, obtained in Pasni (Baluchistan coast) in 1936.


FIG. 3.—Head and pronotum of the same specimen as in Fig. 2 ; enlarged. Note the eye-stripes.



1



2



3

Roonwal.


Desert Locust

PLATE XVII

Hoppers of the Desert Locust, *Schistocerca gregaria* (Forsk.) in the fifth or final hopper stage ; these, on moulting, give rise to the fully-winged adults. The *central* figure in each row is the hopper in side view, about $1\frac{1}{2}$ times natural size ; the figures on the *right* represent the pronotal shield in lateral view, greatly enlarged ; those on the *left* represent a portion of the abdominal segments in lateral view, greatly enlarged. (After Roonwal, 1950).

Upper row : In the *gregaria* or swarming phase, produced by breeding in crowded conditions.

Lower row : In the *solitaria* or non-swarming phase, produced by breeding in comparatively thinned conditions.



Roonwal

Hoppers of the Desert Locust

PLATE XVIII

Compound eyes of *Schistocerca gregaria* (Forsk.) All figures are semi-diagrammatic camera-lucida drawings of sections of eyes, to show the differential pigment distribution in the *solitaria* and the *gregaria* phases. (After Roonwal, 1947a.)

FIG. 1.—Median-longitudinal section of an ommatidium of the unstriped eye of a black (phase *gregaria*) fifth-stage hopper, showing the component parts. Section bleached in Grenacher's mixture and stained with Delafield's haematoxylin and eosin.

FIG. 2.—Median-longitudinal section of an ommatidium of the unstriped eye of a black (phase *gregaria*) fifth stage hopper, showing the distribution of pigment. Section unbleached and unstained.

FIG. 3.—Ditto, in the interstripe region of the striped eye of a green (phase *solitaria*) fifth stage hopper. Note the absence of pigment in the secondary pigment cells.

FIG. 4.—Transverse section of an ommatidium of the striped eye of a green (phase *solitaria*) fifth-stage hopper passing across the distal region of the crystalline cone (line A-A in Fig. 1), to show the structure of the various parts. Section bleached in Grenacher's mixture and stained with Delafield's haematoxylin and eosin. Greatly enlarged.


FIG. 5.—Ditto, across the middle region of the primary pigment cells (line B-B in Fig. 1). Greatly enlarged.

FIG. 6.—Ditto, across the proximal region of the primary pigment cells (line C-C in Fig. 1). Greatly enlarged.

FIG. 7.—Ditto, across the distal nuclei of the reticular cells (line D-D in Fig. 1). Greatly enlarged.

FIG. 8.—Ditto, across the proximal nuclei of the reticular cells (line E-E in Fig. 1). Greatly enlarged.

FIG. 9.—Ditto, across the basement membrane. Greatly enlarged.



Roonwal.

Ommatidia of compound eyes of the Desert Locust

SILUROID FISHES OF INDIA, BURMA AND CEYLON.

XIV.—FISHES OF THE GENUS *MYSTUS* SCOPOLI*

By K. C. JAYARAM, M.Sc., *Zoological Survey of India, Indian Museum*
CALCUTTA.

(Plate XIX.)

The fishes of the Bagrid genus *Mystus* Scopoli are widely distributed in the Oriental and Palaearctic regions. The nomenclature of the genus has been a subject matter of controversy and in a recent note¹ it has been shown that *Mystus* Gronow as emended by Scopoli with *Silurus pelusius* Solander as the type is the correct name for this group of fishes. In this article a systematic revision of the genus and its geographic distribution is given.

I am indebted to Dr. S. L. Hora, for his kindness in suggesting the revision, in placing the material at my disposal and also for constant guidance, supervision and encouragement.

Genus *Mystus* Gronow (*emend.* Scopoli)

1763. *Mystus*, Gronovius, *Zoophylacii Gronoviana* I, no. 388, p. 124, pl. viiia, fig. 6 (*Type-species* :—*Mystus anguillaris* Meuschen).
1777. *Mystus*, Scopoli, *Introd. Hist. Nat.* p. 451.
1781. *Mystus* (in part), Meuschen in Gronow's *Zoophylacii Gronoviana* III.
1794. *Silurus* (in part), Solander in Russell's *Nat. Hist. Aleppo* II, p. 210, pl. vii, fig. 1.
1797. *Silurus* (in part), Bloch, *Ichthyol. Hist. Nat.* XI, p. 40, pl. ccclxxi, fig. 2.
1822. *Pimelodus* (in part), Hamilton, *Fish. Ganges*, p. 160.
1839. *Bagrus* (in part), Valenciennes, *Hist. Nat. Poiss.* XIV, p. 8.
1841. *Platystoma* (in part), *Pimelodus* (in part), Sykes, *Trans. zool Soc. Lond.* II, pp. 371,374.
1841. *Bagrus* (in part), Valenciennes in Jacquemont's *Voy. Ind.* pls. xvi & xvii
1849. *Bagrus* (in part), Jerdon, *Madras J. Lit. Sci.* XV, part. 2, p. 335.
1856. *Macrones*, Duméril, *Mem. Acad. Sci. Paris* XXVII, part I, p. 484 (*Type species* :—*Bagrus lamarii* Valenciennes; name preoccupied in Coleoptera, *Entomologist* I (3), p. 33 (1841)).
1858. *Bagrus* (in part), Bleeker, *Ichthyol. Archipel. Indi. Prodr.* I *Siluri.*, p. 144.
1859. *Bagrus* (in part), Bleeker, *Act. Soc. Sc. Indo-Neerl.* VI, p. 257.
1862. *Bagrus*, *Macrones*, *Hypselobagrus*, *Hemibagrus* & *Aspidobagrus*, Bleeker, *Atlas Ichthyol. Ind. Orient. Neerl.* II, pp. 9, 10.
1863. *Bagrus*, *Macrones*, *Hypselobagrus*, *Hemibagrus*, *Aspidobagrus*, Bleeker, *Ned. Tijdschr. Dierk.* I, p. 96.

*Part of thesis (condensed), approved for the M.Sc., Research degree of the University of Madras (1952).

¹Jayaram, K. C. *Copeia*, (In press).

1864. *Macrones*, Günther, *Cat. Fish. Brit. Mus.* V, p. 74.
 1865. Hara, *Hemibagrus, Hypselobagrus*, Day, *Fish. Malabar*, pp. 184, 186, 187.
 1871. *Macrones*, Day, *Proc. zool. Soc. Lond.*, p. 705.
 1877. *Macrones*, Day, *Fish. India*, p. 442.
 1889. *Macrones*, Day, *Fauna Brit. India Fishes I*, p. 146.
 1890. *Macrones*, Vinciguerra, *Ann. Mus. Stor. nat. Genova IX*, part 2, p. 211.
 1901. *Hypselobagrus, Hemibagrus*, Stiendachner *Abh. Senckenb. naturf. Ges.* XXV, p. 446.
 1911. *Macrones*, Regan, *Ann. Mag. nat. Hist.* (8) VIII, p. 562.
 1913. *Macrones*, Weber & de Beaufort, *Fish. Indo. Austral. Archipel.* II, p. 336.
 1919. *Aoria*, Jordan, *Proc. Acad. nat. Sci. Philad.* LXX, p. 341 (name pre-occupied in Coleoptera, *J. ent. London II*, p. 149 (1863)).
 1925. *Macrones*, Nichols, *Amer. Mus. Novit.* (185), p. 2.
 1926. *Aoria*, Oshima, *Annot. zool. jap.* p. 4.
 1928. *Mystus*, Fowler, *J. Bombay nat. Hist. Soc.* XXXIII, p. 105.
 1929. *Aoria*, Prashad & Mukerji, *Rec. Indian Mus.* XXXI, p. 178 (foot-note).
 1934. *Mystus*, Hora & Mukerji, *ibid.* XXXVI, p. 335 (foot-note).
 1934. *Mystus*, Smith, *J. Siam Soc. nat. Hist.* IX, number 3, p. 294.
 1937. *Mystus*, Herre & Myers, *Bull. Raffles Mus.* XIII, p. 68 (foot-note).
 1939. *Sperata*, Holly, *Zool. Anz.*, CXXV, p. 43 (*Type-species* :—*Silurus vittatus* Bloch).
 1939. *Aorichthys*, Wu, *Sinensia*, X, p. 131.
 1940. *Macronoichthys*, White & Moy Thomas, *Ann. Mag. nat. Hist.* V, p. 505
 1943. *Aoria*, Nichols in *Nat. Hist. Central Asia*, IX, p. 36.
Mystus, Smith, *Bull. U. S. nat. Mus.* (188), p. 382.
 1948. *Mystus*, Hora, *Rec. Indian Mus.* XLVI, p. 72 (foot-note).

DESCRIPTION OF THE GENUS

Body moderately elongated and naked. Head depressed or compressed and ventrally flattened. Eyes small, superior, nostrils with a free circular eyelid; placed above angle of mouth thus being not visible from below ventral surface; orbital rim free. Nostrils small, in two pairs, set apart before the anterior end of orbit. A pair of barbels on the anterior pair. Mouth wide and transverse, cleft of mouth not extending half-way to orbit. Jaws subequal, upper jaw slightly longer; but in some species almost equal. Lips cutaneous and surrounding gape of mouth. Teeth numerous, villiform, in an uninterrupted semilunar band across palate and on upper jaw; band on lower jaw interrupted in middle and deeply curved. Gill openings very wide extending forward for a considerable distance on the ventral surface. Gill membranes free but overlapping each other and free from the isthmus. Upper surface of head plain or rough, with a median longitudinal groove. Occipital process of varying length and concealed by skin or slightly naked. Interneural shield may or may not be present.

Four pairs of barbels, one pair of maxillary barbels always longer than head, one pair of thin nasal barbels, two pairs of mandibular barbels, one inner pair always shorter than head and another outer pair of varying length. Two dorsal fins, an anterior rayed dorsal with seven rays and a serrated spine; a posterior soft adipose dorsal fin, free behind rayed dorsal fin placed nearer tip of snout than base of caudal fin.

Pectoral fin with 8-10 rays and a pungent spine, stronger than dorsal spine. Pelvic fin with six rays and not reaching base of anal fin. Paired fins almost horizontally inserted. Anal papilla may or may not be present. Anal fin with less than 15 rays, short and not reaching base of caudal fin. Caudal fin bilobed or forked, lobes being unequal and sometimes prolonged into filamentous extensions.

Alimentary canal.—Stomach usually spacious, with the oesophagus united at the right side of stomach; intestines coiled.

Air-bladder.—Of varying shapes, placed immediately below the alimentary canal, anteriorly connected but posteriorly free. Internally a longitudinal complete or incomplete septum present.

KEY TO THE SUB-GENERA OF *Mystus* SCOPOLI

- | | |
|---|------------------------------------|
| I. Interneural shield in between occipital process and basal bone of dorsal fin absent.. .. | <i>Mystus</i> Scopoli, (p. 49). |
| II. Interneural shield present. | <i>Osteobagrus</i> , nov, (p. 67). |

Sub-genus **Mystus** Scopoli.

Geno-type.—*Silurus pelusius* Solander=*Mystus pelusius* (Solander) : *Type-locality* :—River Coic, Aleppo (Syria); type in British Museum, Natural History, London.

Diagnosis.—Interneural shield in between basal bone of dorsal fin and occipital process absent. Posterior extremity of air bladder not elongated; an incomplete longitudinal septum present; the lateral chambers communicate with each other by an anterior opening.

Distribution.—Asia, from Mesopotamia, Syria in North west to Ceylon and the Indo-Australian Archipelago in the south; Yangtzeikiang basin in the North East and Malabar coast of India in the west.

KEY TO THE SPECIES OF THE SUB-GENUS *Mystus*.


- | | |
|---|---------------------------|
| A. Occipital process 1.5 to 2 times as long as wide at base | 1 |
| B. Occipital process 3—5 times as long as wide at base .. | 5 |
| 1. Occipital process reaching basal bone of dorsal fin .. | 2 |
| Occipital process not reaching basal bone of dorsal fin .. | 4 |
| 2. Diameter of eyes contained 3.8 to 4.3 times in length of head, 1.25 to 1.8 times apart | <i>pulcher</i> , (p. 52). |
| Diameter of eyes contained more than 4.3 times in length of head .. | 3 |

3. Diameter of eyes 5 times in length of head and not more than 2 times apart. (Caudal peduncle normally high) *bleekeri*, (p. 50).
 Diameter of eyes 7 times in length of head; 2 to 2.25 times apart. (Caudal peduncle narrow and whip like) *vittatus horai*, sub-sp. nov, (p. 56).
4. Interorbital width 2 times the diameter of eyes. (Occipital region of head not covered with skin but rough and rugose) *gilio*, (p. 62).
 Interorbital width 1.5 to 1.75 times the diameter of eyes. (Occipital region of head covered with thick skin; occipital process not visible) *malabaricus*, (p. 64).
5. Maxillary barbels extend beyond Pelvic fins 6
 Maxillary barbels do not extend beyond Pelvic fins .. 9
6. Occipital process 3 times as long as wide at base. (Interorbital width 1 diameter of eyes. Maxillary barbels reaching just base of caudal fin) .. *oculatus*, (p. 60).
 Occipital process more than 3 times as long as wide at base 7
7. Occipital process 3.5 to 4 times as long as wide at base. (Interorbital width 1 to 1.5 diameter of eyes. Maxillary barbels extend even beyond caudal fin's base) *cavasius*, (p. 52).
 Occipital process 4 to 4.5 times as long as wide at base 8
8. Interorbital width 1 to 2 diameter of eyes. (Punctated marks along lateral line may be present) .. *montanus*, (p. 62).
 Interorbital width 2.5 to 3 diameter of eyes. (Punctated marks absent) *menoda*, (p. 64).
9. Median longitudinal groove on head reaching base of occipital process. (Interorbital width 1 to 1.5 diameter of eyes. Height of body not more than 5 times in total length. Pectoral spine with 8 to 10 teeth) *tengara*, (p. 58).
 Median longitudinal groove on head not reaching base of occipital process 10
10. Interorbital width 1.5 to 2 diameter of eyes. (Pectoral spine with 16 teeth. Height of body more than 5 times in total length.) *vittatus*, (p. 54).
 Height of body 5.25 to 5.5 times in total length 11
11. No punctated marks along the lateral line. Diameter of eyes 5 times in length of head *armatus*, (p. 59).
 Height of body 7 to 9 times in total length. Punctated marks along the lateral line .. *punctatus*, (p. 66).

Mystus (Mystus) bleekeri (Day).

1846. *Bagrus keletius*, Bleeker, *Nat. & Geneesk. Arch. Ned. Ind.* III (2), p. 135. (Type-locality.—Bengal).
1853. *Bagrus keletius*, Bleeker, *Verh. Bat. Gen.* XXV, p. 115.
1864. *Macrones keletius*, Günther, *Cat. Fish. Brit. Mus.* V, p. 84.
1877. *Macrones bleekeri*, Day, *Fish. India*, p. 451, pl. ci, fig. 1.
1889. *Macrones bleekeri*, Day, *Fauna Brit. India Fish I*, p. 162.
1890. *Macrones bleekeri*, Vinciguerra, *Ann. Mus. Stor. nat. Genova IX*, part. 2, p. 219.
1903. *Macrones bleekeri*, Volz, *Zool. Anz.* XXVI, p. 556.

1904. *Macrones bleekeri*, Regan, *Ann. Mag. nat. Hist.* (7) XIV, p. 194.
 1910. *Macrones bleekeri*, var. *burmanicus*, Jenkins, *Rec. Indian Mus.* V, p. 138.
 1911. *Macrones bleekeri*, Chaudhuri, *ibid* VI, p. 22.
 1921. *Macrones bleekeri*, Hora, *ibid* XXII, p. 179.
 1929. *Aoria bleekeri*, Prashad and Mukerji, *ibid* XXXI, p. 168.
 1934. *Aoria bleekeri*, *Mystus bleekeri*, Hora and Mukerji, *ibid* XXXVI, pp. 124, 125.
 1935. *Mystus bleekeri*, Hora and Mukerji, *ibid* XXXVII, p. 385.



TEXT-FIG. 1.—*Mystus (Mystus) bleekeri* (Day).

a. Dorsal view of head and anterior portion of body : $\times 1\frac{3}{8}$; b. dentition : $\times 4\frac{1}{5}$;
 c. alimentary canal : $\times 3\frac{1}{2}$; d. air-bladder : $\times 3\frac{1}{5}$;

B. X ; D. 1/7 ; P. 1/9-10 ; V. 6 ; A. 9-10/(3/6-7) ; C. 17.

Length of head 4 to 5 times, width of head equalling length of head behind nostrils. Height of body 4.5 to 5 times in total length. Eyes

5 diameters in length of head, 1.25 to 1.5 from end of snout and 2 apart. Median groove on head reaching base of occipital process. Maxillary barbels reach anal fin or beyond.

Distribution.—Sind, Jumna, upper waters of Ganges, Assam, Burma and Sumatra.

Remarks.—This species is closely allied to *cavasius* from which it could be distinguished by the shorter maxillary barbels. Day includes *Bagrus tengara* Blyth with a query under the synonymy of this species, which Hora & Law (1941)¹ have merged under *Batasio* Blyth.

Mystus (Mystus) pulcher (Chaudhuri).

1911. *Macrones pulcher*, Chaudhuri, *Rec. Indian Mus.*, VI, pp. 22-20, pl. i, fig. 4: (*Type-locality.*—Bhamo, close to Yunnan border).
 1929. *Aoria pulcher*, Prashad & Mukerji, *ibid* XXXI, p. 180.
 1943. *Aoria pulcher*, Nichols, *Freshw. Fish China in Nat. Hist. Cent. Asia IX*, p. 37.

B. VI : D. 1/7-0 : P. 1/9 : V. 6 : A. 12 (2/10) : C. 17.

Length of head 3.3 to 3.6 times, height of body 3.8 times in total length. Eyes 3.8 to 4.3 diameters in length of head, 1.2 to 1.4 from end of snout and 1.25 to 1.8 apart. Median groove on head does not reach base of occipital process. Maxillary barbels reach beyond anal fin. Dorsal spine with 8 teeth; pectoral spine with 10 to 12 teeth. A series of minute openings along lateral line present.

Distribution.—Bhamo, close to Yunnan border; Indawgyi lake, Myitkyana district, Burma.

Remarks.—This species is closely allied to *M. (Mystus) bleekeri*.


Mystus (Mystus) cavasius (Hamilton)

1822. *Pimelodus cavasius*, Hamilton, *Fish. Ganges* pp. 203, 379, pl. xi. fig. 67. (*Type-locality.*—Gangetic Province).
 1877. *Macrones cavasius*, Day, *Fish. India*, p. 447; pl. c, fig. 1.
 1889. *Macrones cavasius*, Day, *Fauna Brit. India Fish I*, p. 155.
 1890. *Macrones cavasius*, Vinciguerra, *Ann. Mus. Stor. nat. Genova. IX*, part 2, p. 218.
 1909. *Macrones cavasius*, Jenkins, *Rec. Indian Mus.* III, p. 287.
 1910. *Macrones cavasius*, Jenkins, *ibid*, V, pp. 137, 140.
 1911. *Macrones cavasius*, Chaudhuri, *ibid*, VI, p. 20.
 1916. *Macrones cavasius*, Chaudhuri, *Mem. Indian Mus.* V, p. 434.
 1916. *Macrones cavasius*, Raj, *Rec. Indian Mus.* XII, p. 264.
 1921. *Macrones cavasius*, Hora, *ibid* XXII, p. 179.
 1929. *Aoria cavasius*, Prashad & Mukerji, *Rec. Indian Mus.* XXXI, pp. 162, 179.

¹ Hora, S. L. & Law, N. C., *Rec. Indian Mus.*, XLIII, p. 36 (1941).

1934. *Mystus cavasius*, Smith, *J. nat. Hist. Soc. Siam.* IX, p. 294.

1935. *Macrones cavasius*, White, *Bull. geol. Surv. Nigeria* part 14, fig. 14. p. 54.



TEXT FIG. 2.—*Mystus (Mystus) Cavasius* (Hamilton).

a. dorsal view of head and anterior portion of body : $\times 1\frac{1}{2}$; b. Dentition : $\times 4$
 c. alimentary canal : $\times 2$; d. air-bladder : $\times 1\frac{1}{2}$.

1935. *Mystus rhegma*, Fowler, *Proc. Acad. nat. Sci. Philad.* LXXXVII, fig. 27, p. 102.

1936. *Mystus cavasius*, Hora, *Rec. Indian Mus.* XXXVIII, p. 1.

1937. *Mystus cavasius*, Hora, *ibid* XXXIX, pp. 17, 19.

1938. *Mystus cavasius*, Hora, *ibid* XI, p. 178.

1938. *Mystus cavasius*, Misra, *ibid*, XI, p. 256.

1938. *Mystus cavasius*, Hora & Misra, *J. Bombay nat. Hist. Soc.* XL, p. 23.

1940. *Mystus cavasius*, Hora, *Rec. Indian Mus.* XLII, pp. 366, 368, 370.

1941. *Mystus cavasius*, Hora & Nair, *ibid* XLIII, p. 369.

1941. *Mystus cavasius*, Hora & Law, *Rec. Indian Mus.*, XLIII, pp. 238, 254.
 1942. *Mystus cavasius*, Hora, *ibid* XLIV, p. 197.
 1942. *Mystus cavasius*, Hora & Misra, *J. Bombay nat. Hist. Soc.* XLII, p. 221.
 1945. *Mystus cavasius*, Smith, *Bull. U. S. nat. Mus.* number 188, p. 389.
 1948. *Mystus cavasius*, Hora, *Rec. Indian Mus.* XLVI, pp. 65, 72.
 1949. *Mystus cavasius*, Hora, *J. Zool. Soc. India* I (1), p. 2.

B. VI : D. 1/7 : P. 1/8 : V. 6 : A. 11-13 (4/7-9) : C. 16.

Length of head 4.5 to 6.5 times, height of body 5.5 to 6 times, width of head 6.5 to 6.75 times in total length. Eyes 3.5 to 4 diameters in length of head, 1 to 2 from end of snout and 1 to 1.5 apart. Median groove on head reaches base of occipital process. Maxillary barbels extend beyond base of caudal fin. Pectoral spine with 11-12 teeth.

Distribution.—Sind, Punjab, North West Frontier Provinces, Delhi, U. P., Bihar and Bengal to Assam and Burma. It extends up to Chilka Lakes and Madras in the south.

Remarks.—2 specimens from a mixed lot, from River Jumna, collected by Malaria Medical Officer are remarkable in that the union of the gill membranes over the isthmus is more rounded instead of being very sharp. This is considered to be an artifact and within the range of variation of this species.

***Mystus (Mystus) vittatus* (Bloch).**


1797. *Silurus vittatus*, Bloch, *Ichthyol. Hist. nat.* XI, p.40, pl. cccclxxi, fig. 2.
 (Type-locality.—Tranquebar, S. India).
 1822. *Pimelodus tengara*, Hamilton, *Fish. Ganges* pp. 183, 377, pl. iii, fig. 61.
 1839. *Bagrus tengara*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 414.
 1849. *Bagrus vittatus*, Jerdon, *Madras J. Lit. Sci.* XV, part, 2, p. 338.
 1853. *Bagrus vittatus*, Bleeker, *Verh. Bat. Gen.* XXV, p. 56, 114.
 1868. *Macrones tengara*, Peters, *Mber. K. Preuss. Akad. Wiss.* p. 271.
 1877. *Macrones vittatus*, Day, *Fish. India* p. 448, pls. xexiii & xcxix, figs. 3 & 4.
 1883. *Macrones vittatus*, Vinciguerra, *Ann. Mus. Stor. nat. Genova* XVIII, p. 657.
 1889. *Macrones vittatus*, Day, *Fauna. Brit. India* Fish I, p. 157.
 1890. *Macrones vittatus*, Vinciguerra, *Ann. Mus. Stor. nat. Genova* IX, part. 2, p. 229.
 1909. *Macrones vittatus*, Jenkins, *Rec. Indian Mus.* III, p. 289.
 1910. *Macrones vittatus*, Jenkins, *ibid* V, p. 135.
 1913. *Macrones vittatus*, Chaudhuri, *ibid* VIII, p. 255.
 1916. *Macrones vittatus*, Raj, *ibid* XII, p. 250.
 1916. *Macrones vittatus*, Chaudhuri, *Mem. Indian Mus.* V, p. 435.
 1921. *Macrones vittatus*, Hora, *Rec. Indian Mus.* XXII, p. 173.
 1923. *Macrones vittatus*, Hora, *Mem. Indian Mus.* V, p. 765.
 1929. *Macrones vittatus*, Pillay, *J. Bombay nat. Hist. Soc.* XXXIII, p. 358.
 1932. *Aoria vittatus*, Deraniyagala, *Spolia zeylan.* XVI, p. 283.
 1936. *Mystus vittatus*, John, *J. Bombay nat. Hist. Soc.* XXXVIII, pp. 706, 709.
 1937. *Mystus vittatus*, *Mystus atrifasciatus*, Fowler, *Proc. Acad. nat. Sci. Philad.* LXXXIX, p. 146.
 1937. *Mystus vittatus*, hora, *Rec. Indian Mus.* XXXIX, p. 19.
 1938. *Mystus vittatus*, Misra, *ibid* XL, p. 256.
 1940. *Mystus vittatus*, Shaw & Shebbeare, *J. roy. Asiat. Soc. Beng.* III, pp. 93, figs. 1, 93.

1940. *Mystus vittatus*, Hora, *Rec. Indian Mus.* XLII, pp. 366, 368.
 1941. *Mystus vittatus*, Hora & Gupta, *J. roy. Asiat. Soc. Beng.* VI, 80.
 1941. *Mystus vittatus*, Hora & Nair, *Rec. Indian Mus.* XLIII, pp. 366, 367.
 1941. *Mystus vittatus*, Hora & Law, *ibid* XLIII, p. 238.
 1942. *Mystus vittatus*, Hora, *ibid* XLIV, p. 197.
 1945. *Mystus vittatus*, Smith, *Bull. U. S. nat. Mus.* 188, p. 385, fig. 85.
 1948. *Mystus vittatus*, Hora, *Rec. Indian Mus.* XLVI, pp. 65, 70.

Length of head 4.5 to 5 times, height of body 4.5 to 5 times in total length, eyes 4 to 5 diameters in length of head, 1.5 to 2 from end of snout and apart. Median groove on head shallow and not reaching base of occipital process. Maxillary barbels reach pelvic fin. Pectoral spine with 15 to 16 teeth.

Distribution.—Throughout India, Burma, Siam and Ceylon.

Remarks.—There has been lot of confusion regarding the identity



TEXT-FIG. 3.—*Mystus (Mystus) vittatus* (Bloch).

a. dorsal view of head and anterior portion of body : $\times 1\frac{1}{2}$; b. dentition ; $\times 5\frac{1}{2}$;
 c. alimentary canal : $\times 4$; d. air-bladder : $\times 2\frac{1}{2}$.

of *vittatus* and *tengara* since the time of Day and Günther. Day (1865, pp. 190-191), describing *Hypselobagrus tengara* observed as below.—

“ I have with some hesitation considered this fish to be *Hamilton Buchanan's* “ *Pimelodns tengara* ”, but it is without doubt *Dr. Günther's* “ *Macrones tengara* ” and as he has

probably described his specimen from *Hamilton Buchanan's* typical one, I conclude some mistake may have arisen in the "Fishes of the Ganges", especially as it is there asserted that the fish is small and inhabits ponds, that its dorsal spine is smooth, and that its maxillary cirri reach to the end of the caudal fin. I should remark that this can hardly be identical as *Dr. Günther* suggests, with *Ham. Buchanan's* "*Pimelodus batasius*", pl. 23, f. 60, the figure of which is incorrect, for in the description, pp. 179, 377 it is expressly stated that it has eight cirri *shorter* than head, and only grows to about three inches in length. Again it can scarcely be the "*Pimelodus carcio*" of the same author, which is a pond fish of about three inches in length, from North Bengal, in which the maxillary cirrus does not extend to the posterior extremity of the first dorsal fin, whilst the dorsal spine is said to have a strong prickle barbed behind, & c. & c."

Hora (1929)¹, after examining Hamilton's manuscript drawings, remarked that "Day's presumption of Buchanan's published figure of *P. batasio* being a drawing of *P. carcio* is correct"

Smith (*op. cit.*), on the basis of material in the British Museum and as well as on the basis of an examination of large material from Siam, concluded that the "true *tengara* does not occur in Siam"

Also it is of interest to observe here that *tengara* does not occur in South India, but only this widely distributed *vittatus* has been recorded even from Ceylon.

***Mystus (Mystus) vittatus horai*, sub-sp. nov.**

Material.—4 specimens from River Indus, Kalabagh, collected by Dr. S. L. Hora on 25-7-23.

B. X. : D. I/7 : P. I/8 : V-6 : A. 11 (3/8) : C-17.


Head depressed, length of head 3 to 3.75 times, height of body about 5 times in total length or 3 and about 4 times in standard length respectively. Dorsal profile arched with a steep rise from tip of snout to base of occiput. Eyes round, placed laterally above angle of mouth, about 7 diameters in length of head, about 1.5 from end of snout and about 2.25 apart. Jaws equal; snout obtuse. Cleft of mouth not extending half way to orbit. Greatest width of head equals length of head behind angle of mouth. Teeth villiform and in bands. In 2 specimens teeth on palate interrupted whilst in other two continuous, but in a very narrow band; 3 to 4 rows on upper jaw and about 5 rows, interrupted on lower jaw. Upper surface of head rough. Median longitudinal groove on head does not reach base of occiput. Occipital process reaches basal bone of dorsal fin and is two times as long as broad at base. Maxillary barbels reaching pelvic fin; nasal barbels nearly as long as head, inner mandibular barbels short and outer pair extending to $\frac{3}{4}$ th of pectoral fin.

Dorsal spine small with 5 to 6 teeth on upper half equalling half the length of head. Adipose dorsal fin commences immediately after rayed fin with no intervening space in between them; its base about 2.25 times the length of base of anal fin. Pectoral fin laterally placed and equalling twice the width of gape of mouth. Pectoral spine strong, internally serrated with about 12 strong teeth. Immediately after the

¹ Hora, S. L. *Mem. Indian Mus.* IX, p. 182, (1929).

adipose fin and anal fin, the caudal peduncle is very much constricted giving the posterior end a tapering appearance. Longest ray of anal fin 1.75 to 2 times the length of base of anal fin. No anal papilla. Caudal fin bilobed. A faint black shoulder mark present. No parallel stripes along sides. Lateral line complete. Colour in spirit brownish yellow on dorsal surface with dull grey beneath.

Type-specimens.—4 specimens collected from River Indus, Kalabagh by Dr. S. L. Hora after whom the sub-species is named. Types deposited in the Zoological Survey of India, Calcutta.



TEXT-FIG. 4.—*Mystus (Mystus) vittatus horai*, sub-sp., nov.

a. dentition : $\times 7\frac{1}{2}$; b. alimentary canal : $\times 2\frac{1}{2}$; c. air-bladder : $\times 3$.

The new sub-species is very closely allied to *Mystus (Mystus) vittatus* from which it differs in its dentition; a constricted caudal peduncle; an interrupted median longitudinal groove on dorsal surface of head; eyes contained 7 diameters in length of head (*versus* 4 to 5 times in

vittatus); number of teeth on the pectoral spine (12 *versus* 15-16 in *vittatus*) and number of teeth on the dorsal spine (5 to 6 *versus* 3 to 5 in *vittatus*).

Measurements in Millimeters

Total length	106.0	96.0	82.0	76.0
Standard length	84.5	78.0	64.2	59.5
Length of head	28.5	28.5	22.8	21.0
Height of head at occiput		17.5	16.5	16.8	12.0
Width of head	19.0	18.2	15.0	12.8
Diameter of eyes		5.0	4.0	3.8	3.2
Length of snout		7.5	6.5	5.5	5.2
Interorbital width		9.2	9.0	6.2	6.0
Depth of body		21.0	20.0	16.0	15.5
Length of caudal peduncle		5.2	3.0	2.0	2.0
Least height of caudal peduncle		11.0	10.0	7.0	7.0
Longest ray of dorsal fin	..	13.8	8.0	9.0	10.0
Length of pectoral fin		15.5	17.0	13.0	12.2
Length of pelvic fin		14.0	13.2	11.0	9.0
Longest ray of anal fin		16.5	15.0	13.2	11.5
Length of base of anal fin		10.0	9.0	8.2	6.5
Length of base of adipose fin		22.5	21.0	18.8	14.5

***Mystus (Mystus) tengara* (Hamilton).**

1822. *Pimelodus carcio*, Hamilton, *Fish Ganges*, 377, pl. clxxxi, pl. xxiii, fig. 60 (written erroneously below the figure as *P. batasius*; *Type locality*: northern parts of Bengal).
1877. *Macrones tengara*, Day, *Fish India*, p. 447, pl. ci, fig. 5.
1889. *Macrones tengara*, Day, *Fauna Brit. India Fish I*, p. 156.
1913. *Macrones tengara*, Chaudhuri, *Rec. Indian Mus.*, VIII, p. 255.
1929. *Pimelodus carcio*, Hora, *Mem. Indian Mus.*, IX, p. 187.
1934. *Mystus tengara*, Smith, *J. nat. Hist. Soc. Siam IX*, part, 3, pp. 294-295.
1936. *Mystus tengara*, Prashad & Hora, *Rec. Malaria Surv. India*, VI, p. 639, pl. ix, fig. 2.
1940. *Mystus tengara*, Hora, *Rec. Indian Mus.*, XLII, p. 367.
1948. *Mystus vittatus*, Hora, *ibid XLVI*, p. 70, number 32.

B. X. : D. I/7 : P. I/8 : V-6 : A. 11-13(2-3/9-10) : C. 19.

Length of head 4 to 4.5 times, height of body 4.5 to 5.5 times in total length. Eyes 4 to 4.5 diameters in length of head, 1 to 1.5 from end of snout and 1.5 to 2 apart. Median groove on head reaches base of occipital process. Maxillary barbels extend to base of pelvic fin. Dorsal spine with 3 to 4 teeth; pectoral spine with 8-10 teeth.

Distribution.—Northern India up to Assam in the east and Madhya Pradesh in the south.

Remarks.—This species is allied to *M. (Mystus) vittatus* from which it is distinguished by the anteriorly serrated teeth in the dorsal spine; 8 to 10 teeth in the pectoral spine; median groove reaching base of occipital process and the length of caudal fin contained 4·3 times in total length (*versus* 5·5 times in *M. (Mystus) vittatus*).


***Mystus (Mystus) armatus* (Day).**

1865. *Hypselobagrus armatus*, Day, *Proc. zool. Soc. London* p. 289; (*Type-locality*:—Cochin, Malabar); *Fish Malabar*, p. 187.

1877. *Macrones armatus*, Day, *Fish India*, p. 450, pl. ci, fig. 3.

1889. *Macrones armatus*, Day, *Fauna British India Fish I*, p. 161.

1931. *Aoria armatus*, Hora, *Rec. Indian Mus.*, XXXII, p. 1.



TEXT-FIG. 5.—*Mystus (Mystus) armatus* (Day).

a. dorsal view of head and anterior portion of body: $\times 2\frac{2}{3}$; b. dentition: $\times 5\frac{1}{2}$; c. alimentary canal: $\times 1\frac{1}{2}$; d. air-bladder: $\times 1\frac{1}{2}$.

B. X.: D. I/7 : P. I/9 : V-6 : A/11(3/8) ; C. 17.

Length of head 5 to 5·5 times, height of body 5·25 to 5·5 times in total length. Eyes 5 diameters in length of head, 1·25 from end of snout

and 1.5 to 2 apart. Median groove on head does not reach base of occipital process. Maxillary barbels extend to base of pelvic fin. Pectoral spine with 10-14 teeth.

Distribution :—Malabar and Wynaad range of hills ; ? Burma.

Remarks.—Hora (*op. cit*) recorded without any comment a single specimen (F.11150/1) from Kyenchong river in Cinchona Reserve, 10 miles off La-poke-chong, Mergui district, Lower Burma and collected by K. Biswas. The specimen on thorough examination proves to belong to this species as has been rightly registered under this name. Subsequent literature reveals no further evidence of its occurrence in Burma. Although it is too presumptuous to state definitely that the species occurs in Burma also, it is worthwhile recording this remarkable zoogeographic locality record. A further collection may settle this problem. The specimen measures 202.5 mm. in total length.

This species is closely allied to *M. (Mystus) malabaricus* from which it differs in its darker colouration, an arched dorsal profile, a longer occipital process and in the denticulation of the pectoral spine (10-14 teeth *versus* 8-10).

***Mystus (Mystus) keletius* (Valenciennes).**

1839. *Bagrus keletius*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 411 (*Type locality* :—Pondicherry, S. India).
 1877. *Macrones keletius*, Day, *Fish India*, p. 449, pl. xviii, fig. 5 ; (number of figure is stated as 3 in the text erroneously).
 1889. *Macrones keletius*, Day, *Fauna Brit. India Fish I*, p. 160.
 1916. *Macrones keletius*, Raj, *Rec. Indian Mus.*, XII, p. 265.
 1932. *Aoria keletius*, Deraniyagala, *Spolia zeylan.* XVI, p. 284.
 1942. *Mystus keletius*, Hora, *Rec. Indian Mus.*, XLIV, p. 197.

B. X. ; D. I/7 ; P. I/10 ; V. 6 ; A. 9-10(2-3/7-8) ; C. 17

Length of head 4 to 4.5 times, height of body about 6 times in total length. Eyes 4.75 to 5 diameters in length of head, 1.5 to 1.75 from end of snout and 1.75 to 2 apart. Median groove on head indistinct extending to a little distance beyond eyes. Maxillary barbels reach nearly end of pelvic fin. Dorsal spine smooth ; pectoral spine with 12-13 teeth.

Distribution.—Madras, Pondicherry, Coimbatore, Mysore and Ceylon.

Remarks.—The species is very closely allied to *M. (Mystus) cavasius* from which it differs in its shorter maxillary barbels, in the median groove not reaching base of occipital process and in the brighter colouration. Günther¹, doubtfully included Valenciennes' *Bagrus keletius* under *Macrones keletius*. It is to be noted that Valenciennes' *keletius* is the south Indian species described above, whereas Günther's doubtful *keletius* is *M. (Mystus) bleekeri* from Hooghly.

***Mystus (Mystus) oculatus* (Valenciennes).**

1839. *Bagrus oculatus*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 424 (*Type-locality* :—Malabar).
 1877. *Macrones oculatus*, Day, *Fish. India*, p. 448, pl. XCVIII, fig. 4.
 1889. *Macrones oculatus*, Day, *Fauna Brit. India Fish I*, p. 156.
 1929. *Macrones oculatus*, Pillay, *J. Bombay nat. Hist. Soc.* XXXIII, p. 358.

¹ Günther, A. *Cat. Fish. Brit. Mus.*, V, p. 84 (1864).


1936. *Macrones oculatus*, John, *ibid* XXXVIII, pp. 706, 709.

1941. *Mytus oculatus*, Hora and Law, *Rec. Indian Mus.*, XLIII, p. 238.

1942. *Mytus oculatus*, Hora, *ibid* XLIV, p. 197.

B. X ; D. I/7 ; P. I/6 ; V. 6 ; A. 11-13(3-4/8-9) ; C. 15.

Length of head 4 to 5 times, height of body 4.5 to 5 times in total length. Eyes 3.75 to 4 diameters in length of head, 1 to 1.5 from end of snout and about 1 apart. Width of head nearly twice the length of snout. Median groove on head interrupted in middle, reaching base of occipital process. Maxillary barbels reach base of caudal fin. Dorsal spine with 5 to 6 teeth ; pectoral spine with 10-13 teeth.



TEXT-FIG. 6.—*Mystus (Mystus) oculatus* (Val.)

a. dorsal view of head and anterior portion of body : $\times 1\frac{1}{2}$; b. alimentary canal. $\times 3\frac{3}{4}$; c. air-bladder : $\times 4\frac{1}{2}$.

Distribution.—Malabar coast and Coimbatore district.

Remarks.—The species is very closely allied to *M. (Mystus) montanus* from which it is distinguished by the larger eyes, and denticulation on the dorsal spine (2-3 teeth in *montanus* versus 5 to 6).

***Mystus (Mystus) montanus* (Jerdon).**

1849. *Bagrus montanus*, Jerdon, *Madras J. Lit. Sci.* XV, part 2, p. 337 (*Type-locality*:—Manantoddy, Wynaad).
 1877. *Macrones montanus*, Day, *Fish India*, p. 449, pl. ci, fig. 4.
 1889. *Macrones montanus*, Day, *Fauna Brit. India Fish*, I, p. 159.
 1929. *Macrones montanus*, Pillay, *J. Bombay nat. Hist. Soc.* XXXIII, p. 358.
 1936. *Macrones montanus*, John, *ibid* XXXVIII, pp. 706, 709.
 1937. *Mystus montanus*, Hora, *Rec. Indian Mus.* XXXIX, p. 19.
 1941. *Mystus montanus*, Hora & Law, *ibid*, XLIII, pp. 254, 255.
 1942. *Mystus montanus*, Hora, *ibid* XLIV, p. 197.

B. X : D. I/7 : P. I/6 : V 6 : A. 12(3/9) : C. 19.

Length of head 4.5 times, height of body 5 to 6 times in total length. Width of head 2.5 times the length of snout. Eyes oval, 4 to 6 diameters in length of head, 1 to 2 from end of snout and 1.5 to 2 apart. Median groove on head extends to midway between eye and base of occiput. Maxillary barbels extend beyond hind edge of eyes. Dorsal spine with 2 to 3 teeth; pectoral spine with 7-8 teeth.

Distribution.—Wynaad range of hills, Manantoddy. The range is extended by 3 specimens from Hoshangabad district (Madhya Pradesh).

Remarks.—This species is closely allied to *M. (Mystus) malabaricus* from which it differs in having a more steep dorsal profile.

***Mystus (Mystus) montanus* var. *dibrugarensis* (Chaudhuri).**

1913. *Macrones montanus* var. *dibrugarensis*, Chaudhuri, *Rec. Indian Mus.*, VIII, p. 254, pl. ix, figs. 2, 2a, 2b, (*Type-locality*:—Dibrugarh, Assam).

B. X. : D. I/7 : P. I/6 : A. 9(2/7) : C. 20.

This variety differs from the *forma typica* in having the length of head 3.75 times in total length versus (4.5 times in *montanus*); height of body 4.2 times in total length (versus 5 to 6 times); dorsal spine smooth (versus 2 to 3 teeth); and pectoral spine with 12 teeth (versus 7-8 teeth).

Not so far recorded from any other locality excepting Dibrugarh Assam. Type F.7795/1, total length 68 mm., very much damaged.


***Mystus (Mystus) gulio* (Hamilton).**

1822. *Pimelodus gulio*, Hamilton, *Fish Ganges*, pp. 201, 379, pl. xxiii, fig. 66. (*Type-locality*:—Gangetic estuaries, upper parts).
 1877. *Macrones gulio*, Day, *Fish India*, p. 445, pl. xcix, fig. 2.
 1889. *Macrones gulio*, Day, *Fauna Brit. India Fish* I, p. 151, fig. 64.
 1890. *Macrones gulio*, Vinciguerra, *Ann. Mus. Stor. nat. Genova*, IX, part 2, p. 229.
 1904. *Macrones gulio*, Volz, *Zool. Jb. Abt. I*, XIX, p. 387.
 1913. *Macrones gulio*, Weber & Beaufort, *Fish. Indo-Austral. Archipel.* II, p. 344.
 1916. *Macrones gulio*, Chaudhuri, *Mem. Indian Mus.* V, p. 434.
 1917. *Macrones gulio*, Kemp, *Rec. Indian Mus.*, XIII, p. 234.
 1923. *Macrones gulio*, Hora, *J. nat. Hist. Soc. Siam* VI, p. 171.
 1929. *Aoria gulio*, Prashad & Mukerji, *Rec. Indian Mus.*, XXXI, pp. 162, 179.
 1932. *Aoria gulio*, Deraniyagala, *Spolia zeylan.* XVI, p. 285.
 1933. *Aoria gulio*, Hora, *Curr. Sci.* I, p. 382.

1934. *Mystus gulio*, Smith, *J. nat. Hist. Soc. Siam* IX, p. 294.
 1936. *Macrones gulio*, John, *J. Bombay nat. Hist. Soc.* XXXVIII, pp. 706, 709.
 1937. *Mystus gulio*, Herre & Myers, *Bull. Raffles Mus.* XIII, p. 68.
 1939. *Mystus gulio*, Herre, *Rec. Indian Mus.*, XLI, p. 111.
 1941. *Mystus gulio*, Hora & Gupta, *Bull. Raffles Mus.* XVII, p. 22.
 1941. *Mystus gulio*, Hora & Law, *Rec. Indian Mus.*, XLIII, pp. 255, 238.
 1942. *Mystus gulio*, Hora & Misra, *J. Bombay nat. Hist. Soc.* XLIII, p. 222.
 1945. *Mystus gulio*, Smith, *Bull. U. S. nat. Mus.* (188), p. 384.
 1948. *Mystus gulio*, Hora, *Rec. Indian Mus.*, XLVI, pp. 65, 72.

B. IX : D. I/7 : P. I/8-9 : V. 6 : A. 12-15 (4/9-11) : C. 17.

Length of head about 4.5 times, height of body 5 to 5.75 times in total length. Eyes 5 to 6 diameters in length of head, 1.5 to 2 from end of snout and 2 apart. Upper surface of head rough and granulated. Median groove on head shallow, lanceolate, extending to opposite hind edge of orbit. Maxillary barbels reach middle or nearly end of pelvic fin.




TEXT-FIG. 7.—*Mystus (Mystus) gulio* (Hamilton).

a. dorsal view of head and anterior of body : $\times 3$; b. dentition : $\times 1$; c. alimentary canal : $\times 2$; d. air-bladder : $\times 1\frac{1}{2}$.

Distribution.—Seas, estuaries and tidal waters from Sind, India, Burma to the Malay Archipelago and Ceylon as also Sumatra and Siam.

Mystus (Mystus) malabaricus (Jerdon).

1849. *Bagrus malabaricus*, Jerdon, *Madras J. Lit. Sci.* XV, p. 338 (*Type-locality* :—Mountain streams in Malabar).
1869. *Hara malabarica*, Day, *Proc. zool. Soc. London* p. 524.
1877. *Macrones malabaricus*, Day, *Fish India*, p. 450, pl. ci. fig. 2.
1889. *Macrones malabaricus*, Day, *Fauna Brit. India Fish I*, p. 160.
1929. *Macrones malabaricus*, Pillay, *J. Bombay nat. Hist. Soc.* XXXIII, p. 359.
1936. *Macrones malabaricus*, John, *ibid.* XXXVIII, pp. 706, 709.
1937. *Macrones malabaricus*, Hora, *Rec. Indian Mus.* XXXIX, p. 19.
1941. *Mystus malabaricus*, Hora & Law, *ibid.*, XLIII, pp. 254, 255, fig. 3.
1942. *Mystus malabaricus*, Hora, *ibid.*, XLIV, p. 197.



TEXT-FIG. 8.—*Mystus (Mystus) malabaricus* (Jerdon).

a. dorsal view of head and anterior portion of body : $\times 1\frac{1}{2}$; b. dentition : $\times 5\frac{1}{2}$; c. alimentary canal : $\times 2\frac{2}{3}$; d. air-bladder : $\times 2\frac{2}{3}$.


B. X : D. I/7 : P. I/9 : V. 6 : A. 10-11 : C. 18.

Length of head 4.5 to 5 times, height of body 6 to 7 times in total length. Eyes 4.5 to 5 diameters in length of head, 1.5 to 2 from end of

snout and 1.5 to 1.75 apart. Median groove on head extends to hind edge of eye or slightly beyond. Maxillary barbels reach middle or end of pelvic fin.

Distribution.—Malabar coast of India and Wynaad hills and hill ranges of Travancore. The range is now extended by 3 specimens from Palgad, Ratnagiri district (Bombay state), and one specimen from Poona (F. 12141/1).

Remarks.—The species is closely allied to *M. (Mystus) montanus* from which it differs in the nature of the occipital process and serrations of the pectoral spine.



TEXT-FIG. 9.—*Mystus (Mystus) menoda* (Hamilton).

a. dorsal view of head and anterior portion of body : $\times 2/3$; b. dentition : $\times 2\frac{2}{3}$;
 c. alimentary canal : $\times 1\frac{1}{3}$; d. air-bladder ; $\times 1\frac{1}{3}$.

Mystus (Mystus) menoda (Hamilton).

1822. *Pimelodus menoda*, Hamilton, *Fish Ganges*, pp. 203, 379, pl. i, fig. 72
(*Type-locality*.—Kosi, Mahanada, North Bihar & Bengal).
1839. *Bagrus corsula*, Valenciennes, *Hist. Nat. Poiss.* XIV, p. 408.
1853. *Bagrus corsula*, Bleeker, *Verh. Bat. Gen.* XXV, p. 56.
1864. *Pimelodus menoda*, Günther, *Cat. Fish. Brit. Mus.* V, p. 74 (foot-note).
1877. *Macrones corsula*, Day, *Fish India*, p. 446, pl. ci, fig. 5.
1889. *Macrones corsula*, Day, *Fauna Brit. India* Fish I, p. 153.
1890. *Macrones menoda*, Vinciguerra, *Ann. Mus. Store nat., Genova* IX, p. 223.
1911. *Macrones corsula*, Chaudhuri, *Rec. Indian Mus.*, VI, p. 24.
1912. *Macrones menoda*, Chaudhuri, *ibid.*, VII, p. 211.
1929. *Pimelodus menoda*, Hora, *Mem. Indian Mus.*, IX, p. 183, 188.
1938. *Mystus menoda*, Shaw & Shebbeare, *J. roy. Asiat. Soc. Beng.* III, p. 92,
fig. 91, pl. ii, fig. 2.
1948. *Mystus menoda*, Hora, *Rec. Indian Mus.*, XLVI, pp. 65, 72.

B. X : D. I/7 : P. I/9 : V 6 : A. 11-13 (3-5/8) : C. 17

Length of head 3.75 to 4.25 times, height of body 5 to 6 times in total length. Eyes 7 to 8 diameters in length of head, 2 to 4 from end of snout and 2.5 to 3 apart. Median groove on head do not reach base of occipital process. Maxillary barbels reach base of anal fin. Dorsal spine with 2 to 3 teeth ; pectoral spine with 20-23 teeth.

Distribution.—From Orissa to Bengal, Assam and Burma.

Remarks.—Attention is invited to the note published by Chaudhuri in 1912 (*op. cit.*) which cleared the confusion regarding the name *corsula* and *menoda*.

Day (1877, p. 446), described *M. (Mystus) microphthalmus* from the Irrawady valley (Burma) as a new species and observed that "possibly all these last forms are local races of one species". The differences between *M. (Mystus) menoda* and *M. (Mystus) microphthalmus* being of a very minor nature, I have treated the latter species as a second subspecies of the former one. The two subspecies of *M. menoda* now known, could be distinguished by the following key :—

- | | | |
|--|----|---|
| 1. Caudal fin's lower lobe prolonged into a filament | .. | <i>M. (Mystus) menoda</i> var.
<i>trachacanthus</i> . |
| 2. Caudal fin's upper lobe prolonged into a filament | .. | <i>M. (Mystus) menoda</i> var.
<i>microphthalmus</i> . |

Mystus (Mystus) menoda trachacanthus (Valenciennes).

1839. *Bagrus trachacanthus*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 419 (*Type-locality*:—Bengal).
1864. *Bagrus trachacanthus*, Günther, *Cat. Fish. Brit. Mus.* V, p. 75
1912. *Macrones menoda*, var. *trachacanthus*, Chaudhuri, *Rec. Indian Mus.*, VII, p. 210.

B. X : D. I/7 : P. I/9 : A. 12 : C. 17.

Length of head 4 times, height of body about 4.4 times in total length. Eyes 4.6 diameters in length of head, 2 from end of snout and also apart. Median groove on head not reaching base of occipital

process. Maxillary barbels extending to middle of pelvic fin. Pectoral spine with 12-13 teeth. Lower lobe of caudal fin filamentous.

Distribution.—Bombay, Bengal.

***Mystus (Mystus) menoda microphthalmus* (Day).**

1877. *Macrones microphthalmus*, Day, *Fish India*, p. 446, pl. ci, fig. 4 (*Type-locality.*—Irrawaddy, Burma).
 1889. *Macrones microphthalmus*, Day, *Fauna Brit. India Fish I*, p. 154.
 1890. *Macrones microphthalmus*, Vinciguerra, *Ann. Mus. Stor. nat. Genova IX*, p. 225.

B. X : D. 1/7 : P. 1/9 : V 6 : A. 12 (3/9) : C. 17.

Length of head 3.5 to 4 times, height of body 7 to 8 times in total length. Eyes 6 to 8 diameters in length of head, 2 to 3 from end of snout and about 2.5 apart. Median groove on head reaches base of occipital process. Maxillary barbels reach anal fin or beyond. Pectoral spine with 9-10 teeth. Upper lobe of caudal fin filamentous.

Distribution.—Burma along the valley or Irrawady.

***Mystus (Mystus) punctatus* (Jerdon).**

1849. *Bagrus punctatus*, Jerdon, *Madras J. Lit. Sci.* XV, p. 339 (*Type-locality* —Cauvery River, W. Ghats).
 1877. *Macrones punctatus*, Day, *Fish India*, p. 445, pl. c, fig. 3.
 1889. *Macrones punctatus*, Day, *Fauna Brit. India Fish I*, p. 153.
 1937. *Mystus punctatus*, Hora, *Rec. Indian Mus.*, XXXIX, p. 19.
 1942. *Mystus punctatus*, Hora, *ibid.*, XLIV, p. 197.

B. XI : D. 1/7 : P. 1/7-8 : V. 6 : A. 11-13 (3-4/8-9) : C. 17.

Length of head 4.5 to 5 times, height of body 7 to 9 times in total length. Eyes 6 to 6.5 diameters in length of head, about 2.5 from end of snout and about 2.25 apart. Width of head nearly 1.5 times height of head at occiput. Median groove on head shallow, broad, reaching halfway between occipital process and eyes. Maxillary barbels extend upto pelvic fin. Pectoral spine with 15 to 16 teeth.

Distribution.—Coorg, Nilgiris, Mysore, Malabar and Travancore. The range is extended by a specimen from Bombay (Cat. 426).

Sub-genus ***Osteobagrus***, nov.

Geno-type.—*Pimelodus aor* Hamilton-*Mystus aor* (Hamilton) :
Type-locality.—Ganges river.

Diagnosis.—A distinct interneural shield between basal bone of dorsal fin and occipital process present; posterior extremity of

air-bladder elongated and pyriform ; a longitudinal complete septum in air-bladder present ; two lateral chambers of air-bladder not communicating with each other.

Day as early as 1871¹ realised the necessity of creating a subgenus for *Macrones aor* and *M. lamarrii* (Valenciennes), but did not do so either that year or in his subsequent work on the *Fishes of India*. In the earlier work, however, he stated :

“ The subgenera, or those with a separate interneural shield on the nape and those destitute of such, appear, at least sometime to denote other internal structural differences in those which I have examined. Amongst the former are *M. aor* and *M. lamarri* in which the anterior portion of the air-vessel is attached to the under surface of the bodies and expanded processes of the anterior vertebrae, but its posterior extremity is elongated and pyriform ; internally it has a longitudinal septum. On the contrary in those not having this separate shield, the posterior extremity of the air-vessel is not elongated, the longitudinal septum (as in all those of this genus which I have examined) has a communicating opening anteriorly ; and sometimes there are transverse partitions forming chambers which freely communicate with those on the same and by means of the anterior one with those on the opposite side as in *M. cavasius*, *M. tengara* *M. carcio*.”

In 1877², he stated as below :—

“ Whether *M. aor* or *M. seenghala* which have a separate interneural shield and an air-vessel with a posterior elongated extremity, should not be separated from the remainder which have no such distinct shield and no prolongation of the air-vessel, may be open to question. ”

From the above it would appear that this subgenus would include *M. seenghala* (Sykes) also besides the geno-type. *Bagrus leucophasis* Blyth should also fall under this subgenus in view of the presence in it of the above-noted characters.

Key to the species of the sub-genus *Osteobagrus*.

- | | |
|---|---|
| 1. Snout spatulate | 2. |
| Snout non-spatulate .. | 3. |
| 2. Width of gape of mouth $\frac{2}{5}$ ths to $\frac{3}{7}$ ths length of head | <i>M. (Osteobagrus) aor</i>
(Hamilton). |
| Width of gape of mouth $\frac{1}{3}$ rd length of head | <i>M. (Osteobagrus) seenghala</i>
(Sykes). |
| 3. Snout rounded. Width of gape of mouth less than $\frac{1}{3}$ rd length of head .. | <i>M. (Osteobagrus) leucophasis</i>
(Blyth). |

Mystus (Osteobagrus) aor (Hamilton).

1822. *Pimelodus aor*, Hamilton, *Fish Ganges*, pp. 205, 379, pl. xx, fig. 68
(Type-locality.—Rivers of Bengal and upper parts of Gangetic estuaries).
1877. *Macrones aor*, Day, *Fish India*, p. 444.
1886. *Macrones aor*, Lydekker, *Palaeont. indica* (10) III, p. 250, pl. xxxvi, fig. 5.
1889. *Macrones aor*, Day, *Fauna Brit. India* Fish I, p. 149.
1890. *Macrones aor*, Vinciguerra, *Ann. Mus. stor. nat. Genova* IX, part 2, p. 217.
1910. *Macrones aor*, Jenkins, *Rec. Indian Mus.*, V, p. 140.
1934. *Macrones aor*, Bhimachar, *Half-yrly. J. Mysore Univ.* VII, pp. 253-254 pl. v, fig. 14 (Skull).


¹ Day, F. *Proc. Zool. Soc. Lond.*, p. 705.

² Day, F. *Fish India*, p. 442 (1877).

1937. *Mystus aor*, Hora, *Rec. Indian Mus.*, XXXIX, p. 19.

1940. *Mystus aor*, Hora, *ibid.*, XLII, p. 368.

1948. *Mystus aor*, Hora, *ibid.*, XLVI, p. 72.



TEXT-FIG. 10.—*Mystus (Osteobagrus) aor* (Hamilton).

a. dorsal view of head and anterior portion of body : $\times 1\frac{1}{3}$; b. dentition : $\times 4$;
c. alimentary canal ; $\times 1\frac{1}{3}$; d. air-bladder ; $\times 1\frac{1}{3}$.

B. XII : D. I/7 : P. I/9-10 : V 6 : A. 12-13 (3-4/8-9) : C. 17.

Length of head 4.5 to 5 times, width of head about 8 times, height of body 6.5 to 7.5 times in total length. Eyes 4.5 to 5.5 diameters in head, 1.5 to 2 from end of snout, and same in interorbital width. Median groove on head reaches base of occipital process. Maxillary barbels reach nearly caudal fin or even beyond. Dorsal spine with 4 to 5 teeth ; pectoral spine with 20 to 25 teeth.

Distribution.—Throughout Sind, Punjab, Delhi, Assam and Burma.

Remarks.—This is closely allied to *seenghala* from which it differs in the height of body (5 to 6 times in total length *versus* 7.5 to 8 times in *seenghala*); width of gape of mouth (2/5ths to 3/7ths of length of head *versus* 1/3rd length of head).

***Mystus (Osteobagrus) seenghala* (Sykes).**

- 1839¹ (May). *Platystoma seenghala*, Sykes, *Trans. zool. Soc. Lond.* II, p. 371, pl. lxxv, fig. 2 (*Type-locality.*—Mota Mala river, Poona).
1839. *Bagrus lamarrii*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 407, pl. cxlv.
1849. *Bagrus seenghala*, Jerdon, *Madras J. Lit. Sci.* XV, part 2, p. 337.
1853. *Bagrus lamarrii*, Bleeker, *Verh. Bat. Gen.* XXV, p. 56.
1856. *Bagrus lamarrii*, Dumeril, *Mem. Acad. Sci. Paris* XXVII, part 1, p. 484 (Dumeril's genotype for genus *Macrones*).
1863. *Macrones lamarrii*, Bleeker, *Ned. Tijdschr. Dierk.* I, p. 96.
1864. *Macrones lamarrii*, Günther, *Cat. Fish Brit. Mus.* V, p. 79.
1871. *Macrones lamarrii*, Day, *Proc. zool. Soc. Lond.* p. 705.
1877. *Macrones seenghala*, Day, *Fish India*, p. 444, pl. xcix, fig. 1.
1889. *Macrones seenghala*, Day, *Fauna Brit. India Fish* I, p. 150.
1911. *Macrones seenghala*, Chaudhuri, *Rec. Indian Mus.*, VI, p. 20.
1942. *Mystus seenghala*, Hora & Misra, *J. Bombay nat. Hist. Soc.* XLIII, p. 222.
1943. *Aoria seenghala*, Nichols, *Freshw. Fish China in Nat. Hist. Cent. Asia* IX, p. 37.
1949. *Mystus seenghala*, Hora, *J. zool. Soc. India* I, p. 2.

B. XII : D. I/7 : P. I/9 : V. 6 : A. 11-12 (3/8-9) : C. 19-21.

Length of head 4 to 4.5 times, width of head 7 to 8 times, height of body 7.5 to 8 times in total length. Eyes 6 to 7 diameters in head, about 2 from end of snout and 1.5 to 2 apart. Median groove on head reaches base of occipital process. Maxillary barbels extend to middle of dorsal fin or slightly beyond. Dorsal spine with 2 to 3 teeth; pectoral spine with about 20 teeth.

Distribution.—Punjab, Delhi, U. P., Bengal and Burma. In the south it extends upto Kistna river and its termination. Recorded from Yunnan also.

Remarks.—Sykes in describing this species mentions that “this fish is remarkable for having the first ray of the ventral as well as that of the pectoral serrated posteriorly, and the first dorsal spine *not* serrated”. His drawing on plate lxxv, fig. 2 also, clearly depicts this. I have not met with any such instance of a serrated first ray of the pelvic fins in any species of the genus *Mystus*.

***Mystus (Osteobargus) leucophasis* (Blyth).**


1860. *Bagrus leucophasis*, Blyth, *J. Asiat. Soc. Beng.* XXIX, p. 148 (*Type-locality.*—Sittang and other Burmese rivers).
1864. *Macrones leucophasis*, Günther, *Cat. Fish Brit. Mus.* V, p. 78.
1877. *Macrones leucophasis*, Day, *Fish India*, p. 449, pl. c, fig. 2.
1883. *Macrones leucophasis*, Vinciguerra, *Ann. Mus. Stor. nat. Genova*, XVIII, p. 659.

¹ Although Sykes' work is often quoted as of 1841, Sherborn gives the actual date of publication as May, 1839, thus giving priority to Sykes' work over Valenciennes' which was published in January, 1840. Prior to 1877 earlier workers seem to have been unaware of this and have used *lamarrii* instead of *seenghala* for the trivial name.

1889. *Macrones leucophasis*, Day, *Fauna Brit. India Fish I*, p. 158.
 1890. *Macrones leucophasis*, Vinciguerra, *Ann. Mus. Stor. nat. Genova*, IX,
 p. 220.
 1921. *Macrones leucophasis*, Hora, *Rec. Indian Mus.* XXII, p. 179.
 1929. *Aoria leucophasis*, Prashad & Mukerji, *ibid.*, XXXI, p. 179

B. XI : D. I/7 : P. I/8-10 : V. 6 : A. 9-10 (2-3/7-8) : C. 17.

Length of head 4.5 times, width of head 5 to 5.5 times, height of body 5.5 times in total length. Eyes about 6 diameters in length of head, 2 from end of snout and about 2.5 apart. Median groove on head not reaching base of occipital process. Maxillary barbels reach anal fin. Dorsal spine serrated posteriorly.



TEXT-FIG. 11.—*Mystus (Osteobagrus) leucophasis* (Blyth).

a. dorsal view of head and anterior portion of body : $\times 1\frac{1}{2}$; b. dentition : $\times 4$
 c. alimentary canal : $\times 1\frac{1}{2}$; d. air-bladder : $\times 2$.

Distribution.—Rivers of Burma.

SPECIES NOT LISTED ABOVE.

Macrones chryseus (Day).

The systematic position of this south Indian species has been elucidated in a note published by the writer¹. Further examination of the allied *Pseudobagrus* material have proved this to be warranting a generic status. The species is not a *Mystus* but allied to *Pelteobagrus* Bleeker. Description of the new genus is being published separately.

Mystus (Mystus) peguensis (Boulenger).

1894. *Macrones peguensis*, Boulenger, *Ann. Mag. nat. Hist.* (6) XIV, p. 196.

Distribution.—Sittang river near Toungoo (Burma).

Remarks.—This species is not represented in the Z. S. I.² collection. This has not also been recorded subsequently.

Mystus (Mystus) rufescens (Vinciguerra).

1890. *Macrones rufescens*, Vinciguerra, *Ann. Mus. Stor. nat. Genova*, IX (2), p. 226.

Distribution.—Meetan (Burma).

Remarks.—This species also is not represented in Z. S. I. collection nor it has been recorded subsequently from Burma.

Mystus (Mystus) pelusius var. **colvillii** (Günther).

1794. *Silurus pelusius*, Solander in Russell's *Nat. Hist. Aleppo* II, p. 210, pl. vii, fig. 1.

1841. *Bagrus halepensis*, Heckel in Russeger's *Reisen Europa, Asien und Africa*, I, p. 1091, pl. viii, fig. 2.

1864. *Macrones aleppensis*, Günther, *Cat. Fish. Brit. Mus.* V, pp. 75, 431.

1874. *Macrones Colvilli*, Günther, *Ann. Mag. nat. Hist.* (4) XIV, p. 36, pl. viii,

1943. *Mystus halepensis colvilli*, Hora and Misra, *J. roy. Asiat. Soc. Beng.* IX, p. 8, fig. 4.

Distribution.—Bagdad, River Tigris.

Remarks.—This species is also neither represented in Z. S. I. collection nor it has been subsequently recorded after 1943.

Mystus (Mystus) pelusius (Solander).

1794. *Silurus pelusius*, Solander in Russell's *Hist. Aleppo* II, p. 210, pl. vii, fig. 1.

1864. *Bagrus aleppensis, halepensis*, Günther, *Cat. Fish. Brit. Mus.* V, pp. 75, 431.

Distribution.—River Coic, Aleppo, Syria.

Remarks.—This species is the geno-type of the genus *Mystus*. This is also not represented in Z. S. I. collection, nor it is subsequently recorded.

¹ Jayaram, K. C. *Ann. Mag. nat. Hist.* V (12), pp. 980-983 (1952).

² Z.S.I. = Zoological Survey of India, Calcutta.

SPECIES FOUND IN SIAM, MALAY PENINSULA AND ARCHIPELAGO.

Mystus (Mystus) wolffi (Bleeker).

1858. *Bagrus wolffi*, Bleeker, *Ichth. Arch. Ind. Prodr.* I, *Siluri*, p. 160.
 1913. *Macrones wolffi*, Weber & Beaufort, *Fish Indo. Austral. Archeipel.* II, p. 340.
 1945. *Mystus wolffi*, Smith, *Bull. U. S. nat. Mus.* (188), p. 383.

Distribution.—Thailand, Perak, Malacca, Borneo and Sumatra.

Remarks.—This species is not represented in Z. S. I. collection.

Mystus (Mystus) pahangensis Herre.

1940. *Mystus pahangensis*, Herre, *Bull. Raffles Mus.* (16), p. 14.

Distribution.—Pahang (Malay Peninsula). Not further Recorded.

Mystus (Mystus) nemurus (Valenciennes).

1839. *Bagrus nemurus*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 423.
 1913. *Macrones nemurus*, Weber & Beaufort, *Fish. Indo-Austral. Archipel.* II, p. 341.
 1945. *Mystus nemurus*, Smith, *Bull. U. S. nat. Mus.* (188), p. 386.

Distribution.—Thailand, Malay Archipelago.

Remarks.—This species is represented in Z. S. I. collection.

Mystus (Mystus) nigriceps (Valenciennes).

1839. *Bagrus nigriceps*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 412.
 1913. *Macrones nigriceps*, Weber & Beaufort, *Fish. Indo-Austral. Archipel.* II, p. 337.
 1945. *Mystus nigriceps*, Smith, *Bull. U. S. nat. Mus.* (188), p. 389 (merged under *M. cavasius*).

Distribution.—Thailand, Malay Peninsula, Java, Sumatra and Borneo.

Remarks.—This species is very closely allied to *M. cavasius* but distinct enough to warrant a specific rank. This species is also represented in Z. S. I. collection.

Mystus (Mystus) planiceps (Valenciennes).

1839. *Bagrus planiceps*, Valenciennes, *Hist. nat. Poiss.* XIV, p. 421.
 1913. *Macrones planiceps*, Weber & Beaufort, *Fish. Indo. Austral. Archeipel.* II, p. 342.
 1945. *Mystus planiceps*, Smith, *Bull. U. S. nat. Mus.* (188), p. 387.

Distribution.—Thailand, Java, Sumatra, Borneo and Malay Peninsula.

Remarks.—I have examined the collection in Z. S. I.

Mystus (Mystus) wyckii (Bleeker).

1858. *Bagrus wyckii*, Bleeker, *Ichth. Arch. Ind. Prodr.* I, *Siluri*, p. 156.
 1913. *Macrones wyckii*, Weber & Beaufort, *Fish. Indo. Austral. Archipel.* II, p. 343.
 1945. *Mystus wyckii*, Smith, *Bull. U. S. nat. Mus.* (188), p. 388.

Distribution.—Thailand, Perak, Malay Peninsula, Sumatra and Java.

Remarks.—This species is not represented in the Z. S. I. collection.

Mystus (Mystus) johorensis Herre.

1940. *Mystus johorensis*, Herre, *Bull. Raffles Mus.* (16), p. 13.

Distribution.—Johore (Malay Peninsula). Not subsequently recorded.

Mystus (Mystus) micracanthus (Bleeker).

1846. *Bagrus micracanthus*, Bleeker, *Nat. & Geneesk. Arch. Ned. Ind.* III (2), p. 151.

1913. *Macrones micracanthus*, Weber & Beaufort, *Fish. Indo. Austral. Archipel.* II, p. 454.

1945. *Mystus micracanthus*, Smith, *Bull. U. S. nat. Mus.* (188), p. 391.

Distribution.—Thailand, Malay Peninsula, Java, Sumatra and Borneo.

Remarks.—This species is not represented in Z. S. I. collection.

SPECIES RESTRICTED TO THE INDO-AUSTRALIAN ARCHIPELAGO.

Mystus (Mystus) baramensis (Regan).

1906. *Macrones baramensis*, Regan, *Ann. Mag. nat. Hist.* XVIII, p. 68.

1913. *Macrones baramensis*, Weber & Beaufort, *Fish. Indo. Austral. Archipel.* II, p. 338.

1937. *Mystus baramensis*, Herre & Myers, *Bull. Raffles Mus.* (13), p. 19.

Distribution.—Baram river (Borneo), Malacca.

Remarks.—This species is not present in Z. S. I. collection.

SPECIES OF CHINESE REGION.

Mystus (Mystus) argentivittata (Regan).

1905. *Macrones argentivittata*, Regan, *Rev. Suisse. Zool.* XIII, p. 390, pl. v, fig. 2.

1943. *Aoria argentivittata*, Nichols, *Freshw. Fish. China in nat. Hist. Cent. Asia* IX, p. 37.

Distribution.—China.

Mystus (Mystus) henryi (Herre).

1942. *Aoria henryi*, Herre, *Lingnan Sci. J.* XI, p. 342.

1943. *Aoria henryi*, Nichols, *Freshw. Fish. China in nat. Hist. Cent. Asia* IX, p. 37.

Distribution.—Canton, S. China.

Mystus (Mystus) cornula (Chu).

1931. *Aoria cornula*, Chu, *Biol. bull. St. John's Univ.* 76.

1943. *Aoria cornula*, Nichols, *Freshw. Fish. China in nat. Hist. Cent. Asia* IX, p. 36.

Distribution.—China. Nichols has recorded this as a doubtful Chinese species.

***Mystus (Mystus) hoi* (Pellegrin & Fang).**

1940. *Mystus hoi*, Pellegrin and Fang, *Bull. Soc. Zool. France* LXIV, p. 338.

Distribution.—Yangtse basin (China). Nichols has not recognised this species in his work.

***Mystus (Mystus) pluriradiatus* (Vaillant).**

1892. *Macrones pluriradiatus*, Vaillant, *Bull. Soc. Phil.* IV (8), p. 126.

1904. *Macrones pluriradiatus*, Vaillant, *Rech. Hist. Indo-China* III, p. 462, pl. xxiii, fig. 2.

Distribution.—Tonkin (South China).

***Mystus (Mystus) rendahli* (Pellegrin & Fang).**

1940. *Mystus rendahli*, Pellegrin & Fang, *Bull. Soc. Zool. France*, LXIV, p. 338.

Distribution.—Yangtse basin (China).

The following is the systematic position of some of the other species formerly described under this genus from China (Nichols, 1943).

<i>Aoria virgatus</i> Oshima	<i>Pseudobagrus virgatus.</i>
<i>Macrones medianlis</i> Regan	<i>Pseudobagrus medianalis.</i>
<i>Macrones pratti</i> Günther	<i>Leiocassis pratti.</i>
<i>Macrones taeniatus</i> Günther	<i>Leiocassis taeniatus.</i>
<i>Macrones tenuis</i> Günther	<i>Leiocassis tenuis.</i>
<i>Aoria macroptera</i> Chu	<i>Hemibagrus macropterus.</i>
<i>Aoria amemiyae</i> Kimura	<i>Hemibagrus amemiyae.</i>
<i>Macrones elongatus</i> Günther	<i>Hemibagrus elongatus.</i>
<i>Macrones chinensis</i> Steindachner	<i>Hemibagrus elongatus.</i>
<i>Macrones sinensis</i> Bleeker	<i>Unidentifiable.</i>
<i>Macrones ussuriensis</i> Her. Warp.	<i>Leiocassis ussuriensis.</i>

GEOGRAPHIC DISTRIBUTION.

Fishes of the genus *Mystus* are widely distributed from Asia Minor in the west to southern China in the east. They live in marshy and sluggish rivers generally, with the exception of certain peninsular species like *M. (Mystus) punctatus*, *M. (Mystus) montanus*, etc., which live in hill streams. Amongst the living species *Mystus (Osteobagrus) aor* is known¹ from the Pliocene Siwalik beds of India and *Mystus (Mystus) cavasius*² from Eocene Nigeria of Africa.

¹ Lyddekkar, R. *Palaeont. indica* (10) III, p. 250, pl. xxxvi, fig. 5 (1886).

² White, E. I. *Bull. Geol. Surv. Nigeria* (14), pp. 53-55, fig. 14 (1935).

The table of distribution of the speices given below leads to the following conclusions. Out of the 39 species and subspecies of *Mystus*, 2 are endemic in Asia Minor, 1 in East Indies, 6 in China, 3 in Malay Peninsula, 4 in Burma, 1 in Assam, 4 in Peninsular India and 1 in Northern India. Seven species are widely distributed while the rest 10 species are found in two areas together. Of the last mentioned 10 species, excepting *Mystus (Mystus) armatus*, *M. (Mystus) montanus*, *M. (Mystus) menoda* var. *trachacanthus*, rest are distributed homogeneously in adjacent areas. The above cited species are discontinuous in distribution and their pattern of distribution is interesting.

M. (Mystus) montanus has been recorded from Hoshangabad (Madhya Pradesh), far from its range of Peninsular India. Similarly *M. (Mystus) armatus* is recorded from Burma. In the case of *M. (Mystus) menoda*, another peninsular species, two geographic races are known. Thus there seems to be a particular east to west spread of these fishes which has resulted in such anomalies.

No.	Name of species.	1	2	3	4	5	6	7	8	
1.	<i>M. (Mystus) bleekeri</i> (Day)	..	—	X	X	X	—	X	—	—
2.	<i>M. (Mystus) pulcher</i> (Chaudhuri)	..	—	—	—	X	—	—	X	—
3.	<i>M. (Mystus) cavasius</i> (Hamilton)	..	X	X	X	X	—	—	—	—
4.	<i>M. (Mystus) vittatus</i> (Bloch)	..	X	X	X	X	X	—	—	—
5.	<i>M. (Mystus) vittatus horai</i> Jayaram		—	X	—	—	—	—	—	—
6.	<i>M. (Mystus) armatus</i> (Day)		X	—	—	X	—	—	—	—
7.	<i>M. (Mystus) keletius</i> (Valenciennes)	..	X	—	—	—	—	—	—	—
8.	<i>M. (Mystus) tengara</i> (Hamilton)		—	X	X	—	—	—	—	—
9.	<i>M. (Mystus) oculatus</i> (Valenciennes)	..	X	—	—	—	—	—	—	—
10.	<i>M. (Mystus) montanus</i> (Jerdon)		X	X	—	—	—	—	—	—
11.	<i>M. (Mystus) montanus</i> var. <i>dibrugarensis</i> (Chaudhuri).		—	—	X	—	—	—	—	—
12.	<i>M. (Mystus) gulio</i> (Hamilton)		X	X	—	X	X	X	—	—
13.	<i>M. (Mystus) malabaricus</i> (Jerdon)		X	—	—	—	—	—	—	—
14.	<i>M. (Mystus) menoda</i> (Hamilton)		X	X	X	X	—	—	—	—

The eight distribution regions (artificial) here recognised are :—

- 1.—Peninsular India and Ceylon ; 2.—Northern India ; 3.—Assam ; 4.—Burma ; 5.—Siam and Malay Archipelago ; 6.—East Indies ; 7.—China ; 8.—Asia Minor.

No.	Name of species.	1	2	3	4	5	6	7	8
15.	<i>M. (Mystus) menoda</i> var. <i>trachacanthus</i> (Valenciennes).	X	X	—	—	—	—	—	—
16.	<i>M. ((mystus) menoda microphthalmus</i> (Day).	—	—	—	X	—	—	—	—
17.	<i>M. (Mystus) punctatus</i> (Jerdon)	..	X	—	—	—	—	—	—
18.	<i>Mystus (Osteobagrus) aor</i> (Hamilton)	—	X	X	X	—	—	—	—
19.	<i>Mystus (Osteobagrus) seenghala</i> (Sykes)	X	X	—	X	—	X	—	—
20.	<i>Mystus (Osteobagrus) leucophasis</i> (Blyth)	—	—	—	X	—	—	—	—
21.	<i>M. (Mystus) peguensis</i> (Boulenger)	..	—	—	X	—	—	—	—
22.	<i>M. (Mystus) refescens</i> (Vinciguerra)	..	—	—	X	—	—	—	—
23.	<i>M. (Mystus) pelusius</i> (Solander)	..	—	—	—	—	—	X	—
24.	<i>M. (Mustus) pelusius</i> var. <i>colvilli</i> (Gunther)	—	—	—	—	—	—	X	—
25.	<i>M. (Mystus) wolffi</i> (Bleeker)	..	—	—	—	X	X	—	—
26.	<i>M. (Mystus) pahangensis</i> Herre	..	—	—	X	—	—	—	—
27.	<i>M. (Mystus) nemurus</i> (Valenciennes)	..	—	—	X	—	—	—	—
28.	<i>M. (Mystus) nigriceps</i> (Valenciennes)	—	—	—	X	X	—	—	—
29.	<i>M. (Mystus) planiceps</i> (Valenciennes)	—	—	—	X	X	—	—	—
30.	<i>M. (Mystus) wyckii</i> (Bleeker)	—	—	—	X	X	—	—	—
31.	<i>M. (Mystus) johorensis</i> Herre	..	—	—	—	X	—	—	—
32.	<i>M. (Mystus) micracanthus</i> (Bleeker)	—	—	—	X	X	—	—	—
33.	<i>M. (Mystus) baramensis</i> (Regan)	..	—	—	—	X	—	—	—
34.	<i>M. (Mystus) argentivittatus</i> (Regan)	—	—	—	—	X	—	—	—
35.	<i>M. (Mystus) henryi</i> (Herre)	..	—	—	—	X	—	—	—
36.	<i>M. (Mystus) cornula</i> (Chu)	—	—	—	—	X	—	—	—
37.	<i>M. (Mystus) hoi</i> (Pellegrin & Fang)	..	—	—	—	X	—	—	—

No.	Name of species.	1	2	3	4	5	6	7	8
38.	<i>M. (Mystus) pluriradiatus</i> (Vaillant)	—	—	—	—	—	—	X	—
39.	<i>M. (Mystus) rendahli</i> Pellegrin & Fang	—	—	—	—	—	—	X	—

To account for such an east to west dispersal, Hora (1937)¹ propounded the Satpura Hypothesis and since then it has been enlarged with interesting results. From the available Palaeontological evidence, it is known that the invasion of Siluroid fishes to India took place in the Pliocene period, which fact is evidenced by the presence of fishes like *M. (Osteobagrus) aor* in the Siwalik beds (Hora, 1952,² 1953³ for details). Peninsular India would seem to have received its fauna during the Pluvial periods of the Pleistocene Glaciation.⁴ The fauna that migrated along the Western Ghats was also aided into fast rate of speciation by the physiographic change & undergone by the Peninsula (Menon, 1952⁵). The presence of hill-stream inhabiting species like *M. (Mystus) punctatus* *M. (Mystus) montanus* would seem to be the resultants of such a physiographic change.

M. (Mystus) pelusius and *M. (Mystus) pelusius* var. *colwilli* which are related to *M. (Mystus) vittatus*, would appear to have spread along a cis-Himalayan watershed from an eastern place of origin.


¹ Hora, S. L., *Rec. Indian Mus.*, XXXIX, pp. 251-259 (1937).

² Hora, S. L., *Proc. Nat. Inst. Sci. India*, XVIII, (1952).

³ Hora, S. L., *Sci. Progress*, XLI, pp. 245-255 (1953).

⁴ Silas, E. G., *Proc. Nat. Inst. Sci. India*, XVIII, pp. 423-446 (1952)

⁵ Menon, A. G. K., *Ibid.*, XVII, pp. 475-497 (1951).



Mystus (Mystus) vittatus horai sub. sp. nov.

ON SOME VARIATIONS IN THE NEMATODE GENERA *PORROCAECUM* AND *ECHINOCEPHALUS*

By T. D. SOOTA, M.Sc., Zoological Assistant, Zoological Survey of India, Calcutta.

While working on an unnamed collection of nematodes in the Zoological Survey of India, which contained *Porrocaecum angusticolle* Baylis and Daubney, 1922 and *Echinocephalus spinosissimus* Baylis and Lane, 1920 the author noticed certain important variations in them. In his opinion they are not regarded significant enough for the present to treat them as new species. However, it is considered desirable to record the variations as below :—

Order : ASCAROIDEA Railliet and Henry, 1915.

Family : ASCARIDAE COBBOLD, 1864.

Sub-family : *ANISAKINAE* Railliet and Henry, 1912, emend. Baylis, 1920.

Genus : **Porrocaecum** Railliet and Henry, 1912.

1. **Porrocaecum angusticolle** Baylis and Daubney, 1922.

Male.—The male specimens measure 42-88 * in length and 0.61-0.95 in maximum width, in the posterior half of the body. The nerve ring is situated at a distance of 0.65-1.1 and the excretory pore 0.79-1.3 from the anterior end. The oesophagus with the oblong ventriculus, measures 3.59-5.6 in length. The ventriculus varies in length from 0.44-0.6 and 0.28-0.48 in width. The intestinal caecum is 2.3-4 in length. The tail measures 0.26-0.39 and the spicules 0.89-1.25 in length.

Female.—The female specimens vary in length from 57-108 with the maximum width of 1.04-1.4 in the posterior half of the body. Nerve ring is situated 0.9-1.21 from the anterior end and excretory pore at a distance of 1.05-1.44. Oesophagus with the oblong ventriculus, measures 4-6.4 in length. Ventriculus is 0.5-0.8 in length and 0.4-0.57 in width. Intestinal caecum varies in length from 2.3-3.7.

The vulva is situated at a distance of 21-44 from the anterior end. The eggs in the uterii measure 0.07-0.0812 in length and 0.0525-0.0696 in width. Anus lies 0.6-0.7 from the tip of the tail.

* All measurements are given in millimeters.

Reg. No. W.3807/1, Zoological Survey of India, Calcutta.

Host.—*Gyps bengalensis* Gmelin.

Locality —Hoshiarpur (Punjab).

Remarks.—The largest specimen of this species recorded by Baylis and Daubney (1922¹), measures only 55 in the male and 90 in the female while in a more recent record by Morgan and Schiller (1950²) from American Hawks the male is 78 and female 140 ; whereas the specimens the author has examined have attained a length of 88 in male and 108 in female indicating by far the greatest length for male. Besides this, the large number of preanal papillae in these specimens is also noteworthy. So far, the number recorded is not more than 20 pairs while the present author has counted 25-33 pairs in the males.

Order : FILARIOIDEA Weinland, 1858.

Family : GNATHOSTOMIDAE RAILLIET, 1895.

Sub-family : GNATHOSTOMINAE Baylis and Lane, 1920.

Genus : **Echinocephalus** Molin, 1858.

2. **Echinocephalus spinosissimus** Baylis and Lane, 1920.

Male.—The male specimen measures 58 in length and 1.04 in maximum thickness, in the middle of the body. Oesophagus is 6.7 long. Tail is 1.2 in length.

Female.—The female specimens measure upto 59 in length and 1.09 in maximum thickness. The oesophagus is 6.09 long. Tail is 1.25 in length.

Reg. No. W.3808/1, Zoological Survey of India, Calcutta.

Host.—Shark.

Locality.—Sandheads, mouth of the River Hooghly.

Remarks.—Out of 8 worms in this lot of collection only three, *i.e.*, one male and two females are in good state of preservation. These specimens differ from others so far recorded, in the great length attained by both the sexes. The largest male measures 58 and female 59 while in the earlier record by Baylis and Lane (1920³) it is 35.4 and 31 respectively.

The author is thankful to Dr. B. S. Chauhan, Assistant Superintendent for guidance and help during the course of this work.

¹ Baylis, H. A., *Fauna Brit. India, Ceylon and Burma*, Nematoda, I, pp. 71-73 (1936).

² Morgan, Banner Bill and Schiller, Everett L., *Trans. Amer. Micros. Soc.*, 69 (4), p. 371-372 (1950).

³ Baylis, H. A., *Fauna Brit. India, Ceylon and Burma*, Nematoda, II, pp. 193-195 (1939).